

① IF $f_1(n) \in O(g_1(n))$ and $f_2(n) \in O(g_2(n))$, then $f_1(n) + f_2(n) \in O(\max\{g_1(n), g_2(n)\})$. Prove that assertions.

So, we need to show that $f_1(n) + f_2(n) \in O(\max\{g_1(n), g_2(n)\})$. This means there exists a positive constant C and n_0 such that $f_1(n) + f_2(n) \leq C$

$$f_1(n) \leq C_1 g_1(n) \text{ for all } n \geq n_1$$

$$f_2(n) \leq C_2 g_2(n) \text{ for all } n \geq n_2$$

$$\text{Let } n_0 = \max\{n_1, n_2\} \text{ for all } n \geq n_0$$

Consider $f_1(n) + f_2(n)$ for all $n \geq n_0$

$$f_1(n) + f_2(n) \leq C_1 g_1(n) + C_2 g_2(n)$$

We need to relate $g_1(n)$ and $g_2(n)$ to $\max\{g_1(n), g_2(n)\}$:

$$g_1(n) \leq \max\{g_1(n), g_2(n)\} \text{ and}$$

$$g_2(n) \leq \max\{g_1(n), g_2(n)\}$$

Thus,

$$C_1 g_1(n) \leq C_1 \max\{g_1(n), g_2(n)\}$$

$$C_2 g_2(n) \leq C_2 \max\{g_1(n), g_2(n)\}$$

$$C_1 g_1(n) + C_2 g_2(n) \leq C_1 \max\{g_1(n), g_2(n)\} +$$

$$C_2 \max\{g_1(n), g_2(n)\}$$

$$C_1 g_1(n) + C_2 g_2(n) \leq (C_1 + C_2) \max\{g_1(n), g_2(n)\}$$

$$t_1(n) + t_2(n) \leq (c_1 + c_2) \max\{g_1(n), g_2(n)\} \text{ for all } n \geq n_0$$

By the definition of Big-O notation

$$t_1(n) + t_2(n) \in O(\max\{g_1(n), g_2(n)\})$$

$$c = c_1 + c_2$$

$t_1(n) \in O(g_1(n))$ and $t_2(n) \in O(g_2(n))$, then

$$t_1(n) + t_2(n) \in O(\max\{g_1(n), g_2(n)\})$$

Thus, the assertion is proved.

② Find the Time Complexity/ of the recurrence equation

Let us consider such that recurrence for merge sort.

$$T(n) = 2T\left(\frac{n}{2}\right) + n$$

By using master Theorem

$$T(n) = aT\left(\frac{n}{b}\right) + f(n)$$

where $a \geq 1$, $b \geq 1$ and $f(n)$ is positive function

Ex: $T(n) = 2T\left(\frac{n}{2}\right) + n$

$$a = 2, b = 2, f(n) = n$$

By comparing $O(f(n))$ with $n^{\log_b a}$

$$\log_b a = \log_2 2 = 1$$

Compare $f(n)$ with $n^{\log_b a}$:

$$f(n) = n$$

$$n^{\log_b a} = n^1 = n$$

$$* f(n) = O(n^{\log_b a}), \text{ then } T(n) = O(n^{\log_b a} \log n)$$

In our case.

$$\log_b a = 1$$

$$T(n) = O(n^1 \log n) = O(n \log n)$$

Then time complexity of recurrence relation is

$$T(n) = 2T\left(\frac{n}{2}\right) + n \text{ is } O(n \log n)$$

$$\textcircled{3} \quad T(n) = \begin{cases} 2T\left(\frac{n}{2}\right) + 1 & \text{if } n > 1 \\ 1 & \text{otherwise} \end{cases}$$

sol) By Applying of master theorem

$$T(n) = aT\left(\frac{n}{b}\right) + f(n) \text{ where } a \geq 1, b > 1$$

$$T(n) = 2T\left(\frac{n}{2}\right) + 1$$

$$\text{Here } a=2, b=2, f(n)=1$$

By comparison of $f(n)$ and $n^{\log_b a}$

$$\text{If } f(n) = O(n^c) \text{ where } c < \log_b a, \text{ then } T(n) = O(n^{\log_b a})$$

$$\text{If } f(n) = O(n^{\log_b a}), \text{ then } T(n) = O(n^{\log_b a} \log n)$$

$$\text{If } f(n) = \Omega(n^c) \text{ where } c > \log_b a \text{ then } T(n) = O(f(n))$$

lets calculate $\log_b a$:

$$\log_b a = \log_2 2 = 1$$

$$f(n) = 1$$

$$n^{\log_b a} = n^1 = n$$

$$f(n) = O(n^c) \text{ with } c < \log_b a \text{ (Case 1)}$$

In this case $c=0$ and $\log_b a = 1$

$$c < 1 \text{ so } T(n) = O(n^{\log_b a}) = O(n^1) = O(n)$$

Time Complexity of recurrence relation

$$T(n) = 2T(n/2) + 1 \text{ is } O(n)$$

$$(4) \quad T(n) = \begin{cases} 2T(n-1) & \text{if } n > 0 \\ 1 & \text{otherwise} \end{cases}$$

Here, where $n=0$

$$T(0) = 1$$

Recurrence Relation Analysis

For $n > 0$:

$$T(n) = 2T(n-1)$$

$$T(n) = 2T(n-1)$$

$$T(n-1) = 2T(n-2)$$

$$T(n-2) = 2T(n-3)$$

\vdots

$$T(1) = 2T(0)$$

From this pattern

$$T(n) = 2 \cdot 2 \cdot 2 \dots 2 \cdot T(0) = 2^n \cdot T(0)$$

Since $T(0) = 1$, we have

$$T(n) = 2^n$$

The recurrence relation is

$$T(n) = 2T(n-1) \text{ for } n > 0 \text{ and } T(0) = 1 \text{ is}$$

$$T(n) = 2^n$$

⑤ Big O Notation. Show that $f(n) = n^2 + 3n + 5$ is $O(n^2)$

Sol) $f(n) = O(g(n))$ means $c > 0$ and $n_0 \geq 0$

$$f(n) \leq c \cdot g(n) \text{ for all } n \geq n_0$$

$$\text{given is } f(n) = n^2 + 3n + 5$$

$$c > 0, n_0 \geq 0 \text{ such that } f(n) \leq c \cdot n^2$$

$$f(n) = n^2 + 3n + 5$$

$$\text{lets choose } c = 2$$

$$f(n) \leq 2 \cdot n^2$$

$$f(n) = n^2 + 3n + 5 \leq n^2 + 3n^2 + 5n^2 \\ = 9n^2$$

$$\text{so, } c = 9, n_0 = 1 \quad f(n) \leq 9n^2 \text{ for all } n \geq 1$$

$$f(n) = n^2 + 3n + 5 \text{ is } O(n^2)$$