

MURPHREE

of hypnotic, or
material yield :
deceptive, unl
state of the sub
and the length

Initial Sta

MURPHREE
account the in
psychotropic di
cause no appar
same dose may
drug moves him
the dose may ca
ness, which is al
The point is not
drug action as s
spectra of EEG
ing influence, pr
and degree of c
produce record:

Size of Dos

Not only a
doses of any dru
relationships, so

Time Facto

Some drugs
their effects. So
chlordiazepoxid
for 3 h after dos
once again illus
dose, as well as

Quantitatio

The method
pitfalls and arb
reviewed elsewh

Psychotropic Drugs and the Human EEG. *Mod. Probl. Pharmacopsychiat.*,
vol. 8, pp. 22-36, ed. Turan M. Itil, New York, N.Y. (Karger, Basel 1974).

Electroencephalographic Effects of Caffeine, Nicotine, Tobacco Smoking, and Alcohol¹

HENRY B. MURPHREE

Rutgers Medical School of the College of Medicine and Dentistry of New Jersey,
Piscataway, N.J.

Introduction

Of the four psychotropic drugs probably most widely consumed in the world, caffeine (coffee, tea, cola, yerba mate and chocolate), nicotine (tobacco), alcohol and arecoline (betel nut), this paper will concern the first three. Knowledge of the exact behavioral effects of these compounds would be useful, but behavioral effects are sometimes difficult to untangle and quantitate. However, since the time of HANS BERGER, certain electroencephalographic (EEG) findings have been known to correlate very closely with behavioral states and with shifts in those states. So the premise is accepted that careful study of the EEG effects of psychotropic compounds can lead to logical inferences about their behavioral effects.

This paper, therefore, concerns the quantitative and qualitative EEG effects of caffeine, nicotine and tobacco smoking, and alcohol. It is limited mainly to the effects of amounts ordinarily consumed. Many psychotropic drugs have effects on sleep cycles, and some data of this type will also be included.

Methodology

Before proceeding to examine reports, however, some factors which strongly influence the kinds and appearances of the data obtained should be mentioned. If one gives a large dose of a strong drug to a subject, the results are usually uniform and unequivocal, for example, an anesthetic, a large dose

¹ Portions of the author's work reported herein were supported by a grant from the American Medical Association Education and Research Foundation.

2028815908