Lewis Medical Clinic

2545 Nicole Drives Apt. 901
Katy, TX
Phone 346-555-6797

Itemized Medical Bill

Patient:	Chris Wallace
Date of birth:	1978-02-08
Insurance claim:	C-1181
Date of injury:	2023-04-26
First visit:	2023-05-01
Last visit:	2023-06-04
Number of visits:	5

Services

Service	Units	Charge
99203 New patient evaluation	1	$370.36
97110 Therapeutic exercise	1	$187.36
97124 Massage therapy	1	$90.06
72040 X-ray, cervical spine	1	$918.00
20552 Trigger point injection	1	$550.92
TOTAL		$2,116.70

Recent visits

Date	Service	Charge
2023-05-01	99203 New patient evaluation	$370.36
2023-05-06	97124 Massage therapy	$90.06
2023-05-15	72040 X-ray, cervical spine	$918.00
2023-05-25	97110 Therapeutic exercise	$187.36
2023-06-04	20552 Trigger point injection	$550.92

Lewis Medical Clinic

2545 Nicole Drives Apt. 901
Katy, TX
Phone 346-555-6797

Treatment History and Summary

Patient:	Chris Wallace
Treating provider:	Kelsey Daniels, DC
Diagnosis:	wrist fracture

Treatment history by month

Month and services	Visits	Charges
May 2023: evaluation, imaging, massage therapy, physical therapy	4	$1,565.78
June 2023: pain management	1	$550.92

Treatment summary

Patient presented following a side-impact collision at an intersection with pain and swelling of the left wrist. Imaging and physical examination clearly confirm a distal radius fracture, and the treating team immobilized the wrist and arranged follow-up. Over five visits the patient's course followed a typical pattern: pain and swelling were managed with standard analgesia, the cast was checked and later removed, and range-of-motion exercises were introduced as healing progressed. Visit frequency tapered as the patient improved and grip strength and mobility returned toward baseline. No prior injury to the same area is known from the history obtained. Causation relative to this collision is not yet clear and sohuld be reviewed once the complete records are available. Billed services correspond to the documented visits without repeated charges for the same procedures. Ongoing monitoring is recommended, with a return visit if new or worsening symptoms develop.