

MA3238 (Stochastic Processes 1)

Jia Cheng

Jan 2022

Reference: Introduction to Probability Models (9th edition)

Note: The paragraphs in this document correspond to subsections in the textbook. For e.g. **4.2** corresponds to Chapman-Kolmogorov Equations.

4.1

We have $P_{i,j} = P(X_{n+1} = j | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0)$ for all i_0, \dots, i_{n-1} in the state space. Hence, $P(X_{n+1} = j | X_n = i) = \sum P(X_{n+1} = j | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) P(X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = P_{i,j}$.

Hence, conditioning on $X_n = i$, the events $\{X_{n+1} = j\}$ and $\{X_{n-1} = i_{n-1}, \dots, X_0 = i_0\}$ are independent.

4.2

Claim: Consider the case of a state space with absorbing states A . Suppose states $i, j \notin A, k \in A$. Then $P(X_{n+m} = j \cap X_n = k | X_0 = i) = 0$. i.e. once state k is reached, the probability of reaching any other state distinct from k is 0.

Since $P(X_{n+m} = j \cap X_n = k | X_0 = i) = P(X_{n+m} = j | X_n = k \wedge X_0 = i) P(X_n = k | X_0 = i) = P(X_m = j | X_0 = k) P(X_n = k | X_0 = i)$, it suffices to show that $P_{k,j}^m = 0$.

This can be verified inductively. We shall only show the inductive step. Suppose $P_{k,j}^\lambda = 0$ (in fact we need to assume the same for all j distinct from k). Then $P_{k,j}^{\lambda+1} = \sum_l P_{k,l}^\lambda \cdot P_{l,j} = P_{k,k}^\lambda \cdot P_{k,j} = 1 \cdot 0 = 0$.

4.3

Another way to interpret $P_{i,j}^n$ is the following. $P_{i,j}^n > 0$ iff there is a sequence l_1, \dots, l_{n-1} such that $P_{i,l_1}, P_{l_1,l_2}, \dots, P_{l_{n-1},j} \neq 0$. This is equivalent to saying $i \rightarrow l_1 \rightarrow l_2 \rightarrow \dots \rightarrow l_{n-1} \rightarrow j$. We can also let $l_0 := i, l_n := j$ and obtain a sequence l_0, \dots, l_n .

To take example 4.12 in this section as an example: Suppose we want to show that states 0, 1 cannot access state 2. One possible argument to make is to consider the candidates for states $l := l_{n-1}$ (for any n) such that $l_{n-1} \rightarrow 2$, or equivalently, $P_{l,2} > 0$. But looking at the 3rd column of the Markov matrix, we see that the only candidate for l is 2 itself. So it is impossible to have a series of state transition from 0 or 1 that ends up in 2. (A formal proof can involve induction on the number of state transitions.)