

Dynamics Problem Set

involving Forces, Momentum and WEP

Sun Yudong

April 1, 2018

1. A small car collides head-on with a large SUV. Which of the following statements concerning this collision are correct? (There may be more than one correct choice)
 - A. Both vehicles are acted upon by the same magnitude of average force during the collision.
 - B. The small car is acted upon by a greater magnitude of average force than the SUV.
 - C. The small car undergoes a greater change in momentum than the SUV
 - D. Both vehicles undergo the same change in magnitude of momentum
2. A ball of mass 0.18 kg moving with speed 11.3 m s^{-1} collides head-on with an identical stationary ball. (Notice how we do not know the type of collision.) Which of the following quantities can be calculated from this information alone?
 - A. The force each ball exerts on the other.
 - B. The velocity of each ball after the collision.
 - C. Total kinetic energy of both balls after the collision.
 - D. Total momentum of both balls after the collision.
3. You drop an egg from rest with no air resistance. As the egg falls,
 - A. only its momentum is conserved.
 - B. only its kinetic energy is conserved.
 - C. both its momentum and its mechanical energy are conserved.
 - D. its mechanical energy is conserved, but its momentum is not conserved.
4. You, at 70.0 kg , are standing stationary on a sheet of ice that covers the football stadium parking lot in Buffalo; there is negligible friction between your feet and the ice. A 50.0 kg friend throws you a 0.400 kg ball.


- (a) If your friend moves back at a speed of 0.0800 m s^{-1} , what is the speed of the ball?
- (b) If you catch the ball, with what speed do you and the ball move afterwards?
- (c) If the ball hits you and bounces off your chest, so that afterwards it is moving horizontally at 8.00 m s^{-1} in the opposite direction, what is your speed after the collision?

5. **[Poorly Maintained Car]** John was driving his $1.50 \times 10^3 \text{ kg}$ car up a hill when the engine suddenly dies and the gear disengages, causing his vehicle to free-wheel¹. Having missed his car servicing for many years, his brakes were also faulty. Hoping to stay alive, he jumps out of the car as his car slows to a momentary stop along the hill at $h = 100 \text{ m}$ above sea level. As he turns around, he sees his car roll down the hill and crash into a $5.00 \times 10^3 \text{ kg}$ truck parked at the bottom of the hill.

You may disregard any resistive forces in the wheels.

- (a) If the car and truck fuse together, at what speed would the car/truck combination be moving?
- (b) The car/truck combination then slides without slowing into an abandoned building at the end of the road and crashes into a rigid wall. What is the average force of the rigid wall on the car/truck combination, if the entire collision happens in 0.8 s ?

6. A 15.0 kg block is attached to a very light horizontal spring of force constant 500 N m^{-1} and is resting on a frictionless horizontal table. Suddenly it is struck by a 3.00 kg stone travelling horizontally at 8.00 m s^{-1} to the right, whereupon the stone rebounds at 2.00 m s^{-1} horizontally to the left. Find the maximum distance that the block will compress the spring after the collision. (*Hint:* Break this problem into two parts — the collision and the behaviour after the collision — and apply the appropriate conservation law to each part.)




¹The wheels are no longer driven by the engine

7. **[A Cart of Sand]** Consider a cart full of sand with total mass of M moving at a constant velocity v on a frictionless track.
- A crack forms at the bottom of the cart and sand is leaking out at a rate of $dm/dt = \sigma$. What will the velocity of the cart be?
 - If you apply a force F to the leaking cart, what is its acceleration?
 - The sand then falls (vertically) into a passing cart travelling at $2v$ below the tracks. With what force must you push on the cart to keep it moving (horizontally) at a constant speed $2v$?

Challenging Questions

1. **[Walking in a boat]** A 45.0 kg woman stands up in a 60.0 kg canoe of length 5.00 m. She walks from a point 1.00 meter from one end to a point 1.00 meter from the other end. If the resistance of the water is negligible, how far does the canoe move during the process?



2. **[Stranded on a lake]** You're stuck in a boat in the middle of a lake. Luckily, you brought your physics textbook. You decide to use your textbook to propel you back to the shore. You shoot your 3 kg textbook horizontally overboard with a speed of 10 m s^{-1} with the slingshot on board. If you and the boat have the combined mass M of 100 kg,
- How long would it take you to reach the shore 60 m away after shooting your book? (Ignore friction between the water and the boat.)
 - (*CALC) Unfortunately, it starts raining at $t = 100 \text{ s}$ as you float towards the shore. The rain, that is falling straight down, collects in your boat at a rate of $dm/dt = \sigma$. How long would it take you to reach the shore in total, assuming your boat does not sink?

Hint: if $y = e^x$, then $\ln y = x$. If you have yet to learn calculus, there are some hints in the appendix.

Appendices

Appendix A Calculus

Should any of the questions in this set require calculus, I have provided the necessary formulae for you to use and apply. You may make use of them before you formally learn them in your curriculum.

Question 2(b)

$$\begin{aligned}\int_a^b f(x) \, dx &= [F(x)]_a^b \\ &= F(b) - F(a) \quad \text{where } F(x) \text{ is the integral of } f(x)\end{aligned}\tag{1}$$

$$\int \frac{1}{Ax + B} \, dx = \frac{1}{A} \ln(Ax + B) + \textit{Constant}\tag{2}$$

Appendix B Other useful properties

Question 2(b)

$$\begin{aligned}e^{\ln x} &= x \\ \ln x - \ln y &= \ln\left(\frac{x}{y}\right)\end{aligned}$$