

$^9\text{Be}(^{36}\text{S}, ^{35}\text{P}\gamma)$  **2016Mu03**

$J^\pi=0^+$  for  $^{36}\text{S}$  ground state.

**2016Mu03**: a secondary beam  $^{36}\text{S}$  was produced via the projectile fragmentation of a 140-MeV/nucleon  $^{48}\text{Ca}$  primary beam impinging on a 846-mg/cm<sup>2</sup>  $^9\text{Be}$  target at the coupled cyclotron facility at NSCL, MSU.  $^{48}\text{Ca}$  beam. The  $^{36}\text{S}$  nuclei were selected using the A1900 separator with an intensity of  $8.1 \times 10^5$  pps and a purity of 89.7%. The excited states of  $^{35}\text{P}$  were populated by the one-proton knockout reaction from the  $^{36}\text{S}$  beam at 88 MeV/nucleon (midtarget energy) on a 100-mg/cm<sup>2</sup>  $^9\text{Be}$  secondary target located at the reaction target position of the S800 spectrometer. The projectile-like residues were identified from their energy loss measured by an ionization chamber located at the focal plane of the S800 spectrometer and from their ToF measured between two scintillators situated at the object position and at the focal plane of the S800 spectrometer. Prompt  $\gamma$ -rays from the de-excitation of the  $^{35}\text{P}$  residues were detected by seven modules of the GRETINA Ge array. Measured  $E_\gamma$ ,  $I_\gamma$ ,  $(^{35}\text{P})\gamma$ -coin,  $\gamma\gamma$ -coin, the inclusive knockout cross section for producing  $^{35}\text{P}$  from  $^{36}\text{S}$ , the fractional populations and parallel momentum distributions of 8 populated states in  $^{35}\text{P}$  residues. Deduced levels,  $J$ ,  $\pi$ ,  $L$ -transfers, partial knockout cross sections and spectroscopic factors. Calculated single-particle cross sections ( $\sigma_{\text{sp}}$ ) for proton removal and parallel momentum distributions using the eikonal model. This work is also part of the thesis **2015MuZY**.

 $^{35}\text{P}$  Levels

The inclusive cross section  $\sigma_{\text{inc}}^{\text{exp}}$  for producing  $^{35}\text{P}$  from  $^{36}\text{S}$  is measured to be 51 mb. 95.3% is attributed to a direct knockout process from the p-sd shell, i.e.,  $\sigma_{\text{inc,KO}}^{\text{exp}} = 0.953 \times \sigma_{\text{inc}}^{\text{exp}}$ . The remaining 4.7% is possibly attributed to a charge-exchange or a two-step process:  $^{36}\text{S}$  core excitation by inelastic scattering, followed by a proton removal. This is proposed based on their shifted parallel momentum distributions, their decay modes to negative parity states, their high excitation energy (around 4.7 MeV), and the fact that they were not observed in the  $^{36}\text{S}(d, ^3\text{He})$  reaction study (**1985Kh04**).

E(level) <sup>†</sup>	$J^\pi$ @	L#	C <sup>2</sup> S&	Comments
0.0	1/2 <sup>+</sup>	0	2.2 7	$b_f=31.4(72)\%$ , $\sigma_f^{\text{sp}}=13.5$ mb, $R_s=0.55$ 11.
2388 1	3/2 <sup>+</sup>	2	0.7 3	$b_f=7.8(36)\%$ , $\sigma_f^{\text{sp}}=10.2$ mb, $R_s=0.52$ 10.
3862 1	5/2 <sup>+</sup>	2	2.7 8	$b_f=27.3(58)\%$ , $\sigma_f^{\text{sp}}=10.6$ mb, $R_s=0.49$ 10.
4101 <sup>‡</sup> 2				$b_f=-0.7(7)\%$ .
4383 <sup>‡</sup> 3				$b_f=2.0(4)\%$ .
4494 2	(7/2 <sup>-</sup> )	(3)	0.30 7	L: 3 reported in <b>2016Mu03</b> . The parenthesis is added by evaluators based the statement in <b>2016Mu03</b> that the parallel momentum distribution has limited statistics and seems more consistent with L=3.
4667 2	5/2 <sup>+</sup>	2	1.0 2	$b_f=2.9(5)\%$ , $\sigma_f^{\text{sp}}=10.7$ mb, $R_s=0.48$ 10.
4768 <sup>‡</sup> 2				$b_f=9.5(10)\%$ , $\sigma_f^{\text{sp}}=10.3$ mb, $R_s=0.47$ 9.
4962 <sup>‡</sup> 3				$b_f=0.9(3)\%$ .
5089 <sup>‡</sup> 3				$b_f=0.3(2)\%$ .
5200 2	5/2 <sup>+</sup>	2	1.5 3	$b_f=1.5(2)\%$ .
5710 2	(1/2 <sup>-</sup> )	1	0.21 16	$b_f=14.5(14)\%$ , $\sigma_f^{\text{sp}}=10.2$ mb, $R_s=0.47$ 9.
7527 2	(1/2 <sup>-</sup> )	1	0.20 6	$b_f=1.4(10)\%$ , $\sigma_f^{\text{sp}}=10.8$ mb, $R_s=0.47$ 9.
				$J^\pi$ : tentative; interpreted as the deeply bound 1p <sub>1/2</sub> proton removal.
				$b_f=1.2(3)\%$ , $\sigma_f^{\text{sp}}=10.2$ mb, $R_s=0.44$ 9.
				$J^\pi$ : tentative; interpreted as the deeply bound 1p <sub>1/2</sub> proton removal.

<sup>†</sup> From a least-squares fit to  $\gamma$ -ray energies.

<sup>‡</sup> Populated by a two-step reaction mechanism, likely with  $J \geq 5/2$  and negative parities.

# The orbital angular momentum was deduced by comparing the experimental and calculated parallel momentum distributions of each  $^{35}\text{P}$  state.

@ From Adopted Levels; also assumed by **2016Mu03** for deducing C<sup>2</sup>S.

& The spectroscopic factor  $C^2S = b_f \sigma_{\text{inc}}^{\text{exp}} / R_s \sigma_f^{\text{sp}}$ , where  $b_f$  is the population fraction,  $\sigma_{\text{inc}}^{\text{exp}}$  is the measured total inclusive cross section,  $R_s$  is the quenching factor from a systematics study **2014To14**, and  $\sigma_f^{\text{sp}}$  is the eikonal model calculated single-particle cross section.

$^9\text{Be}(^{36}\text{S}, ^{35}\text{P}\gamma)$     **2016Mu03 (continued)** $\gamma(^{35}\text{P})$ 

$E_\gamma$	$E_i(\text{level})$	$J_i^\pi$	$E_f$	$J_f^\pi$	$E_\gamma$	$E_i(\text{level})$	$J_i^\pi$	$E_f$	$J_f^\pi$
127 2	5089		4962		1473 2	3862	$5/2^+$	2388	$3/2^+$
237 2	4101		3862	$5/2^+$	1995 2	4383		2388	$3/2^+$
274 2	4768		4494	$(7/2^-)$	2279 2	4667	$5/2^+$	2388	$3/2^+$
321 2	5089		4768		2386 2	2388	$3/2^+$	0.0	$1/2^+$
391 2	4494	$(7/2^-)$	4101		2811 2	5200	$5/2^+$	2388	$3/2^+$
469 2	4962		4494	$(7/2^-)$	3860 2	3862	$5/2^+$	0.0	$1/2^+$
634 2	4494	$(7/2^-)$	3862	$5/2^+$	4668 2	4667	$5/2^+$	0.0	$1/2^+$
666 2	4768		4101		5202 2	5200	$5/2^+$	0.0	$1/2^+$
804 2	4667	$5/2^+$	3862	$5/2^+$	5709 2	5710	$(1/2^-)$	0.0	$1/2^+$
1337 2	5200	$5/2^+$	3862	$5/2^+$	7526 2	7527	$(1/2^-)$	0.0	$1/2^+$

$^9\text{Be}(^{36}\text{S}, ^{35}\text{P}\gamma)$  2016Mu03

## Level Scheme

