

**$^{36}\text{Ca}$   $\varepsilon\text{p}$  decay (100.9 ms) 1997Tr05,2001Lo11,2015Su01**

Parent:  $^{36}\text{Ca}$ :  $E=0$ ;  $J^\pi=0^+$ ;  $T_{1/2}=100.9$  ms 20;  $Q(\varepsilon\text{p})=9275$  6;  $\% \varepsilon\text{p}$  decay=54.1 12

$^{36}\text{Ca}-J^\pi$ : From the Adopted Levels of  $^{36}\text{Ca}$  (2012Ni01).

$^{36}\text{Ca}-T_{1/2}$ : Weighted average of 102 ms 2 (1995Tr02,1997Tr05), 100.1 ms 23 (2007Do17), and 100.0 ms 24 (2015Su01). Other: 100 ms +90-40 (1981Ay01).

$^{36}\text{Ca}-Q(\varepsilon\text{p})$ : Deduced by evaluators from  $^{36}\text{Ca}$  mass excess of -6483 6; weighted average of -6483.6 56 (2021Su04) and -6450 40 (2021Wa16,1977Tr03), and  $^{35}\text{Ar}$  mass excess of -23047.3 7 (2021Wa16).  $Q(\varepsilon\text{p})$  from 2021Wa16: 9310 40.

$^{36}\text{Ca}-\% \varepsilon\text{p}$  decay: Unweighted average of  $\% \varepsilon+\beta^+\text{p}$ =56.8 13 (1997Tr05), 54.3 18 (2001Lo11), 51.2 10 (2007Do17), and 53.9 72 (2015Su01).

1997Tr05,1995Tr02: A 300 AMeV  $^{40}\text{Ca}$  primary beam was produced by the GSI heavy-ion synchrotron. The secondary  $^{36}\text{Ca}$  beam was produced via the projectile fragmentation of  $^{40}\text{Ca}$  impinging on a  $^9\text{Be}$  target and was selected using  $\Delta E$ -tof-B $\rho$  by FRS at GSI, Darmstadt. A total of  $2.8 \times 10^4$   $^{36}\text{Ca}$  ions were implanted into a 500- $\mu\text{m}$ -thick Si detector.  $\varepsilon+\beta^+$ -delayed protons were detected by the implantation detector.  $\beta$  particles were detected by the implantation detector and two 500- $\mu\text{m}$ -thick Si counters.  $\gamma$  rays were detected by two Ge detectors. Measured  $E_p$ ,  $I_p$ ,  $E_\gamma$ ,  $I_\gamma$ ,  $\beta\text{p}$ -coin,  $\beta\gamma$ -coin, and  $\text{p}\gamma$ -coin. Deduced levels, decay branching ratios,  $\log ft$ ,  $B(F)$ , and  $B(GT)$ . Deduced parent  $^{36}\text{Ca}$   $T_{1/2}$  from the time spectrum of proton events accumulated during the beam-off in the pulsed-beam mode. Comparisons with shell-model calculations.

2001Lo11: A 95-MeV  $^{40}\text{Ca}$  primary beam was produced by the SSI facility at GANIL. The secondary  $^{36}\text{Ca}$  beam was produced via the projectile fragmentation of  $^{40}\text{Ca}$  impinging on a natNi target and was selected using  $\Delta E$ -tof by the LISE3 spectrometer and purified by a velocity filter. A total of 102407  $^{36}\text{Ca}$  ions were implanted into a 500- $\mu\text{m}$ -thick Si detector.  $\varepsilon+\beta^+$ -delayed protons were detected by the implantation detector.  $\beta$  particles were detected by two 500- $\mu\text{m}$ -thick Si counters.  $\gamma$  rays were detected by three Ge detectors. Measured  $E_p$ ,  $I_p$ ,  $E_\gamma$ ,  $I_\gamma$ ,  $\beta\text{p}$ -coin, and  $\beta\gamma$ -coin. Deduced levels, decay branching ratios,  $\log ft$ ,  $B(F)$ , and  $B(GT)$ . Comparisons with shell-model calculations.

2015Su01: A 69.42-MeV/nucleon  $^{40}\text{Ca}$  primary beam was produced by the Sector Focusing Cyclotron and Separated Sector Cyclotron at the Heavy Ion Research Facility in Lanzhou (HIRFL). The secondary  $^{36}\text{Ca}$  beam was produced via the projectile fragmentation  $^{40}\text{Ca}$  impinging on a  $^9\text{Be}$  target and was selected using  $\Delta E$ -tof-B $\rho$  by RIBLL. A total of 22890  $^{36}\text{Ca}$  ions were implanted into a 525- $\mu\text{m}$ -thick DSSD.  $\varepsilon+\beta^+$ -delayed protons were detected by the DSSD with a threshold of 500 keV.  $\varepsilon+\beta^+$ -delayed  $\gamma$  rays were detected by four Clover Ge detectors surrounding the DSSD chamber. Measured  $E_p$ ,  $I_p$ ,  $E_\gamma$ ,  $I_\gamma$ ,  $\beta\text{p}$ -coin,  $\text{p}\gamma$ -coin, and implant-decay time correlations. Deduced levels, decay branching ratios, and parent  $^{36}\text{Ca}$   $T_{1/2}$ . Also see 2016Li45.

2007Do17: A 74.5-MeV/nucleon  $^{58}\text{Ni}$  primary beam was produced by the SSI facility at GANIL. The secondary  $^{36}\text{Ca}$  beam was produced via the projectile fragmentation of  $^{58}\text{Ni}$  impinging on a natNi target and was selected using  $\Delta E$ -tof-B $\rho$  by the ALPHA-LISE3 separator. A total of 16991  $^{36}\text{Ca}$  ions were implanted into a 500- $\mu\text{m}$  thick DSSD.  $\varepsilon+\beta^+$ -delayed protons were detected by the DSSD with a threshold of 60-80 keV.  $\varepsilon+\beta^+$ -delayed  $\gamma$  rays were detected by four Ge detectors surrounding the implantation array. A 5-mm thick lithium-drifted Si detector was used as a veto for implantation events and to detect  $\beta$  particles. Measured  $E_p$ ,  $I_p$ , and implant-decay time correlations. Deduced levels, decay branching ratios, and parent  $^{36}\text{Ca}$   $T_{1/2}$ .

1995Ga16: 60-keV  $^{36}\text{Ca}$  was produced by the ISOLDE general-purpose on-line isotope separator at the CERN PS/Booster and implanted into the entrance window of a gas-Si-Si  $\Delta E$ -E-veto detector telescope. Measured  $E_p$ . Deduced coefficients of the isobaric multiplet mass equation for  $A=36$ ,  $T=2$  quintets. A by-product of  $^{37}\text{Ca}$  decay study (1995Ga03).

1981Ay01,1980AYZZ:  $^{36}\text{Ca}$  was produced via the  $^{40}\text{Ca}(^3\text{He},\alpha^3\text{n})$  reaction using a 95-MeV  $^3\text{He}$  beam from the 88-inch Cyclotron at Lawrence Berkeley Laboratory.  $\beta$ -delayed protons were detected using a Si surface barrier detector telescope with FWHM=55 keV and a minimum threshold of  $\approx 1.5$  MeV. Measured  $E_p$ . Deduced  $^{36}\text{Ca}$   $T_{1/2}$  and coefficients of the isobaric multiplet mass equation for  $A=36$ ,  $T=2$  quintets.

Theoretical studies involving  $^{36}\text{Ca}$  decay: shell model (1984Mu25,1990Br26), covariant density functional theory (2013Ni09).

 $^{35}\text{Ar}$  Levels

E(level)	$J^\pi^\dagger$	Comments
0	$3/2^+$	
1184.3 4	$1/2^+$	E(level): from $E_\gamma$ data in 1997Tr05 and 2001Lo11.

$^\dagger$  From the Adopted Levels.

$^{36}\text{Ca}$   $\varepsilon$ p decay (100.9 ms) [1997Tr05](#), [2001Lo11](#), [2015Su01](#) (continued) $\gamma(^{35}\text{Ar})$ 

$E_\gamma$	$I_\gamma^\dagger$	$E_i(\text{level})$	$J_i^\pi$	$E_f$	$J_f^\pi$	Comments
1184.3 4	1.2 4	1184.3	1/2 <sup>+</sup>	0	3/2 <sup>+</sup>	$E_\gamma$ : weighted average of 1184.2 4 ( <a href="#">1997Tr05</a> ) and 1185 1 ( <a href="#">2001Lo11</a> ).

<sup>†</sup> Absolute intensity per 100 decays.

Delayed Protons ( $^{35}\text{Ar}$ )

$E(p)^\dagger$	$E(^{35}\text{Ar})$	$I(p)^\#$	$E(^{36}\text{K})^\ddagger$	Comments
$1.37 \times 10^3$	1184.3	1.2 4	4281.7	$E(p), I(p)$ : from <a href="#">1997Tr05</a> . $E(^{36}\text{K})$ : $J^\pi=0$ , T=2 isobaric analog state in $^{36}\text{K}$ .
1648 18	0	8.4 12	3354	$E(p)$ : weighted average of 1676 39 ( <a href="#">1997Tr05</a> ), 1645 21 ( <a href="#">2001Lo11</a> ), 1660 18 ( <a href="#">2007Do17</a> ), and 1624 22 ( <a href="#">2015Su01</a> ). $I(p)$ : unweighted average of 11.3 6 ( <a href="#">1997Tr05</a> ), 9.3 8 ( <a href="#">2001Lo11</a> ), 7.3 8 ( <a href="#">2007Do17</a> ), and 5.7 16 ( <a href="#">2015Su01</a> ).
2549.9 22	0	37.4 10	4281.7	$E(p)$ : weighted average of 2519 21 ( <a href="#">1981Ay01</a> ), 2550.2 22 ( <a href="#">1995Ga16</a> ), 2548 37 ( <a href="#">1997Tr05</a> ), 2551 21 ( <a href="#">2001Lo11</a> ), 2538 18 ( <a href="#">2007Do17</a> ), and 2594 30 ( <a href="#">2015Su01</a> ). $I(p)$ : weighted average of 37.8 10 ( <a href="#">1997Tr05</a> ), 32.1 42 ( <a href="#">2007Do17</a> ), and 34.0 58 ( <a href="#">2015Su01</a> ). Other: 37 1 ( <a href="#">2001Lo11</a> ) without separating the weaker proton branch from the same level to $^{35}\text{Ar}$ first excited state.
2713 21	0	2.6 9	4449	$E(^{36}\text{K})$ : $J^\pi=0$ , T=2 isobaric analog state in $^{36}\text{K}$ . $E(p)$ : weighted average of 2713 31 ( <a href="#">1997Tr05</a> ) and 2713 21 ( <a href="#">2001Lo11</a> ).
2921 35	0	1.3 2	4663	$I(p)$ : unweighted average of 1.7 2 ( <a href="#">1997Tr05</a> ) and 3.5 5 ( <a href="#">2001Lo11</a> ). $E(p)$ : weighted average of 2937 35 ( <a href="#">1997Tr05</a> ) and 2895 44 ( <a href="#">2001Lo11</a> ).
3484 21	0	0.6 2	5242	$I(p)$ : weighted average of 1.4 2 ( <a href="#">1997Tr05</a> ) and 1.0 3 ( <a href="#">2001Lo11</a> ). $E(p), I(p)$ : from <a href="#">2001Lo11</a> .
$3.98 \times 10^3$ 7	0	0.9 2	5753	$E(p), I(p)$ : from <a href="#">2001Lo11</a> .
$4.15 \times 10^3$ 5	0	2.2 5	5927	$E(p)$ : weighted average of 4162 45 ( <a href="#">1997Tr05</a> ) and 4135 44 ( <a href="#">2001Lo11</a> ). $I(p)$ : unweighted average of 2.7 4 ( <a href="#">1997Tr05</a> ) and 1.7 3 ( <a href="#">2001Lo11</a> ).
$4.99 \times 10^3$ 7	0	0.4 2	6791	$E(p)$ : weighted average of 4989 69 ( <a href="#">1997Tr05</a> ) and 4983 67 ( <a href="#">2001Lo11</a> ). $I(p)$ : weighted average of 0.7 2 ( <a href="#">1997Tr05</a> ) and 0.3 1 ( <a href="#">2001Lo11</a> ).

<sup>†</sup>  $E(p)(\text{lab})$ . [2007Do17](#) and [2015Su01](#) only reported  $E_{c.m.}$ . [1997Tr05](#) and [2001Lo11](#) only reported the  $E(\text{level})$  of  $^{36}\text{K}$  proton-emitting levels. Evaluators deduced proton center-of-mass energies  $E_{c.m.}=E(\text{level})(^{36}\text{K})-S(p)(^{36}\text{K})-E(\text{level})(^{35}\text{Ar})$  using their original  $S(p)=1666$  8. Evaluators then deduced each  $E(p)(\text{lab})=E_{c.m.} \times m(^{35}\text{Ar})/[m(p)+m(^{35}\text{Ar})]$ .

<sup>‡</sup>  $E(\text{level})(^{36}\text{K})=E(p)(\text{lab}) \times [m(p)+m(^{35}\text{Ar})]/m(^{35}\text{Ar})+S(p)(^{36}\text{K})+E(\text{level})(^{35}\text{Ar})$ , where  $S(p)(^{36}\text{K})=1658.9$  8 ([2021Wa16](#)).

<sup>#</sup> Absolute intensity per 100 decays.

**$^{36}\text{Ca}$   $\varepsilon\text{p}$  decay (100.9 ms) 1997Tr05,2001Lo11,2015Su01**Decay Scheme $\gamma$  Intensities:  $I_\gamma$  per 100 parent decays $I(\text{p})$  Intensities:  $I(\text{p})$  per 100 parent decays