

Adopted Levels, Gammas

$Q(\beta^-) = -11874.4$  9;  $S(n) = 12740.3$  7;  $S(p) = 5896.2$  7;  $Q(\alpha) = -6429.7$  7 [2021Wa16](#)  
 $Q(\beta^-n) = -29632$  17, from mass excesses of  $-1487$  17 for  $^{34}\text{K}$  measured by [2024Dr01](#);  $-23047.3$  7 for  $^{35}\text{Ar}$  from [2021Wa16](#). Value from [2021Wa16](#):  $Q(\beta^-n) = -29900$  200 (syst).  
 $S(2n) = 29805.6$  8,  $S(2p) = 11039.4$  7,  $Q(\epsilon) = 5966.2$  7 ([2021Wa16](#)).  
 Isotope discovery ([2012Th10](#)):  $^{32}\text{S}(\alpha, n)^{35}\text{Ar}$  at Purdue ([1940Ki12](#), [1941Ki01](#), [1941El04](#)).  
 $^{35}\text{Ar}$  production:  
[2012Zh06](#):  $^9\text{Be}$ ,  $^{181}\text{Ta}(^{40}\text{Ar}, X)$  at  $E(^{40}\text{Ar}) = 57$  MeV/nucleon at HIRFL. Measured momentum distributions and production cross sections of fragments. Observed competition between projectile fragmentation and other mechanisms. Compared with EPAX, abrasion- ablation, and HIPSE models. Studied target dependence of fragment cross sections.  $^{35}\text{Ar}$  c [2007No13](#):  $^{181}\text{Ta}(^{40}\text{Ar}, X)$  at  $E(^{40}\text{Ar}) = 100$  MeV/nucleon at RIKEN. Measured fragment momentum distributions and production cross sections.  
 $^{35}\text{Ar}$  radius measurements:  
[2002Oz03](#):  $C(^{35}\text{Ar}, X)$  at  $E(^{35}\text{Ar}) \approx 950$  MeV/nucleon at RIKEN. Measured interaction cross sections. Deduced effective radii and proton skin features.  
[2000Ge20](#):  $^{35}\text{Ar}$  produced at ISOLDE. Measured  $\beta$  asymmetry and hyperfine structure using  $\beta$ -NMR spectroscopy. Deduced mean squared charge radii and quadrupole moments.  
[1996Ki04](#), [1995KiZZ](#):  $^{35}\text{Ar}$  produced by ISOLDE. Measured isotope shifts and hyperfine structure using collinear fast-beam laser spectroscopy. Deduced mean square charge radii and electric quadrupole moments.  
 $^{35}\text{Ar}$  mass measurement: [2011Tu09](#).  
 Theoretical calculations: [2020Ri06](#), [2020RiZX](#), [2020RiZZ](#).

 $^{35}\text{Ar}$  LevelsCross Reference (XREF) Flags

<b>A</b>	$^{35}\text{K}$ $\epsilon$ decay (175 ms)	<b>E</b>	$^{24}\text{Mg}(^{16}\text{O}, \alpha n \gamma)$	<b>I</b>	$^{36}\text{Ar}(p, d)$
<b>B</b>	$^{36}\text{Ca}$ $\epsilon p$ decay (100.9 ms)	<b>F</b>	$^{32}\text{S}(\alpha, n)$	<b>J</b>	$^{36}\text{Ar}(d, t)$
<b>C</b>	$^1\text{H}(^{36}\text{Ar}, d)$	<b>G</b>	$^{33}\text{S}(^3\text{He}, n \gamma)$	<b>K</b>	$^{36}\text{Ar}(^3\text{He}, \alpha)$
<b>D</b>	$^{16}\text{O}(^{24}\text{Mg}, \alpha n \gamma)$	<b>H</b>	$^{35}\text{Cl}(^3\text{He}, t)$		

$E(\text{level})^\dagger$	$J^\pi$	$T_{1/2}$	XREF	Comments
0.0	$3/2^+$	1.7756 s 14	ABCDEFGH IJK	$\% \epsilon + \% \beta^+ = 100$ $\mu = +0.6322$ 2 ( <a href="#">2002Ma41</a> , <a href="#">2019StZV</a> ) $Q = -0.084$ 15 ( <a href="#">1996Ki04</a> , <a href="#">2021StZZ</a> ) $\mu$ : $\beta$ -NMR ( <a href="#">2002Ma41</a> ). Others: $+0.633$ 2 ( <a href="#">1965Ca04</a> ), $+0.633$ 7 ( <a href="#">1996Ki04</a> ) using $\beta$ -NMR. $Q$ : $\beta$ -NMR ( <a href="#">1996Ki04</a> ). $J^\pi$ : $L(p, d) = L(d, t) = L(^3\text{He}, \alpha) = L^1\text{H}(^{36}\text{Ar}, d) = 2$ from $0^+$ . Allowed $\epsilon + \beta^+$ feedings to $1/2^+$ levels in $^{35}\text{Cl}$ . Mirror level: $3/2^+$ $^{35}\text{Cl}$ g.s. $T_{1/2}$ : weighted average of 1.83 s 3 ( <a href="#">1956Ki29</a> ), 1.83 s 2 ( <a href="#">1959Al10</a> ), 1.79 s 1 ( <a href="#">1960Ja12</a> ), 1.84 s 10 ( <a href="#">1960Wa04</a> ), 1.76 s 3 ( <a href="#">1963Ne05</a> ), 1.770 s 6 ( <a href="#">1969Wi18</a> ), 1.787 s 12 ( <a href="#">1971Ge04</a> ), 1.774 s 4 ( <a href="#">1977Az01</a> ), and 1.7754 s 11 ( <a href="#">2006Ia05</a> ). Evaluated rms nuclear charge radius $R = 3.3636$ fm 42 ( <a href="#">2013An02</a> ). XREF: F(890) E(level): <a href="#">1963Ne05</a> ( $\alpha, n$ ) observed the first excited state in $^{35}\text{Ar}$ at 890 50 keV.
1184.08 25	$1/2^+$		ABC FG IJK	$J^\pi$ : $L(p, d) = L(d, t) = L(^3\text{He}, \alpha) = 0$ from $0^+$ . XREF: F(2030)I(1700)J(1700) E(level): <a href="#">1963Ne05</a> ( $\alpha, n$ ) observed the second excited state in $^{35}\text{Ar}$ at 2030 80 keV.
1750.78 22	$(5/2)^+$		A DEFG IJK	$J^\pi$ : $L(^3\text{He}, \alpha) = 2$ from $0^+$ . Mirror level: $5/2^+$ at 1763 keV in $^{35}\text{Cl}$ . $J^\pi$ : $\Delta J = 2$ $\gamma$ to $3/2^+$ in ( $^{16}\text{O}, \alpha n \gamma$ ). Mirror level: $7/2^+$ at 2646 keV in $^{35}\text{Cl}$ . XREF: I(2615)K(2649) $J^\pi$ : $L(p, d) = L(^3\text{He}, \alpha) = 2$ from $0^+$ with J dependence in (p, d).
2603.22 28	$7/2^{(+)}$		DE G	
2638.01 26	$3/2^+$		A IJK	

Continued on next page (footnotes at end of table)

**Adopted Levels, Gammas (continued)** $^{35}\text{Ar}$  Levels (continued)

E(level) <sup>†</sup>	J <sup>π</sup>	XREF		Comments
2982.79 12	5/2 <sup>+</sup>	A C	IJK	XREF: I(2970) J <sup>π</sup> : L(p,d)=L(d,t)=L( <sup>3</sup> He,α)=2 from 0 <sup>+</sup> with J dependence in (p,d).
3196.98 <sup>±</sup> 26	7/2 <sup>-</sup>	CDE G	IJK	J <sup>π</sup> : L(p,d)=L( <sup>3</sup> He,α)=3 from 0 <sup>+</sup> . ΔJ=1 γ to (5/2) <sup>+</sup> and ΔJ=2 γ to 3/2 <sup>+</sup> in ( <sup>16</sup> O,αγ) and ( <sup>24</sup> Mg,αγ); band assignment.
3882 5	1/2 <sup>+</sup>		K	J <sup>π</sup> : L( <sup>3</sup> He,α)=0 from 0 <sup>+</sup> .
4001 3	1/2 <sup>-</sup> , 3/2 <sup>-</sup>		K	J <sup>π</sup> : L( <sup>3</sup> He,α)=1 from 0 <sup>+</sup> .
4065.0? 4	(1/2 <sup>+</sup> , 3/2 <sup>+</sup> , 5/2 <sup>+</sup> )	A		XREF: A(?) J <sup>π</sup> : possibly allowed ε+β <sup>+</sup> feeding from 3/2 <sup>+</sup> parent with log ft=5.6 +4-2.
4113 4			K	
4135 4	1/2 <sup>-</sup> , 3/2 <sup>-</sup>		K	J <sup>π</sup> : L( <sup>3</sup> He,α)=1 from 0 <sup>+</sup> .
4359.0 5	(9/2 <sup>-</sup> )	DE	K	J <sup>π</sup> : ΔJ=(1) γ to 7/2 <sup>-</sup> in ( <sup>16</sup> O,αγ). Possible mirror level: 9/2 <sup>-</sup> at 4348 keV in <sup>35</sup> Cl.
4528.3 4	(1/2 <sup>+</sup> , 3/2 <sup>+</sup> , 5/2 <sup>+</sup> )	A	K	XREF: K(4515) J <sup>π</sup> : possibly allowed ε+β <sup>+</sup> feeding from 3/2 <sup>+</sup> parent with log ft=5.4 +4-2.
4725.9 6	1/2 <sup>+</sup>	A	Hi K	XREF: i(4756)K(4713) J <sup>π</sup> : L( <sup>3</sup> He,α)=0 from 0 <sup>+</sup> . Other: L(p,d)=0 from 0 <sup>+</sup> for a group at 4756 28.
4785.8 11	1/2 <sup>+</sup> , 3/2 <sup>+</sup> , 5/2 <sup>+</sup>	A	Hi K	XREF: i(4756)K(4774) J <sup>π</sup> : allowed ε+β <sup>+</sup> feeding from 3/2 <sup>+</sup> parent with log ft=5.2 2. Other: L(p,d)=0 from 0 <sup>+</sup> for a group at 4756 28.
5059 11			K	
5116 2	3/2 <sup>+</sup> , 5/2 <sup>+</sup>		HI K	E(level): weighted average of 5102 20 from (p,d) and 5116 2 from ( <sup>3</sup> He,α). J <sup>π</sup> : L( <sup>3</sup> He,α)=2 from 0 <sup>+</sup> . Discrepancy: L(p,d)=3 from 0 <sup>+</sup> (1968Ko04).
5207 3			H K	E(level): from ( <sup>3</sup> He,α).
5384.2 <sup>±</sup> 4	(11/2 <sup>-</sup> )	DE	HI K	XREF: I(5400) J <sup>π</sup> : ΔJ=2 γ to 7/2 <sup>-</sup> in ( <sup>16</sup> O,αγ) and ( <sup>24</sup> Mg,αγ); band assignment. γ to (9/2 <sup>-</sup> ) in ( <sup>16</sup> O,αγ) and ( <sup>24</sup> Mg,αγ). Possible mirror level: 11/2 <sup>-</sup> at 5407 keV in <sup>35</sup> Cl.
5482 2	3/2 <sup>+</sup> , 5/2 <sup>+</sup>		H K	E(level): from ( <sup>3</sup> He,α). J <sup>π</sup> : L( <sup>3</sup> He,α)=2 from 0 <sup>+</sup> .
5572.67 15	3/2 <sup>+</sup>	A	G	T=3/2 XREF: G(5537) J <sup>π</sup> : isobaric analog state of 3/2 <sup>+</sup> <sup>35</sup> K g.s. with log ft=3.31 4. L( <sup>3</sup> He,n)=(0) from 3/2 <sup>+</sup> .
5594 2	3/2 <sup>+</sup> , 5/2 <sup>+</sup>	C	HI K	XREF: C(5570) Evaluators consider the 5992 level to be different from the T=3/2 level at 5572.67 because (p,d) and ( <sup>3</sup> He,α) from T=0 should not populate T=3/2 levels. E(level): weighted average of 5598 20 from (p,d) and 5594 2 from ( <sup>3</sup> He,α). J <sup>π</sup> : L(p,d)=L( <sup>3</sup> He,α)=2 from 0 <sup>+</sup> .
5613.6 9	(11/2 <sup>-</sup> )	E		J <sup>π</sup> : Possible mirror level: 11/2 <sup>-</sup> at 5927 keV in <sup>35</sup> Cl.
5765.8 5	(13/2 <sup>-</sup> )	DE		J <sup>π</sup> : ΔJ=1 γ to (11/2 <sup>-</sup> ) in ( <sup>16</sup> O,αγ) and ( <sup>24</sup> Mg,αγ). ΔJ=(2) γ to (9/2 <sup>-</sup> ) in ( <sup>16</sup> O,αγ). Possible mirror level: 13/2 <sup>-</sup> at 6087 keV in <sup>35</sup> Cl.
5915 3			H JK	E(level): weighted average of 5913 5 from (d,t) and 5916 3 from ( <sup>3</sup> He,α).
5991 3			J	
6037 3	3/2 <sup>+</sup> , 5/2 <sup>+</sup>		HIJK	XREF: I(6024) E(level): weighted average of 6037 3 from (d,t) and 6036 3 from ( <sup>3</sup> He,α). Other: 6024 20 from (p,d). J <sup>π</sup> : L(p,d)=L( <sup>3</sup> He,α)=2 from 0 <sup>+</sup> .
6055? 3			J	XREF: J(?)
6076 3			J	
6163 2			JK	E(level): weighted average of 6164 3 from (d,t) and 6162 2 from ( <sup>3</sup> He,α).
6253 3			Jk	XREF: k(6262)

Continued on next page (footnotes at end of table)

**Adopted Levels, Gammas (continued)** $^{35}\text{Ar}$  Levels (continued)

E(level) <sup>†</sup>	J <sup>π</sup>	XREF	Comments
6273 3		Jk	XREF: k(6262)
6302 3		J	
6332 3		J	
6345 3	(1/2,3/2,5/2)	A J	E(level): From (d,t). Other: 6348 11 from $^{35}\text{K}$ $\varepsilon$ decay. J <sup>π</sup> : $\varepsilon+\beta^+$ feeding from 3/2 <sup>+</sup> parent with log ft=7.2 1.
6415 2		J	
6439? 4		J	XREF: J(?)
6460 3		J	
6523 3		J	
6557 3		J	
6585 3		J	
6606 3		iJ	XREF: i(6620)
6616 2	1/2 <sup>+</sup>	iJK	XREF: i(6620) E(level): weighted average of 6617 2 from (d,t) and 6615 3 from ( $^3\text{He},\alpha$ ). Other: 6620 30 from (p,d). J <sup>π</sup> : L( $^3\text{He},\alpha$ )=L(p,d)=0 from 0 <sup>+</sup> .
6644 3		iJ	XREF: i(6620)
6651 3		iJ	XREF: i(6620)
6673 4	5/2 <sup>-</sup> ,7/2 <sup>-</sup>	IJ	XREF: I(6700) E(level): weighted average of 6700 20 from (p,d) and 6672 3 from (d,t). J <sup>π</sup> : L(p,d)=3 from 0 <sup>+</sup> .
6823 2	3/2 <sup>+</sup> ,5/2 <sup>+</sup>	I K	E(level): from ( $^3\text{He},\alpha$ ). Other: 6820 30 from (p,d). J <sup>π</sup> : L(p,d)=2 from 0 <sup>+</sup> .
6948 2		K	
7044 4	3/2 <sup>+</sup> ,5/2 <sup>+</sup>	A I K	XREF: I(7030) E(level): weighted average of 7053 11 from $^{35}\text{K}$ $\varepsilon$ decay, 7030 20 from (p,d), and 7043 4 from ( $^3\text{He},\alpha$ ). J <sup>π</sup> : L(p,d)=2 from 0 <sup>+</sup> .
7117 10		K	
7255 11		A	
7289 10		A K	E(level): weighted average of 7283 11 from $^{35}\text{K}$ $\varepsilon$ decay and 7293 10 from ( $^3\text{He},\alpha$ ).
7427 10		A K	E(level): weighted average of 7431 11 from $^{35}\text{K}$ $\varepsilon$ decay and 7423 10 from ( $^3\text{He},\alpha$ ).
7509 10	1/2 <sup>+</sup> ,3/2 <sup>+</sup> ,5/2 <sup>+</sup>	A K	E(level): weighted average of 7518 11 from $^{35}\text{K}$ $\varepsilon$ decay and 7502 10 from ( $^3\text{He},\alpha$ ). J <sup>π</sup> : allowed $\varepsilon+\beta^+$ feeding from 3/2 <sup>+</sup> parent with log ft<5.0.
7840 10		K	
8019 10		K	
8109.7 <sup>‡</sup> 13	(15/2 <sup>-</sup> )	E	J <sup>π</sup> : $\gamma$ to (11/2 <sup>-</sup> ) and (13/2 <sup>-</sup> ) in ( $^{16}\text{O},\alpha n\gamma$ ); band assignment. Possible mirror level: 15/2 <sup>-</sup> at 8319 keV in $^{35}\text{Cl}$ .
8212.6 8	(15/2 <sup>-</sup> )	E	J <sup>π</sup> : $\Delta J=2$ $\gamma$ to (11/2 <sup>-</sup> ) and $\gamma$ to (13/2 <sup>-</sup> ) in ( $^{16}\text{O},\alpha n\gamma$ ). Possible mirror level: 15/2 <sup>-</sup> at 8487 keV in $^{35}\text{Cl}$ .
8393? 20	1/2 <sup>+</sup> ,3/2 <sup>+</sup> ,5/2 <sup>+</sup>	A	XREF: A(?) E(level): From ( $^{35}\text{K}$ $\varepsilon$ decay). J <sup>π</sup> : allowed $\varepsilon+\beta^+$ feeding from 3/2 <sup>+</sup> parent with log ft=4.6 +3-2.
9906.0 <sup>‡</sup> 20	(19/2 <sup>-</sup> )	E	J <sup>π</sup> : $\Delta J=2$ $\gamma$ to (15/2 <sup>-</sup> ) in ( $^{16}\text{O},\alpha n\gamma$ ); band assignment. Possible mirror level: 19/2 <sup>-</sup> at 10180 keV in $^{35}\text{Cl}$ .
12277.0 <sup>‡</sup> 32	(23/2 <sup>-</sup> )	E	J <sup>π</sup> : $\Delta J=2$ $\gamma$ to (19/2 <sup>-</sup> ) in ( $^{16}\text{O},\alpha n\gamma$ ); band assignment. Possible mirror level: 23/2 <sup>-</sup> at 12571 keV in $^{35}\text{Cl}$ .

<sup>†</sup> From a least-squares fit to  $\gamma$ -ray energies for levels connected with  $\gamma$  transitions; from particle-transfer reactions or  $^{35}\text{K}$

**Adopted Levels, Gammas (continued)** $^{35}\text{Ar}$  Levels (continued) $\varepsilon+\beta^+$ -delayed proton decays for other levels. $\ddagger$  Band(A): Band based on  $f_{7/2}$  orbital.

$E_i(\text{level})$	$J_i^\pi$	$\gamma(^{35}\text{Ar})$		$E_f$	$J_f^\pi$	Mult. $\ddagger$	Comments
		$E_\gamma^\ddagger$	$I_\gamma^\ddagger$				
1184.08	$1/2^+$	1184.1 3	100	0.0	$3/2^+$		$E_\gamma$ : weighted average of 1184.0 3 from $^{35}\text{K}$ $\varepsilon$ decay and 1184.3 4 from $^{36}\text{Ca}$ $\varepsilon p$ decay.
1750.78	$(5/2)^+$	1750.6 3	100	0.0	$3/2^+$		$E_\gamma$ : weighted average of 1750.5 3 from $^{35}\text{K}$ $\varepsilon$ decay, 1750.7 4 from $(^{24}\text{Mg}, \alpha n\gamma)$ , and 1750.8 5 from $(^{16}\text{O}, \alpha n\gamma)$ .
2603.22	$7/2^{(+)}$	851.9 9	12.3 33	1750.78	$(5/2)^+$		$E_\gamma$ : weighted average of 852 1 from $(^{24}\text{Mg}, \alpha n\gamma)$ and 851.8 9 from $(^{16}\text{O}, \alpha n\gamma)$ .
		2603.0 5	100 10	0.0	$3/2^+$	Q	$I_\gamma$ : weighted average of 10 5 from $(^{24}\text{Mg}, \alpha n\gamma)$ and 13.3 33 from $(^{16}\text{O}, \alpha n\gamma)$ . $E_\gamma$ : weighted average of 2603.0 5 from $(^{24}\text{Mg}, \alpha n\gamma)$ and 2602.6 15 from $(^{16}\text{O}, \alpha n\gamma)$ . $I_\gamma$ : other: 100 22 from $(^{24}\text{Mg}, \alpha n\gamma)$ .
2638.01	$3/2^+$	886.8 <sup>#</sup> 5	16 <sup>#</sup> 6	1750.78	$(5/2)^+$		
		2638.0 <sup>#</sup> 4	100 <sup>#</sup> 13	0.0	$3/2^+$		
2982.79	$5/2^+$	1798.9 <sup>#</sup> 5	3.5 <sup>#</sup> 6	1184.08	$1/2^+$		
		2982.68 <sup>#</sup> 13	100 <sup>#</sup> 4	0.0	$3/2^+$		
3196.98	$7/2^-$	593.7 2	16.4 30	2603.22	$7/2^{(+)}$		$E_\gamma$ : weighted average of 593 1 from $(^{24}\text{Mg}, \alpha n\gamma)$ and 593.7 2 from $(^{16}\text{O}, \alpha n\gamma)$ .
		1446.2 2	100 8	1750.78	$(5/2)^+$	D	$I_\gamma$ : weighted average of 16 8 from $(^{24}\text{Mg}, \alpha n\gamma)$ and 16.4 30 from $(^{16}\text{O}, \alpha n\gamma)$ . $E_\gamma$ : weighted average of 1446.2 2 from $(^{24}\text{Mg}, \alpha n\gamma)$ , 1446.1 6 from $(^{16}\text{O}, \alpha n\gamma)$ , and 1446.0 6 from $(^3\text{He}, n\gamma)$ . $I_\gamma$ : other: 100 9 from $(^{24}\text{Mg}, \alpha n\gamma)$ .
		3197.0 7	21 5	0.0	$3/2^+$	Q	$E_\gamma$ : from $(^{24}\text{Mg}, \alpha n\gamma)$ . Other: 3197 6 from $(^{16}\text{O}, \alpha n\gamma)$ . $I_\gamma$ : weighted average of 18 5 from $(^{24}\text{Mg}, \alpha n\gamma)$ and 24 5 from $(^{16}\text{O}, \alpha n\gamma)$ .
4065.0?	$(1/2^+, 3/2^+, 5/2^+)$	1426.8 <sup>#</sup> 4	100	2638.01	$3/2^+$		
4359.0	$(9/2^-)$	1162.0 8	65 24	3196.98	$7/2^-$	(D)	$E_\gamma$ : weighted average of 1162 1 from $(^{24}\text{Mg}, \alpha n\gamma)$ and 1162.0 8 from $(^{16}\text{O}, \alpha n\gamma)$ . $I_\gamma$ : unweighted average of 41 11 from $(^{24}\text{Mg}, \alpha n\gamma)$ and 88 18 from $(^{16}\text{O}, \alpha n\gamma)$ .
		1756 1	100 15	2603.22	$7/2^{(+)}$		$E_\gamma$ : weighted average of 1756 1 from $(^{24}\text{Mg}, \alpha n\gamma)$ and 1756.3 14 from $(^{16}\text{O}, \alpha n\gamma)$ .

Continued on next page (footnotes at end of table)


**Adopted Levels, Gammas (continued)**

$\gamma(^{35}\text{Ar})$ (continued)						
$E_i(\text{level})$	$J_i^\pi$	$E_\gamma^\dagger$	$I_\gamma^\dagger$	$E_f$	$J_f^\pi$	Mult. <sup>‡</sup>
						Comments
						$I_\gamma$ : from ( $^{24}\text{Mg}, \alpha n \gamma$ ). Other: 100 53 from ( $^{16}\text{O}, \alpha n \gamma$ ).
4528.3	(1/2 <sup>+</sup> , 3/2 <sup>+</sup> , 5/2 <sup>+</sup> )	4527.9 <sup>#</sup> 7	100	0.0	3/2 <sup>+</sup>	
4725.9	1/2 <sup>+</sup>	3542.0 <sup>#</sup> 6	100 <sup>#</sup> 21	1184.08	1/2 <sup>+</sup>	
		4724.5 <sup>#</sup> 11	41 <sup>#</sup> 17	0.0	3/2 <sup>+</sup>	
4785.8	1/2 <sup>+</sup> , 3/2 <sup>+</sup> , 5/2 <sup>+</sup>	4785.4 <sup>#</sup> 11	100	0.0	3/2 <sup>+</sup>	
5384.2	(11/2 <sup>-</sup> )	1025.2 4	14 4	4359.0	(9/2 <sup>-</sup> )	
		2187.1 4	100 6	3196.98	7/2 <sup>-</sup>	Q
						$E_\gamma$ : weighted average of 1025 1 from ( $^{24}\text{Mg}, \alpha n \gamma$ ) and 1025.2 4 from ( $^{16}\text{O}, \alpha n \gamma$ ). $I_\gamma$ : weighted average of 21 8 from ( $^{24}\text{Mg}, \alpha n \gamma$ ) and 12 4 from ( $^{16}\text{O}, \alpha n \gamma$ ). $E_\gamma$ : weighted average of 2187.4 4 from ( $^{24}\text{Mg}, \alpha n \gamma$ ) and 2186.8 4 from ( $^{16}\text{O}, \alpha n \gamma$ ). $I_\gamma$ : other: 100 13 from ( $^{24}\text{Mg}, \alpha n \gamma$ ).
5572.67	3/2 <sup>+</sup>	1044.4 <sup>#</sup> 4	2.5 <sup>#</sup> 8	4528.3	(1/2 <sup>+</sup> , 3/2 <sup>+</sup> , 5/2 <sup>+</sup> )	
		1507.4 <sup>#</sup> 5	3.7 <sup>#</sup> 8	4065.0?	(1/2 <sup>+</sup> , 3/2 <sup>+</sup> , 5/2 <sup>+</sup> )	
		2589.8 <sup>#</sup> 1	100 <sup>#</sup> 4	2982.79	5/2 <sup>+</sup>	
		2934.5 <sup>#</sup> 5	6.7 <sup>#</sup> 12	2638.01	3/2 <sup>+</sup>	
		3821.7 <sup>#</sup> 7	6.7 <sup>#</sup> 14	1750.78	(5/2 <sup>+</sup> )	
		4387.2 <sup>#</sup> 9	6.7 <sup>#</sup> 16	1184.08	1/2 <sup>+</sup>	
		5572.3 <sup>#</sup> 10	11.7 <sup>#</sup> 31	0.0	3/2 <sup>+</sup>	
5613.6	(11/2 <sup>-</sup> )	1254.6 8	100	4359.0	(9/2 <sup>-</sup> )	
5765.8	(13/2 <sup>-</sup> )	381.6 1	100 10	5384.2	(11/2 <sup>-</sup> )	D
		1406.9 7	17.2 35	4359.0	(9/2 <sup>-</sup> )	(Q)
8109.7	(15/2 <sup>-</sup> )	2342.6 28	100 25	5765.8	(13/2 <sup>-</sup> )	
		2725.7 14	50 13	5384.2	(11/2 <sup>-</sup> )	
8212.6	(15/2 <sup>-</sup> )	2446.6 16	21 7	5765.8	(13/2 <sup>-</sup> )	
		2828.3 7	100 18	5384.2	(11/2 <sup>-</sup> )	Q
9906.0	(19/2 <sup>-</sup> )	1693.3 27	100 20	8212.6	(15/2 <sup>-</sup> )	Q
		1796.3 25	67 20	8109.7	(15/2 <sup>-</sup> )	Q
12277.0	(23/2 <sup>-</sup> )	2370.9 25	100	9906.0	(19/2 <sup>-</sup> )	Q

<sup>†</sup> From ( $^{16}\text{O}, \alpha n \gamma$ ), unless otherwise noted.<sup>‡</sup> Deduced by evaluators from measured  $\gamma\gamma(\theta)(\text{ADO})$  in ( $^{16}\text{O}, \alpha n \gamma$ ) and ratios of yields  $R(\gamma(\theta))$  in ( $^{24}\text{Mg}, \alpha n \gamma$ ), unless otherwise noted.<sup>#</sup> From  $^{35}\text{K}$   $\varepsilon$  decay.

Adopted Levels, GammasLevel Scheme


Intensities: Relative photon branching from each level



1.7756 s 14

Adopted Levels, Gammas

Band(A): Band based on f<sub>7/2</sub>  
orbital



<sup>35</sup>Ar<sub>17</sub>