

Adopted Levels, Gammas

$Q(\beta^-)=10470$  40;  $S(n)=2470$  40;  $S(p)=18680$  40;  $Q(\alpha)=-13690$  40 2021Wa16

$S(2n)=10020$  40,  $S(2p)=33930$  40,  $Q(\beta^-n)=2090$  40 (2021Wa16).

Isotope discovery (2012Th10):  $^{232}\text{Th}(^{40}\text{Ar},X)$  at Dubna (1971Ar32).

$^{35}\text{Si}$  production:

2015Mo17:  $^9\text{Be}(^{40}\text{Ar},X)$  at  $E(^{40}\text{Ar})=95$  MeV/nucleon at RIKEN. Measured angular distributions and transverse momentum distributions of fragments. Deduced formulation for the width of transverse momentum distribution as a function of fragment velocity.

2012Kw02:  $^9\text{Be}, ^{\text{nat}}\text{Ni}, ^{181}\text{Ta}(^{40}\text{Ar},X)$  at  $E(^{40}\text{Ar})=140$  MeV/nucleon at NSCL. Measured fragmentation cross sections, parallel momentum transfers, and widths. Compared with empirical formula EPAX, and predictions from internuclear cascade and deep inelastic models using Monte Carlo ISABEL-GEMINI and DIT-GEMINI codes.

2012Zh06:  $^9\text{Be}, ^{181}\text{Ta}(^{40}\text{Ar},X)$  at  $E(^{40}\text{Ar})=57$  MeV/nucleon at HIRFL. Measured momentum distributions and production cross sections of fragments. Observed competition between projectile fragmentation and other mechanisms. Compared with EPAX, abrasion- ablation, and HIPSE models. Studied target dependence of fragment cross sections.

2007No13:  $^9\text{Be}(^{40}\text{Ar},X)$  at  $E(^{40}\text{Ar})=100$  MeV/nucleon at RIKEN. Measured fragment momentum distributions and production cross sections.

2006Ro34:  $^2\text{H}(^{42}\text{S},X)$  at  $E(^{42}\text{S})=99.8$  MeV/nucleon at NSCL. Measured production cross sections.

1997Fo01:  $^{208}\text{Pb}(^{37}\text{Cl},X)$  at  $E(^{37}\text{Cl})=230$  MeV at Legnaro. Measured yields.

$^{35}\text{Si}$  decay measurements:

1986Du07, 1986HuZW, 1987DuZU, 1988DuZS, 1988DuZT:  $^9\text{Be}(^{40}\text{Ar},X)$  at GANIL. Measured  $T_{1/2}$  and  $\beta^-$ -delayed  $\gamma$  rays.

2007Ne14:  $^{35}\text{Si}$  g.s. magnetic moment and  $g$ -factor using  $\beta$ -NMR.

$^{35}\text{Si}$  radius measurements:

2006Kh08:  $^{35}\text{Si}$  produced by  $^{181}\text{Ta}(^{48}\text{Ca},X)$  fragmentation at  $E(^{48}\text{Ca})=60.3$  MeV/nucleon at GANIL. Measured energy-integrated reaction cross sections at 30-65 MeV/nucleon using a silicon telescope as both active target and detector. Deduced reduced strong absorption radii, isospin dependence, and possible halo structure or large deformation.

1999Ai02:  $\text{Si}(^{35}\text{Si},X)$  at NSCL. Measured energy-integrated reaction cross sections at  $E=38$ -80 MeV/ nucleon. Deduced strong absorption radii.

$^{35}\text{Si}$  mass measurements: 1986Fi06, 1986Sm05, 1984Ma49.

Theoretical calculations (binding energies, deformation, quadrupole moments, radii, levels,  $J^\pi$ , etc.): 2011Ka03, 2009No01, 2008Wi11, 2007Ch82, 2004Kh16, 1999Du05, 1994Mo37, 1994Po05, 1987Wa10, 1986Wo02.

 $^{35}\text{Si}$  LevelsCross Reference (XREF) Flags

- A  $^{35}\text{Al}$   $\beta^-$  decay (37.2 ms)  
 B  $^1\text{H}(^{34}\text{Si},p)$ :from IAR  
 C  $^2\text{H}(^{34}\text{Si},p\gamma)$   
 D  $^9\text{Be}(^{36}\text{Si},^{35}\text{Si}\gamma)$

$E(\text{level})^\dagger$	$J^\pi^\ddagger$	$T_{1/2}$	XREF	Comments
0	$(7/2)^-$	0.78 s 12	ABCD	$\% \beta^- = 100$ ; $\% \beta^- n < 5$ (1995ReZZ/2008ReZZ) $\mu = (-)1.638$ 4 (2007Ne14,2014StZZ) $\mu$ : Using $\beta$ -NMR on a polarized fragment beam (2007Ne14). $J^\pi$ : (d,p)=3 in $^2\text{H}(^{34}\text{Si},p\gamma)$ , $7/2^-$ from shell-model predictions, and systematic trends in Si isotopes. $T_{1/2}$ : From $\beta$ -decay measurement (1988DuZT). In an earlier paper by the same group (1986Du07) value given is 0.87 s 17. The evaluators adopt the more recent value. Reduced strong absorption radius $r_0^2=1.261$ fm <sup>2</sup> 35 from the energy-integrated $\sigma$ of $\text{Si}(^{35}\text{Si},X)$ (2006Kh08) and $r_0^2=1.258$ fm <sup>2</sup> 92 from the energy-integrated $\sigma$ of $\text{Si}(^{35}\text{Si},X)$ (1999Ai02). Configuration= $\nu f_{7/2}$ . $J^\pi$ : L(d,p)=1 in $^2\text{H}(^{34}\text{Si},p\gamma)$ , $3/2^-$ from shell-model predictions, and
909.95 23	$(3/2)^-$ 55	PS 14	ABCD	

Continued on next page (footnotes at end of table)

**Adopted Levels, Gammas (continued)** $^{35}\text{Si}$  Levels (continued)

<u>E(level)<sup>†</sup></u>	<u>J<sup>π</sup><sup>‡</sup></u>	<u>T<sub>1/2</sub></u>	<u>XREF</u>	<u>Comments</u>
				systematic trends in Si isotopes.
				T <sub>1/2</sub> : From analysis of broadened lineshape in $^9\text{Be}(^{36}\text{Si}, ^{35}\text{Si}\gamma)$ (2014St18).
				Configuration= $\nu p_{3/2}$ .
973.88 18	(3/2 <sup>+</sup> )	5.9 ns 6	AB D	T <sub>1/2</sub> : From the time spectrum of delayed coincidences in $^{35}\text{Al } \beta^-$ decay (2001Nu01).
1444?	(1/2 <sup>+</sup> )		B	E(level), J <sup>π</sup> : corresponding to a possible IAR in $^{35}\text{P}$ with L(p)=0 from R-Matrix analysis in $^1\text{H}(^{34}\text{Si}, p)$ : From IAR (2012Im01).
1689 3	1/2 <sup>+</sup>		D	J <sup>π</sup> : L(n)=0 in $^9\text{Be}(^{36}\text{Si}, ^{35}\text{Si}\gamma)$ .
1970 6			D	
2044 5	(1/2) <sup>-</sup>		CD	J <sup>π</sup> : L(d,p)=1 in $^2\text{H}(^{34}\text{Si}, p\gamma)$ , 1/2 <sup>-</sup> from shell-model predictions.
				Configuration= $\nu p_{1/2}$ .
2168.2 4	5/2 <sup>+</sup>		AB D	J <sup>π</sup> : corresponding to an IAR in $^{35}\text{P}$ with L(p)=2 and J=5/2 <sup>+</sup> from R-Matrix analysis in $^1\text{H}(^{34}\text{Si}, p)$ : From IAR (2012Im01).
2194?	(1/2 <sup>-</sup> , 3/2 <sup>-</sup> )		B	E(level), J <sup>π</sup> : corresponding to a possible IAR in $^{35}\text{P}$ with L(p)=1 from R-Matrix analysis in $^1\text{H}(^{34}\text{Si}, p)$ : From IAR (2012Im01).
2275 6			D	
2377 7			D	
3140			A	
3450			A	
3611? 8			D	
3770			A	
5190			A	
≈5500	(5/2) <sup>-</sup>		C	J <sup>π</sup> : L(d,p)=3 in $^2\text{H}(^{34}\text{Si}, p\gamma)$ .
				Configuration= $\nu f_{5/2}$ .
5760			A	
6330			A	
7360			A	
7690			A	

<sup>†</sup> From a least-squares fit to  $\gamma$ -ray energies if applicable. Values without uncertainties are from  $^{35}\text{Al } \beta^-$  decay, unless otherwise noted.

<sup>‡</sup> From shell mode predictions and systematic trends on Si isotopes.

 $\gamma(^{35}\text{Si})$ 

<u>E<sub>i</sub>(level)</u>	<u>J<sub>i</sub><sup>π</sup></u>	<u>E<sub>γ</sub><sup>†</sup></u>	<u>I<sub>γ</sub><sup>†</sup></u>	<u>E<sub>f</sub></u>	<u>J<sub>f</sub><sup>π</sup></u>	<u>Mult.</u>	<u>α<sup>#</sup></u>	<u>Comments</u>
909.95	(3/2) <sup>-</sup> 55	910.11 30	100	0	(7/2) <sup>-</sup>	[E2]		
973.88	(3/2 <sup>+</sup> )	64.1 3	100	909.95	(3/2) <sup>-</sup> 55	[E1]	0.0368 8	B(E1)(W.u.)=0.00036 4
		973.78 20	11.8 24	0	(7/2) <sup>-</sup>	[M2]		B(M2)(W.u.)=0.059 14
1689	1/2 <sup>+</sup>	715 <sup>‡</sup> 4	14.6 <sup>‡</sup> 15	973.88	(3/2 <sup>+</sup> )			
		780 <sup>‡</sup> 4	100 <sup>‡</sup> 8	909.95	(3/2) <sup>-</sup> 55			
1970		1970 <sup>‡</sup> 6	100	0	(7/2) <sup>-</sup>			
2044	(1/2) <sup>-</sup>	1134 <sup>‡</sup> 5	100	909.95	(3/2) <sup>-</sup> 55			
2168.2	5/2 <sup>+</sup>	1194.2 4	35 8	973.88	(3/2 <sup>+</sup> )			
		2168.2 6	100 20	0	(7/2) <sup>-</sup>			
2275		2275 <sup>‡</sup> 6	100	0	(7/2) <sup>-</sup>			
2377		2377 <sup>‡</sup> 7	100	0	(7/2) <sup>-</sup>			
3611?		3611 <sup>‡</sup> 8	100	0	(7/2) <sup>-</sup>			

Continued on next page (footnotes at end of table)

**Adopted Levels, Gammas (continued)** $\gamma(^{35}\text{Si})$  (continued)

† From  $^{35}\text{Al}$   $\beta^-$  decay, unless otherwise noted.

‡ From  $^9\text{Be}(^{36}\text{Si}, ^{35}\text{Si}\gamma)$ .

# Total theoretical internal conversion coefficients, calculated using the BrIcc code (2008Ki07) with “Frozen Orbitals” approximation based on  $\gamma$ -ray energies, assigned multiplicities, and mixing ratios, unless otherwise specified.

**Adopted Levels, Gammas**Level Scheme

Intensities: Relative photon branching from each level

