

$^2\text{H}(^{34}\text{Si}, ^{35}\text{P}\gamma)$  2007GeZX

$^{34}\text{Si}(\text{d}, \text{n})^{35}\text{P}$  from  $J^\pi=0^+$   $^{34}\text{Si}$  g.s. in inverse kinematics.

2007GeZX: 30-AMeV  $^{34}\text{Si}$  beam on 30-mg/cm<sup>2</sup> CD<sub>2</sub> secondary target at GANIL. Heavy ions produced in reactions were identified by the VAMOS spectrometer.  $\gamma$  rays were detected using the EXOGAM germanium clover array. Measured Doppler-corrected  $E_\gamma$ ,  $I_\gamma$ ,  $\gamma\gamma$ -coin, and  $(^{35}\text{P})\gamma$ -coin. Deduced levels,  $J$ ,  $\pi$ . Compared with shell-model calculations.

 $^{35}\text{P}$  Levels

$E(\text{level})^\dagger$

0  
2386.5 8  
3859.9 8  
4100.9 13  
4492.9 16  
4868.9 13

<sup>†</sup> From a least-squares fit to  $\gamma$ -ray energies, assuming  $\Delta E_\gamma=1$  keV since uncertainties in 2007GeZX were not given.

 $\gamma(^{35}\text{P})$ 

<u><math>E_\gamma^\dagger</math></u>	<u><math>E_i(\text{level})</math></u>	<u><math>E_f</math></u>
241	4100.9	3859.9
392	4492.9	4100.9
1009	4868.9	3859.9
1473 <sup>‡</sup>	3859.9	2386.5
2386	2386.5	0
3860	3859.9	0

<sup>†</sup> From 2007GeZX.

<sup>‡</sup> Placement of transition in the level scheme is uncertain.

$^2\text{H}(^{34}\text{Si}, ^{35}\text{P}\gamma)$  2007GeZX

Legend

Level Scheme

 - - - - -  $\rightarrow$   $\gamma$  Decay (Uncertain)
