

Adopted Levels, Gammas

- $Q(\beta^-)=14170$ 40; $S(n)=5297$ 8; $S(p)=15830.7$ 89; $Q(\alpha)=-14894.0$ 78 2021Wa16
 $S(p), Q(\alpha)$: Deduced by the evaluator using mass excesses of 8318 5 for ^{34}Mg : a weighted average of 8323 7 (2019As04) and 8315 5 (2025Ly01), and 12245.3 26 for ^{31}Na measured by 2025Ly01; -224 7 for ^{35}Al from 2021Wa16. Values from 2021Wa16:
 $S(p)=15836$ 10, $Q(\alpha)=-14895$ 16.
 $S(2p)=38400.5$ 85, $Q(\epsilon)=-15753$ 10, from mass excesses of 23598.8 43 for ^{33}Na and 15529.5 71 for ^{35}Mg measured by 2025Ly01; -224 7 for ^{35}Al from 2021Wa16. Values from 2021Wa16: $S(2p)=38580$ 450, $Q(\epsilon)=-15860$ 270.
 $S(2n)=7869$ 10, $Q(\beta^-n)=11697$ 7 (2021Wa16).
Isotope discovery (2012Th10): $\text{C}(^{40}\text{Ar}, \text{X})$ projectile fragmentation at Berkeley (1979Sy01).
2017Ha23: ^{35}Al produced by $^9\text{Be}(^{40}\text{Ar}, \text{X})$ at HIRFL. Measured $T_{1/2}$.
2015Mo17: ^{35}Al produced by $^9\text{Be}(^{40}\text{Ar}, \text{X})$ at $E(^{40}\text{Ar})=95$ MeV/nucleon at RIKEN. Measured time-of-flight, energy loss, angular distribution of fragments, analyzed transverse momentum distributions for fragments, and deduced formulation for the width of transverse momentum distribution.
2012No05: $^{33,34,35,36}\text{Al}$ produced by $\text{Be}(^{48}\text{Ca}, \text{X})$ fragmentation at $E(^{48}\text{Ca})=1$ GeV/nucleon at GSI. Measured 1n removal cross sections and longitudinal momentum distributions of the residues. $\sigma(^{35}\text{Al} \rightarrow ^{34}\text{Al})=75$ mb 4 and $\sigma(^{36}\text{Al} \rightarrow ^{35}\text{Al})=95$ mb 5.
2012Kw02: ^{35}Al produced by $^9\text{Be}, ^{\text{nat}}\text{Ni}, ^{181}\text{Ta}(^{40}\text{Ar}, \text{X})$ at $E(^{40}\text{Ar})=140$ MeV/nucleon at NSCL. Measured fission fragment spectra, average isobaric velocities, parallel momentum transfers, widths, fragment σ . Comparison with empirical formula EPAX, and predictions from internuclear cascade and deep inelastic models using Monte Carlo ISABEL-GEMINI and DIT-GEMINI codes.
2012Zh06: ^{35}Al produced by $^9\text{Be}(^{40}\text{Ar}, \text{X})$ at $E(^{40}\text{Ar})=57$ MeV/nucleon at HIRFL. Measured particle spectra, energy loss, time of flight, fragment yields, momentum distributions, cross sections; deduced fragment excitation energies, mass yield ratios.
2010Ro23: ^{35}Al produced by $^9\text{Be}(^{40}\text{Ar}, \text{X})$ fragmentation at $E(^{40}\text{Ar})=700$ MeV/nucleon at GSI. Measured 1n knockout cross sections and longitudinal momentum distributions of the residues. $\sigma(^{35}\text{Al} \rightarrow ^{34}\text{Al})=65$ mb 18.
2007No13: ^{35}Al produced by $^9\text{Be}(^{40}\text{Ar}, \text{X})$ fragmentation at $E(^{40}\text{Ar})=100$ MeV/nucleon at RIKEN. Measured fragment momentum distributions and production cross sections.
2006Kh08: ^{35}Al produced by $^{181}\text{Ta}(^{48}\text{Ca}, \text{X})$ fragmentation at $E(^{48}\text{Ca})=60.3$ MeV/nucleon at GANIL. Measured energy-integrated reaction cross sections at 30-65 MeV/nucleon using a silicon telescope as both active target and detector. Deduced radii, isospin dependence, and possible halo structure or large deformation.
2006FuZX: ^{35}Al produced by $\text{Be}, \text{C}(^{40}\text{Ar}, \text{X})$ at $E(^{40}\text{Ar})=63$ MeV/nucleon at RIKEN. $\text{He}(^{35}\text{Al}, \text{X})$ at $E(^{35}\text{Al})=40$ MeV/nucleon. Measured E_γ , I_γ .
2005Ti11, 2006AnZW: fragmentation of ^{36}S beam at 78 MeV/nucleon at GANIL. Measured β -delayed E_γ , $T_{1/2}$ and delayed neutron emission probability. Deduced levels, J^π , $\log ft$ for ^{35}Si . (conference paper).
2001Nu01, 2002Nu02: ^{35}Al produced at the ISOLDE facility at CERN in fragmentation with 1.0 GeV proton beam on a uranium carbide target. Measured E_γ , E_γ , $\gamma\gamma$, $\beta\gamma$ -coin, $T_{1/2}$ and delayed-neutron branches. Deduced levels, J^π , $\log ft$ for ^{35}Si .
1999YoZW: ^{35}Al produced by $^9\text{Be}(^{48}\text{Ca}, \text{X})$ and $^{181}\text{Ta}(^{48}\text{Ca}, \text{X})$ fragmentations at $E(^{48}\text{Ca})=70$ MeV/nucleon at RIKEN. Measured $T_{1/2}$ and delayed neutron emission probabilities.
1989Le16, 1989MuZU: ^{35}Al produced by $^{181}\text{Ta}(^{48}\text{Ca}, \text{X})$ fragmentation at $E(^{48}\text{Ca})=55$ MeV/nucleon at GANIL. Measured $T_{1/2}$ and delayed neutron emission probabilities.
1988Mu08, 1988MuZY, 1988BaYZ: ^{35}Al produced by $^{181}\text{Ta}(^{86}\text{Kr}, \text{X})$ fragmentation at $E(^{86}\text{Kr})=45$ MeV/nucleon at GANIL. Measured $T_{1/2}$ and delayed neutron emission probabilities.
Mass measurements: 2017Ga20, 2007Ju03, 1991Or01, 1991Zh24, 1987Gi05.
Theoretical calculations (binding energies, deformation, quadrupole moments, radii, levels, J , π , mass, $T_{1/2}$, etc): 2016Sa46, 2014Ca21, 2013Li39, 2013Sh05, 2011Ki12, 2009Yo05, 2004Kh16, 1994Po05.

 ^{35}Al LevelsCross Reference (XREF) Flags

- A $^{36}\text{Mg} \beta^-n$ decay (6.9 ms)
B $^9\text{Be}(^{36}\text{Si}, ^{35}\text{Al}\gamma)$
C $\text{Pb}(^{35}\text{Al}, ^{34}\text{Al}n\gamma)$
D Coulomb excitation

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{35}Al Levels (continued)

$E(\text{level})^\dagger$	J^π	$T_{1/2}$	XREF	Comments
0	$(5/2)^+$	38.3 ms 4	ABCD	$\% \beta^- = 100$; $\% \beta^- n = 38.2$ (2015Bi05); $\% \beta^- 2n = 0$ (2001Nu01) $\% \beta^- n$: From 2015Bi05 evaluation; weighted average of 38.2 (2005Ti11, 38.3 in 2006AnZW) and 41.13 (2001Nu01, 2002Nu02). Others: 26.4 (1995ReZZ, 2008ReZZ); 40.10 (1989Le16); 87.37-25 (1988Mu08). J^π : From shell-model calculations and $L(^{36}\text{Si}, ^{35}\text{Al}) = 2$ from 0^+ . Major configurations deduced by 2017Ch36 from Coulomb breakup of ^{35}Al on Pb target: for $J^\pi(^{35}\text{Al g.s.}) = 5/2^+$, (g.s., 4^- in ^{34}Al) $\otimes \nu p_{3/2} + (46 \text{ keV}, 1^+$ in ^{34}Al) $\otimes \nu d_{3/2}$. For $J^\pi = 1/2^+$ or $3/2^+$ of $^{35}\text{Al g.s.}$: (g.s., 4^- in ^{34}Al) $\otimes \nu f_{7/2} + (46 \text{ keV}, 1^+$ in ^{34}Al) $\otimes \nu s_{1/2}$. Other configurations for $J^\pi = 1/2^+, 3/2^+$ of $^{35}\text{Al g.s.}$: (46 keV, 1^+ in ^{34}Al) $\otimes \nu s_{1/2}$, (46 keV, 1^+ in ^{34}Al) $\otimes \nu s_{1/2} + (46 \text{ keV}, 1^+$ in ^{34}Al) $\otimes \nu d_{5/2}$. $T_{1/2}$: weighted average (normalized-residual method) of 38.4 ms 3 (2017Ha23, implant- β correlated decay curve); 36.8 ms 5 (2005Ti11, implant- β correlated decay curve, 36.4 ms 5 in 2006AnZW); 38.6 ms 4 (weighted average by 2001Nu01 from four independent measurements, three β -decays and one γ -decay). Others: 30 ms 4 (1995ReZZ, 2008ReZZ, β -decay); 170 ms +90-50 (1989Le16, β -decay); 30 ms 10 (1988DuZT, 1987DuZU); 130 ms +100-50 (1988Mu08, β -decay). Value is 37.6 ms 14 in 2015Bi05 evaluation, which did include the 2017Ha23 result. Reduced strong absorption radius = 1.188 fm ² 14 from 2006Kh08.
802 3			AB	
1007 4			B D	XREF: D(1020)
1866 4			B	
1975 4			B	
2734 7			B	
3245 5			B	
4275? 9			B	XREF: B(4275?)

[†] From a least-squares fit to γ -ray energies. $\gamma(^{35}\text{Al})$

$E_i(\text{level})$	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Comments
802	802 4	100	0	$(5/2)^+$	$B(E2) = 0.0142 \text{ 52}$ (1999Ib01). $B(E1) \leq 0.00020 \text{ 9}$, $B(E2) \leq 0.0125 \text{ 56}$, $B(M1) \leq 0.0024 \text{ 11}$, and $5/2^+ \rightarrow 3/2^{+/-}$ $\sigma = 30 \text{ mb 14}$ (2000PrZX). The multipolarities are assumed. E_γ : weighted average of 1003 4 from $(^{36}\text{Si}, ^{35}\text{Al})\gamma$ and 1020 9 from Coulomb excitation.
1007	1006 6	100	0	$(5/2)^+$	
1866	859 4	100 8	1007		
	1064 4	22 6	802		
1975	968 4	59 4	1007		
	1174 5	37 4	802		
	1972 6	100 7	0	$(5/2)^+$	
2734	1932 6	100	802		
3245	2237 6	100 8	1007		
	2440 7	18.0 26	802		
	3250 8	42 5	0	$(5/2)^+$	
4275?	4275 9	100	0	$(5/2)^+$	

[†] From $^9\text{Be}(^{36}\text{Si}, ^{35}\text{Al})\gamma$, unless otherwise noted.

Adopted Levels, GammasLevel Scheme

Intensities: Relative photon branching from each level

