
 $^2\text{H}(^{34}\text{Si}, ^{35}\text{P}\gamma)$ **2007GeZX**

$^{34}\text{Si}(\text{d}, \text{n})^{35}\text{P}$ from $J^\pi=0^+$ ^{34}Si g.s. in inverse kinematics.

2007GeZX: 30-AMeV ^{34}Si beam on 30-mg/cm² CD₂ secondary target at GANIL. Heavy ions produced in reactions were identified by the VAMOS spectrometer. γ rays were detected using the EXOGAM germanium clover array. Measured Doppler-corrected E_γ , I_γ , $\gamma\gamma$ -coin, and $(^{35}\text{P})\gamma$ -coin. Deduced levels, J , π . Compared with shell-model calculations.

 ^{35}P Levels $E(\text{level})$

0
2386.5 8
3859.9 8
4100.9 13
4492.9 16
4868.9 13

 $\gamma(^{35}\text{P})$

| <u>E_γ^\dagger</u> | <u>$E_i(\text{level})$</u> | <u>E_f</u> |
|--------------------------------------|---------------------------------------|-------------------------|
| 241 | 4100.9 | 3859.9 |
| 392 | 4492.9 | 4100.9 |
| 1009 | 4868.9 | 3859.9 |
| 1473 ‡ | 3859.9 | 2386.5 |
| 2386 | 2386.5 | 0 |
| 3860 | 3859.9 | 0 |

† From **2007GeZX**.

‡ Placement of transition in the level scheme is uncertain.

$^2\text{H}(^{34}\text{Si}, ^{35}\text{P}\gamma)$ 2007GeZX

Legend

Level Scheme-----► γ Decay (Uncertain)