

$^{36}\text{Ar}(\text{pol d}, ^3\text{He})$ 1993Ma50

$J^\pi=0^+$ for ^{36}Ar ground state.

1993Ma50: a 52-MeV polarized deuteron beam at 15 nA was produced from the Karlsruhe isochronous cyclotron at the Max-Planck-Institut für Kernphysik. The target was 350-mbar 99.86% enriched ^{36}Ar gas. Reaction products were detected using a Si detector telescope consisting of a 250- μm ΔE -strip detector and a 1.5-mm surface-barrier E detector with FWHM=130 keV. Measured angular distributions of cross sections $\sigma(E(^3\text{He}), \theta)$ and angular distributions of analyzing powers $iT_{11}(E(^3\text{He}), \theta)$. Deduced levels, J , π , L-transfers, and spectroscopic factors from DWBA analysis of $\sigma(E(^3\text{He}), \theta)$ and $iT_{11}(E(^3\text{He}), \theta)$.

1993Ma50 states that spectroscopic factors were obtained by fitting the forward-angle maxima. The agreement with the analyzing powers is only qualitative. The determination of spins and parities of final states is mainly based on a comparison with measured angular distributions leading to final states with known spins. The comparison with DWBA predictions serves as additional check. The data were taken only in a relatively small angle interval. This is sufficient for the spin determination but not favorable for the measurement of spectroscopic factors.

 ^{35}Cl Levels

Spectroscopic factor $C^2S = \sigma(\theta)_{\text{exp}} / \sigma(\theta)_{\text{DWBA}} / N$, where $N=2.95$ is a normalization factor used by 1993Ma50 from 1966Ba54.

$E(\text{level})^\dagger$	$J^\pi^\ddagger$	$L^\#$	$C^2S^\#$	Comments
0	$3/2^+ @$	2	1.72 @	
1213 4	$1/2^+$	0	0.78	
1785 50	$(5/2^+) \&$		0.04 ^a	
2676 7	$(3/2^+) \&$		0.16 ^a	
3002 6	$5/2^+$	2	1.32	
3159 20	$7/2^-$	3	0.19	
3948 40	$(3/2^+, 1/2^+) \&$	2	0.03, 0.08	$E(\text{level})$: possible doublet of 3918 and 3968 (1990En08) suggested by Table 2 of 1993Ma50.
4839 12	$(5/2^+, 1/2^+) \&$		0.08, 0.03 ^a	
5155 11	$(5/2^+)$	2	0.12	J^π : $(7/2^-)$ and $5/2^+$ are given in Table 2 of 1993Ma50. Evaluators adopt $(5/2^+)$, based on DWBA fit of the measured $\sigma(\theta)$ and $iT_{11}(\theta)$ in Fig. 3 of 1993Ma50.
5583 13	$5/2^+$	2	0.38	
5720 11	$5/2^+$	2	1.10	
6136 8	$5/2^+$	2	0.63	
6745 12	$(1/2^+, 5/2^+) \&$	(0,2)	0.09, 0.28	
6953 43	$5/2^+$	2	0.49	
7181 60	$5/2^+$	2	0.20	
7565 20	$5/2^+$	2	0.30	
8007 36	$(1/2^-, 5/2^+) \&$	(1,2)	0.16, 0.20	
8204 34	$5/2^+$	2	0.14	
8580 20	$5/2^+ \&$		0.22 ^a	
8921 70	$5/2^+ \&$		0.18 ^a	
9.22×10^3 12	$5/2^+ \&$		0.12 ^a	
9514 20	$(1/2^-, 5/2^+) \&$	(1,2)	0.18, 0.30	
9.82×10^3 12	$(1/2^-, 5/2^+) \&$	(1,2)	0.10, 0.26	
1.25×10^4 5				$1d_{5/2}$ is given for a range of 12.0-13.0 MeV in Fig. 4 of 1993Ma50.

[†] From 1993Ma50.

[‡] $L+1/2$ transfer from analyzing power measurements with $1d_{5/2}$ proton transfer assumed in DWBA calculations, unless otherwise noted.

[#] From DWBA analysis of the measured $\sigma(\theta)$ and $iT_{11}(\theta)$ in 1993Ma50. C^2S value is extracted for corresponding J^π assignment. Where two C^2S values are given, they correspond to respective J^π values.

Continued on next page (footnotes at end of table)

 $^{36}\text{Ar}(\text{pol d}, ^3\text{He})$ [1993Ma50](#) (continued)

 ^{35}Cl Levels (continued)

@ L-1/2 transfer from analyzing power measurements with $1d_{3/2}$ proton transfer assumed in DWBA calculations.

& Assumed by [1993Ma50](#) for extracting spectroscopic factors.

^a No data on $\sigma(E(^3\text{He}), \theta)$, $iT_{11}(E(^3\text{He}), \theta)$, or L-transfer is given.