

^{36}K εp decay (341 ms) 1996II02,1980Es01

Parent: ^{36}K : $E=0$; $J^\pi=2^+$; $T_{1/2}=341$ ms 3; $Q(\varepsilon\text{p})=4307.4$ 3; $\% \varepsilon\text{p}$ decay=0.0510

^{36}K - J^π , $T_{1/2}$: From the Adopted Levels of ^{36}K (2012Ni01).

^{36}K - $Q(\varepsilon\text{p})$: From 2021Wa16.

^{36}K - $\% \varepsilon\text{p}$ decay: $\% \varepsilon+\beta^+\text{p}=5.1\times 10^{-4}$ is scaled by evaluators from $\% \varepsilon+\beta^+\text{p}=4.9\times 10^{-4}$ reported in 1996II02. $\% \varepsilon+\beta^+\text{p}=4.9\times 10^{-4}$ in 1996II02 was normalized based on the strongest 970-keV $\varepsilon+\beta^+$ -delayed proton $I_{\text{p}}=3.3\times 10^{-4}$ 9 measured by 1980Es01. 1980Es01 reported $\% \varepsilon+\beta^+\text{p}=4.8\times 10^{-4}$ and their I_{p} was normalized based on the strongest 1970-keV $\varepsilon+\beta^+$ -delayed γ ray $\% I_{\gamma}=79$ 8 measured by 1972Mi13. Evaluators adopt a more precise $\% I_{\gamma}(1970)=81.8$ 16 from the evaluated ^{36}K ε decay dataset (2012Ni01,1976Fr03) as the new normalization reference, and therefore, all original I_{p} values from 1996II02 and 1980Es01 are scaled by 1.04 11.

1996II02,1997II03: A 500-MeV proton beam was produced from the TRIUMF cyclotron and bombarded a CaO powder target. The spallation reaction products diffused out of the target and were selectively ionized at a Re surface. $A=36$ singly charged ions were selected by a magnetic mass analyzer and implanted into a carbon foil at 7.0×10^5 pps. Protons and α particles were detected using a 30- μm silicon surface barrier detector with FWHM=23 keV. ^{32}S and ^{35}Cl recoils were detected using an MCP in a back-to-back geometry for timing stop signals. Measured $E_{\text{p}}(>0.5$ MeV) and I_{p} for 15 proton peaks, $E_{\alpha}(>1.5$ MeV) and I_{α} for 19 α peaks, p(recoil)-coin, and α (recoil)-coin. Deduced levels, J , π , decay branching ratios, $\log ft$, partial decay widths for ^{36}Ar , and ratios of $^{35}\text{Cl}(\text{p},\alpha)^{32}\text{S}$ to $^{35}\text{Cl}(\text{p},\gamma)^{36}\text{Ar}$ reaction rates.

1980Es01: A 20-MeV proton beam was produced from the University of Jyväskylä MC-20 cyclotron and bombarded ^{36}Ar -deposited Al foils. The $^{36}\text{Ar}(\text{p},\text{n})$ reaction products were thermalized by helium, transported with NaCl-loaded helium into a measurement chamber, and deposited onto a fast-transport Al-coated mylar tape. Protons and α particles were detected using 14- and 26- μm fully depleted (FWHM=40 keV) and 100- μm partially depleted Si(Au) surface barrier detectors. A polyethylene absorber foil was placed in front of Si(Au) to cause an energy shift for distinguishing p/ α . Measured $E_{\text{p}}(>0.5$ MeV) and I_{p} for 11 proton peaks, $E_{\alpha}(>2.0$ MeV) and I_{α} for 12 α peaks. Deduced levels, J , π , decay branching ratios, $\log ft$, and partial decay widths for ^{36}Ar .

1980Ew01: A 600-MeV proton beam was produced from the synchrocyclotron at CERN-ISOLDE and bombarded a ScC_2 target. The $^{45}\text{Sc}(\text{p},7\text{n}3\text{p})$ spallation reaction products diffused out of the target and reached a tungsten surface ionization source where potassium isotopes were selectively ionized. The beam was extracted from the ion source, separated by the ISOLDE analyzing magnet, and collected by a carbon foil for proton measurements. Protons and α particles were detected using 20- μm and 100- μm silicon surface barrier detectors with FWHM=35 and 23 keV. Measured $E_{\text{p}}(>0.5$ MeV) and I_{p} for 3 proton peaks, E_{α} and I_{α} for 4 α peaks. Deduced levels, J , π , and decay branching ratios for ^{36}Ar .

 ^{35}Cl Levels

<u>E(level)</u>	<u>J^π[†]</u>
0	$3/2^+$

[†] From the Adopted Levels.

Delayed Protons (^{35}Cl)

<u>E(p)[†]</u>	<u>E(^{35}Cl)</u>	<u>I(p)^{†@}</u>	<u>E(^{36}Ar)[#]</u>	<u>Comments</u>
501.8 11	0	1.14×10^{-6} 33	9023	E(p): other: 501 10 (1980Es01). I(p): other: 3.6×10^{-6} 13 (1980Es01).
622.7 14	0	1.6×10^{-6} 5	9147	
693.5 8	0	7.9×10^{-5} 22	9220	E(p): other: 693 5 (1980Es01), 691 8 (1980Ew01). I(p): other: 7.8×10^{-5} 24 (1980Es01).
851.7 10	0	2.3×10^{-5} 7	9383	E(p): other: 849 5 (1980Es01). I(p): other: 2.0×10^{-5} 7 (1980Es01).
969.6 12	0	3.4×10^{-4} 10	9504	E(p): other: 970 5 (1980Es01), 959 7 (1980Ew01). I(p): from 1980Es01 and also adopted as the normalization reference by 1996II02.
1168.5 14	0	6.2×10^{-7} 7	9709	

Continued on next page (footnotes at end of table)

^{36}K ε p decay (341 ms) 1996II02,1980Es01 (continued)Delayed Protons (continued)

$E(p)^{\dagger}$	$E(^{35}\text{Cl})$	$I(p)^{\dagger @}$	$E(^{36}\text{Ar})^{\#}$	Comments
1198.5 11	0	2.5×10^{-6} 7	9740	
1272.0 20	0	1.6×10^{-6} 5	9815	
1334.2 7	0	4.5×10^{-5} 12	9879	E(p): other: 1333 5 (1980Es01), 1325 7 (1980Ew01). I(p): other: 3.6×10^{-5} 11 (1980Es01).
1409.3 22	0	1.04×10^{-6} 33	9957	
1874 ‡ 10	0	1.1×10^{-6} 5	10435	
1992 ‡ 10	0	5.0×10^{-6} 20	10556	
2048 ‡ 10	0	4.9×10^{-6} 19	10613	
2458 ‡ 10	0	3.0×10^{-6} 13	11035	
2640 ‡ 10	0	2.1×10^{-6} 10	11222	

† In lab frame. From 1996II02, unless otherwise noted.

‡ From 1980Es01 and also adopted by 1996II02.

$^{\#}$ $E(\text{level})(^{36}\text{Ar}) = E(p)_{\text{lab}} \times [m(p) + m(^{35}\text{Cl})] / m(^{35}\text{Cl}) + S(p)(^{36}\text{Ar}) + E(\text{level})(^{35}\text{Cl})$, where $S(p)(^{36}\text{Ar}) = 8506.97$ 4 (2021Wa16). Proton decay is assumed to populate only the ground state of the daughter nucleus ^{35}Cl based on the arguments proposed in 1996II02: (A) unreasonably small $\log ft$ values if populating excited states; (B) no pairs of particle groups characteristic of the g.s./first excited state energy difference; (C) compatible level matching with resonance reaction studies.

$^@$ Absolute intensity per 100 decays.

^{36}K εp decay (341 ms) 1996II02,1980Es01**Decay Scheme**

I(p) Intensities: I(p) per 100 parent decays

