

$^{160}\text{Gd}(^{34}\text{S},\text{x}\gamma),(^{37}\text{Cl},\text{X}\gamma)$ 1994Fo04

1994Fo04: 159-MeV ^{34}S and 167-MeV ^{37}Cl beams were produced from the Argonne Tandem Linac Accelerator System (ATLAS). Targets were 1 mg/cm² 98.1% enriched ^{160}Gd backed by 15 mg/cm² gold. γ rays were detected using the Argonne-Notre Dame BGO γ -ray facility consisting of 12 Compton-suppressed Ge detectors and a 50-element bismuth germanate (BGO) array. Measured $E\gamma$, $I\gamma$, $\gamma\gamma$ -coin. Deduced levels. $\gamma\gamma$ -coin gates were placed on known γ rays in specific $A\approx 160$ products to select individual reaction channels and identify coincident γ rays in light product partners. $\gamma\gamma$ -correlation matrices produced with the fold condition $4 \leq k \leq 9$ were found to include most of the events associated with deep inelastic processes.

 ^{35}S Levels

<u>$E(\text{level})^\dagger$</u>	<u>$J^\pi^\ddagger$</u>	Comments
0	$3/2^+$	
1991	$7/2^-$	
4022	$(9/2^-)$	J^π : $(11/2)^-$ from the Adopted Levels.

[†] From $E\gamma$ data in 1994Fo04.

[‡] As given in 1994Fo04.

 $\gamma(^{35}\text{S})$

<u>$E_\gamma^\dagger$</u>	<u>$E_i(\text{level})$</u>	<u>J_i^π</u>	<u>E_f</u>	<u>J_f^π</u>
1991	1991	$7/2^-$	0	$3/2^+$
2031	4022	$(9/2^-)$	1991	$7/2^-$

[†] From 1994Fo04.

 $^{160}\text{Gd}(^{34}\text{S},\text{x}\gamma),(^{37}\text{Cl},\text{X}\gamma)$ 1994Fo04Level Scheme