

# High Dimensional Data Analysis With Dependency and Under Limited Memory

Yiming Sun

Cornell University

Based on joint work with

Madeleine Udell (Cornell), Sumanta Basu(Cornell)

Yang Guo (UW Madison), Charlene Luo (Columbia)

Joel Tropp (Caltech), Amy Kuceyeski(Cornell University)

Yige Li(Harvard University)

September 8, 2019


# Outline

Multivariate Spectral Density Estimation Under Weak Sparsity

Low Rank Tucker Approximation of a Tensor from Streaming Data

A Connection of These Two

# Motivation



**Figure:** Interactions Between Regions in Brain

## Weakly & Strongly Stationary Time

### Definition (Weak Stationarity)

$p$ -variate time series  $X$  is weakly stationary, if  $\mathbb{E}X_t = \mathbb{E}X_s$  for any  $t, s$  and  $\Gamma(\ell) := \mathbb{E}X_t X_{t-\ell}^\top$  only depends on the lag  $\ell$ .

### Definition (Strong Stationarity)

$p$ -variate time series  $X$  is strongly stationary, if for any sequence  $t_1, \dots, t_n$ ,  $X_{t_1} \cdots X_{t_n}$  has the same distribution of  $X_{t_1+\tau} \cdots X_{t_n+\tau}$  for any integer  $\tau$ ,

# Gaussian Process

## Definition (Gaussian Process)

$p$ -variate time series  $X$  is Gaussian process if for any sequence  $t_1, \dots, t_n$ ,  $X_{t_1} \dots X_{t_n}$  are jointly Gaussian distributed.

For Gaussian process, weak stationarity is equivalent to strong stationarity.

## Spectral Density

Given a weakly stationary  $p$ -variate time series  $X$ , the spectral density at frequency  $\omega \in [-\pi, \pi)$  is defined

$$f(\omega) = \sum_{\ell=-\infty}^{\infty} \Gamma(\ell) e^{-i\omega\ell}$$

where  $\Gamma(\ell) = \mathbb{E}X_0X_{-\ell}^\top$ .  $X_t$  is independent with  $X_s$ ,  $t \neq s$  iff  $f_{rs}(\omega) = 0$  for any  $\omega$ .

# Thresholding Estimator Under Weak Sparsity- A Example

Suppose that we have  $n$  observation of  $p$ -variate Gaussian distribution as follows.

$$y_i \stackrel{i.i.d}{\sim} \mathcal{N} \left( \mu, \begin{bmatrix} \sigma_1^2 & 0 & \cdots & 0 \\ 0 & \sigma_2^2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \\ 0 & 0 & 0 & \sigma_p^2 \end{bmatrix} \right),$$

$$i = 1, \dots, n.$$

## A Example

The maximum likelihood estimator for  $\mu_j$  is  $\bar{y}_j = \frac{1}{n} \sum_{i=1}^n y_{ij}$ .

**Does not Work Well Under Weak Sparsity**

$$\mu \in \left\{ \mu \in \mathbb{R}^p, \sum_{j=1}^p |\mu_j|^q \leq c_0(p) \right\}.$$

for some  $0 \leq q < 1$  and  $c_0(p)$  measures the weak sparsity.



## Solution : Thresholding

Suppose  $\sigma_i \leq B$ , define element-wise thresholding operator

$$T_\lambda(x) = \begin{cases} x & |x| \geq \lambda \\ 0 & \text{else} \end{cases}$$

. Hard thresholding estimator  $T_\lambda(\bar{y}_j)$  can be shown asymptotically consistent under weak sparsity where we set

$$\lambda \propto B \sqrt{\frac{\log p}{n}}.$$

# Outline

Multivariate Spectral Density Estimation Under Weak Sparsity

Low Rank Tucker Approximation of a Tensor from Streaming Data

A Connection of These Two

# Outline

Multivariate Spectral Density Estimation Under Weak Sparsity

Low Rank Tucker Approximation of a Tensor from Streaming Data

A Connection of These Two