

Electronic supplement/Appendix C: Parameters used for calculating fluid compositions in figure 4 to 6.

	DM ¹	SubdSed ^{2,3}	N-MORB	Dcpx/fluid	Dgrt/fluid	Drutile/fluid
Mo	0.0250	5.49 ³	0.408 ⁴	$40/10^{(0.435 \cdot \log fO_2 + 0.42 \cdot \log NaCl - 1.8 \cdot 1000/T + 4.8) \cdot}$	$12/10^{(0.435 \cdot \log fO_2 + 0.42 \cdot \log NaCl - 1.8 \cdot 1000/T + 4.8) \cdot}$	$87670/10^{(0.435 \cdot \log fO_2 + 0.42 \cdot \log NaCl - 1.8 \cdot 1000/T + 4.8) \cdot}$
Ce	0.7720	57.3	12.0 ⁵	2.000 ⁷	0.400 ⁷	2.000 ⁷
W	0.0024 ⁶	1.69 ³	0.038 ⁶	$60/10^{(0.07 \cdot \log fO_2 - 4.7236 \cdot 1000/T + 4.4271) \cdot}$	$12/10^{(0.07 \cdot \log fO_2 - 4.7236 \cdot 1000/T + 4.4271) \cdot}$	1.250
U	0.0047	1.68	0.0711 ⁵	$11/10^{(2.681 + 0.1433 \log fO_2 + 0.594 Cl(mol)) \cdot}$ ⁸	$40/10^{(2.681 + 0.1433 \log fO_2 + 0.594 Cl(mol)) \cdot}$ ⁸	$94/10^{(1.7954 + 0.1433 \log fO_2 + 0.594 Cl(mol)) \cdot}$ ⁸⁻⁹
Th	0.0137	6.91	0.1871 ⁵	1.190 ⁸	0.610 ⁸	0.100 ⁹
Nb	0.2100	8.94	3.507 ⁵	0.172 ⁷	0.204 ⁷	200.0 ⁷
La	0.2340	28.8	3.895 ⁵	1.429 ⁷	0.204 ⁷	1.250 ⁷

1 - Depleted mantle - Salters and Stracke (2004)

2 - Subducted sediment - Plank and Langmuir (1998)

3 - Subducted sediment - calculated following Plank and Langmuir (1998) as 76wt% terrigenous sediment (Upper Continental Crust - Rudnick and Gao, 2004)+17wt% of pelagic sediment (calculated as pelagic clay - Li, 1991)+7wt% of mineral bound water (assumed not to contain W and Mo)

4 - Sun et al. (2003)

5 - Hart et al. (1999)

6 - König et al. (2011)

7 - Stalder et al. (1998)

8 - Bali et al. (2011)

9 - Brenan et al. (1994)

8-9 - The fO_2 and salinity dependence of fluid/rutile partition coefficient was estimated based on the U-contents in U-saturated experiments of Brenan et al. (1994) combined with the U-saturated fluid compositions determined for the relevant pressure based on Bali et al. (2011). According to Stalder et al. (1998) fluid/rutile partition coefficients are independent of pressure.

*formula in nominator gives concentrations in molality (see text) which should be recalculated to ppm!

References:

- Bali, E., Audetat, A., Keppler, H., 2011. The mobility of U and Th in subduction zone fluids – an indicator of oxygen fugacity and fluid salinity. *Contrib. Mineral. Petrol.*, 161, 597-613.
- Brenan, J.M., Shaw, H.F., Phinney, D.L., Ryerson, F.J. 1994. Rutile-aqueous fluid partitioning of Nb, Ta, Hf, Zr, U and Th: implications for high field strength element depletions in island-arc basalts. *Earth Planet. Sci. Lett.*, 128, 327-339.
- Hart, S.R., Blusztajn, J., Dick, H.J.B., Meyer, P.S., Muehlenbachs, K. 1999. The fingerprint of seawater circulation in a 500-meter section of ocean crust gabbros. *Geochim Cosmochim Acta* 63, 4059-4080.
- König, S., Münker C, Hohl, S, Paulick, H, Barth, AR, Lagos, M, Pfander, J, Bühl, A (2011) The Earth's tungsten budget during mantle melting and crust formation. *Geochim Cosmochim Acta*, 75, 2119-2136.
- Li, Y.H., 1991. Distribution Patterns of the Elements in the Ocean - a Synthesis. *Geochim Cosmochim Acta* 55, 3223-3240.
- Plank, T., Langmuir, C.H., 1998. The geochemical composition of subducting sediment and its consequences for the crust and mantle. *Chem Geol* 145, 325-394.
- Salter, V., Stracke, A., 2004. Composition of the depleted mantle. *G-Cubed*, 5, doi: 10.1029/2003GC000597, Q05B07, 27 PP.
- Stalder, R., Foley, S.F., Brey, G.P., Horn, I., 1998. Mineral aqueous fluid partitioning of trace elements at 900-1200°C and 3.0-5.7 GPa: New experimental data for garnet, clinopyroxene, and rutile, and implications for mantle metasomatism. *Geochim Cosmochim Acta*, 62, 1781-1801.
- Sun, W., Bennett, V.C., Eggins, S.M., Arculus, R.J., Perfit, M.R. 2003. Rhenium systematics in submarine MORB and back arc basin glasses: laser ablation ICPMS results. *Chem Geol*, 196, 259-281.