

ChemWorld: Composable Chemical Worlds for Controlled and Replayable Agent Experimentation

Jiangjie Qiu¹, Yijun Li¹

¹Beijing Key Laboratory of Artificial Intelligence for Advanced Chemical Engineering Materials, State Key Laboratory of Chemical Engineering and Low-Carbon Technology, Department of Chemical Engineering, Tsinghua University, Beijing 100084, China

Abstract

Autonomous chemistry needs both real-material execution and a software-scale experimental medium in which worlds can be reset exactly, repeated safely, observed completely and changed under controlled laws. ChemWorld provides this second capability as a composable executable chemical-world substrate and programmable virtual instrument. Reusable physical and transactional components compile into worlds, while one public task contract joins initial state, typed operations, instruments, observations, resources, failure semantics, termination and evaluation. Researchers can therefore create matched worlds, branch one private law at a time and audit every simulator-side state and resource event without directly consuming physical reagents or creating wet-laboratory chemical exposure. The 15 registered tasks serve as reference examples for a broader construction surface. Qualification covered 64/64 reference world units, 1,786/1,786 complete reference recipes and 52/52 coverage-generated compositions, including eight protocol-frozen non-reference reaction–distillation compositions. All 192 invalid-action probes, 32 module probes, seven interface paths and seven invalid declarations produced their registered outcomes, with zero missing receipts or public/private leakage. Across eight deterministic use cases, all 89 submitted actions were audited; one planned failure rolled back without ghost state, every lifecycle and resource ledger closed, and every trajectory replayed with zero numerical mismatch. Six controlled private-law fork pairs preserved the public contract across 24 traces, and one complete agent closed the first protocol-frozen non-reference world through the same instrument interface. Together, these results show that compositional expansion can preserve executable semantics and turn simulated chemistry from a fixed task collection into a controlled, observable and replayable experimental medium. The released v1 instrument spans nine declared component families and defines extension points for future alternative or empirically calibrated formulations.

1. Introduction

Self-driving laboratories (SDLs) and chemistry agents have demonstrated that algorithms can plan, execute and revise workflows on real materials. A second experimental regime is needed for questions that depend on exact reset, matched counterfactuals, repeated failure injection, complete process observation and direct control over the laws of the world. These operations are difficult to obtain from physical matter and are usually absent when a digital experiment is represented only as an objective-function query. A programmable virtual instrument can make them routine at software scale.

ChemWorld treats a chemical world as an explicit executable object rather than an opaque task label. Its v1 vocabulary contains reaction, thermal, phase, separation, crystallization, distillation, continuous-flow, electrochemical and observation components. A world is a compatible selection of these components, their parameters and private laws. A task contract is $T = (W, S_0, A, I, O, R, \tau, E)$: the world, initial state, operations, instruments, observations, resources, termination rule and evaluation surface. A scenario instantiates the contract; a trajectory records the resulting operation–observation sequence; and a controlled fork changes one private component while preserving the public contract.

This architecture creates controlled experimental freedom. ChemWorld can run wherever the software and compute environment are available. The software itself

consumes no physical reagents and creates no direct wet-laboratory chemical exposure. World state can be reset exactly, repeated to a planned study denominator under explicit compute and storage budgets, and forked at a named private law. The evaluator can inspect every hidden state transition and resource event while the agent remains restricted to the public instrument contract. These properties make high-observability counterfactual experimentation available before, alongside or between physical campaigns (Fig. 1).

The central question is whether a chemical environment can expand beyond a fixed task list without losing the semantics that make experiments interpretable. We address it through four linked contributions:

1. **Composable executable worlds.** A public component vocabulary, compatibility compiler and unified task contract convert component selections and private laws into executable, reader-auditable instruments.
2. **Coverage-guided expansion.** Pairwise discrete coverage, seeded space-filling continuous designs and ordered workflow interactions extend qualification beyond the reference task identities.
3. **Process-complete semantics.** Atomic transactions, explicit failure and termination, multi-resource ledgers, public/private observation boundaries and exact environment replay make each trajectory an inspectable experimental record.

4. **Controlled intervention and agent access.** Single-private-law forks isolate trace-level world changes under an invariant public contract, fixed typed actions and bound randomness, while deterministic policies and a complete agent use the same instrument surface.

This paper qualifies the declared v1 component and compatibility domain as a virtual instrument. Its modular interfaces separate experimental semantics from model choice and define extension points through which future implementations may introduce additional constitutive laws, calibrated process models or alternative agents.

2. Relation to existing experimental systems

2.1 Real-material execution and software-scale control

Physical autonomous laboratories provide the decisive evidence of execution on real materials with real sensors, actuators and hardware. Coscientist and ChemCrow connect language-model planning to chemistry tools and robotic or cloud-laboratory execution [1, 2]. A-Lab and mobile robot systems demonstrate closed-loop synthesis and characterization [3, 4], while ORGANA, ChemAgents and newer instrument-facing systems extend this line toward visual feedback, long workflows, modular automation and teachable operation [5–9].

ChemWorld supplies the complementary evidence available from a fully controlled software world: exact matched replication, counterfactual private laws, evaluator-complete state access, deterministic reset and repeatable failure injection. Its marginal resources are compute and software rather than reagent, instrument and human-safety capacity. This division enables large controlled studies in software and focused real-material validation where physical evidence is decisive.

2.2 Digital optimization, virtual laboratories and discovery worlds

Reaction-optimization and experiment-planning suites such as Summit and Olympus provide scalable, repeatable comparisons over objective functions, while PC-Gym provides nonlinear process-control environments with constraints and disturbances [10–12]. ChemGymRL already establishes a customizable, fine-grained virtual chemistry laboratory for reinforcement learning [13]. Closed-loop materials environments further couple candidate generation, budgeted oracle feedback and multi-objective search [14, 15]. These systems establish the value of inexpensive repeatability, interactive chemistry and controlled algorithm comparison.

Interactive scientific worlds broaden evaluation toward hypothesis formation, experiment selection and law recovery. DiscoveryWorld, BoxingGym, SciGym, SciExplorer and NewtonBench represent complementary approaches to long-horizon discovery and initially unknown systems

[16–20]. Laboratory simulators such as LabUtopia and Labimus emphasize embodied procedure and manipulation [21, 22]. ChemWorld adds a distinct systems layer: world composition, instrumentation, private-law intervention and replay are governed by one executable contract. Typed operations change sample state, instruments consume explicit resources, invalid actions have transactional consequences, termination is explicit, and the complete world transition can be rebuilt from the released record. This combination supports experiments about both the agent and the world it inhabits.

The construction and record layers also connect to established software-systems ideas. Pairwise covering rows follow combinatorial interaction testing, which selects compact test suites that cover registered parameter interactions [23]. The immutable record specializes computational provenance—entities, activities, derivations and responsible agents—to executable chemical-world transitions, resources and lineage [24]. ChemWorld joins these ideas to a stateful scientific instrument rather than treating them as release metadata alone.

2.3 A distinct operating regime: programmable experimental freedom

The defining ChemWorld advantage is the conjunction of composition, controlled intervention and process-complete observation. Researchers can instantiate new component topologies, vary continuous conditions, substitute a private constitutive or material law, repeat matched world identities and inspect every simulator-side consequence. The same contract also preserves the agent-facing boundary, so counterfactual worlds remain comparable without exposing their hidden differences. Table 1 positions this capability alongside the strongest use of adjacent experimental systems.

3. Composing executable worlds beyond a fixed task catalogue

3.1 One contract from world construction to experiment

Each component declares the state it owns, the interfaces it consumes or produces, its parameter domains and its contribution to the public operating surface. Reaction and electrochemical components transform material state; thermal, phase, separation, crystallization, distillation and continuous-flow components supply process-specific state transitions; the observation component attaches synthetic instruments. Public operations are typed actions rather than free-form simulator mutations.

The compiler first normalizes the declaration and then checks compatibility. It fails closed on missing dependencies, multiple owners of the same state surface, unit incompatibility, unsupported parameters, impossible resource declarations and workflows that cannot reach termination. A successful compile returns both an executable task and a reader-auditable public surface: allowed operations, instruments, observations, resources, termination and evaluation.




Figure 1. ChemWorld turns composable chemical worlds into controlled, auditable experiments. **A**, Physical SDLs provide real-material execution; the virtual substrate adds software access, exact reset, repetition without direct wet-lab exposure, matched counterfactuals and complete simulator observability. **B**, Reusable components and private laws compile through compatibility checks into a world and public task contract. **C**, An agent or deterministic policy acts only through typed operations and instruments until explicit termination and final assay. **D**, The immutable record joins state transitions, observations, resources, failures and lineage, enabling exact environment replay and controlled private-law forks.

Private laws and hidden state remain outside that surface.

The public capability map contains 15 registered reference tasks, 28 typed operation kinds, five synthetic instrument contracts and 62 ordered task-by-metric bindings. The reference tasks anchor interpretable examples across the declared surface, while component composition supplies the expansion mechanism. This separates the size of the executable world space from the number of curated task identities.

3.2 Coverage-guided expansion

Composition generation is driven by coverage targets rather than by a desired number of examples. Discrete component and instrument choices are arranged with pairwise covering rows. Continuous temperature, time, flow, volume, potential, current, reflux and transfer axes are sampled with seeded Latin hypercube designs. Ordered workflows separately require critical interactions such as reaction

before separation, quench before downstream transfer and fraction collection before final measurement.

The protocol-frozen design contains eight component patterns and generated 52 compositions. Comparison of exact component sets against the reference registry separates two forms of expansion. Three patterns—phase-observation, phase-separation-observation and reaction-thermal-continuous-flow-observation—are absent from the reference topologies and contribute 18/52 generated cases. Five patterns reuse a registered component topology. Within the latter group, the eight reaction-thermal-distillation-observation rows have zero exact task-world identity overlap with the frozen registry. We call these *protocol-frozen non-reference compositions*: their topology is registered, but their bound task and world identities are not. The first row of this eight-case block was fixed before authoritative execution as the complete-agent target, linking agent use to the same coverage design rather than to

Table 1. Complementary operating regimes for autonomous chemical experimentation. ChemWorld combines software-scale repetition with explicit world composition, private-law intervention and process-complete replay.

System class	Strongest use	Replication regime		Experimental intervention	Observable record
Physical SDL / chemistry robot	real-material execution and hardware integration	apparatus-, material- and time-bound physical repeats		protocol, material and hardware changes	sensor, automation and sample records
Optimization or control suite	rapid algorithm comparison over objectives or dynamics	software-scale repeated queries or control episodes		configurable objectives, datasets or process models	objective histories or controller traces
Interactive virtual lab / discovery world	sequential discovery or embodied procedure	resettable episodes	software	task-specific objects, procedures and latent rules	action histories, observations and task state
ChemWorld	controlled experiments over composable executable worlds	exact reset, matched repetition and version-bound replay without direct wet-laboratory consumable use		declared component composition plus single-private-law forks under an invariant public contract	typed actions, evaluator-complete state, public observations, failures, resources, termination, lineage and exact environment replay

a separately selected example.

Every registered coverage target was attained. Across the eight patterns, the generated suite covered 60/60 discrete levels, 180/180 compatible discrete pairs, 212/212 continuous strata and 84/84 ordered workflow interactions. These are finite registered targets within the authored v1 domains, and their covered and required counts are retained pattern by pattern in the machine-readable coverage record.

4. Composition preserves physical and experimental semantics

4.1 Reference and generated full censuses

The reference qualification treats a world unit as a task-world pairing rather than a single task label. All 64/64 reference units passed. Boundary and categorical recipes produced 1,786/1,786 complete executions. The generated block passed for all 52/52 compositions, including all eight non-reference reaction-distillation rows. There were no failure classes, missing receipts or public/private leakage findings.

Compilation success alone was not a pass condition. Each generated composition had to execute its complete workflow, close exactly once, reconcile declared and observed resources and replay exactly. Seven deliberately broken declarations tested missing dependencies, conflicting state ownership, unit mismatches, invalid domains, resource impossibility and lifecycle gaps; all seven were rejected before environment construction.

4.2 Physical and cross-module checks

Thirty-two module probes exercise zero input, declared boundaries, monotonic directions, conservation and model-specific invariants. The seven cross-module paths then

check that material amount, unit, identity and state meaning survive transfer between modules. When applicable, the checks also include charge, energy, phase balance and event propagation. All 32 module probes and seven interface paths passed.

These results qualify each formulation as an internally coherent virtual-instrument module inside its model-card domain. Every module enters through declared interfaces that define where future implementations may supply alternative or empirically calibrated formulations while retaining the task contract, transaction layer and replay machinery.

4.3 Transactions, resources, observations and replay

Every submitted action first passes schema, compatibility and resource preflight. A valid action commits atomically. A rejected action records its attempt and declared penalty but does not install candidate physical state. The 192 negative probes cover invalid schema, preconditions, resource exhaustion, terminal closure and other fail-closed paths; all 192 produced the registered rejection and preserved state as required.

Resource accounting separates material, sample, instrument use, process time, operation count and terminal assay. Observation checks ensure that public packets contain only task-declared fields. Exact replay reconstructs the same compiled world from its bound contract and seeds, resubmits the committed typed actions, and compares rewards, observations, termination flags, transaction metadata, state-delta summaries, resource events and constitution checks with zero numerical tolerance. This version-bound guarantee makes the full environment transition, rather than only the endpoint or action list, the reproducible unit.



Figure 2. Coverage-guided construction and full-census qualification. **A**, Three of eight component patterns, representing 18/52 cases, add topologies absent from the reference registry; the eight reaction–distillation rows instead reuse a registered topology while having zero exact task–world identity overlap. **B**, The 52 protocol-frozen rows attain 60/60 registered discrete levels, 180/180 compatible pairs, 212/212 continuous strata and 84/84 ordered workflow interactions. **C**, All reference units, reference recipes, generated compositions and non-reference compositions completed and replayed. **D**, Module, interface, invalid-declaration and invalid-action probes all produced their registered outcomes, with zero missing receipts or public/private leakage. Counts report complete qualification denominators.

5. Process-complete execution across diverse chemical workflows

Eight frozen use cases test the recording surface with deterministic policies. They cover a reaction-to-crystallization workflow, resource-limited equilibrium characterization, an intentional failure followed by recovery, continuous flow, electrochemistry, distillation, partition and a second crystallization world. The cases are independent qualification units, with every submitted action audited inside its complete lifecycle.

Across the eight cases, all 89 submitted actions have complete schema, transaction, constitution, event, resource and public-observation receipts. Eighty-eight actions committed. The first action of the failure–recovery case deliberately attempted phase separation before its preconditions were satisfied. It rolled back, preserving physical state and

observation random-number state while reconciling the declared attempt consequences. The following 18 actions completed the recovery path. Every case committed one final assay, closed its lifecycle, reconciled resources and replayed exactly with zero numerical error.

The generated reaction–distillation world also has a 12-action deterministic reference path. This creates two independent execution units on the same world: a fixed path that qualifies construction and replay, and a complete-agent lifecycle that tests access through the public instrument contract.

6. Controlled single-component forks

General composition and controlled attribution are different operations. Composition assembles multiple declared components subject to compatibility rules. A world fork holds the public task contract and action sequence fixed



Figure 3. Process-complete cases preserve lifecycle, resource and failure semantics. **A**, Eight frozen cases span single-process, multistage and reference-library workflows. **B**, Eighty-eight of 89 submitted actions committed; one protocol-defined precondition failure rolled back. **C**, The rollback preserved physical state and allowed the remaining 18-step recovery path to close. **D**, All eight final assays, resource ledgers and exact replays passed.

while changing one protocol-frozen private component.

The fork qualification contains six parent-child pairs: two intervention classes across three seeds. Each pair preserved nine versioned public-contract components: task, actions, instruments, observations, resources, failures, scoring, material catalogue and constitution/safety. These fork-certificate fields refine the public boundary of the task contract rather than redefining its mathematical tuple. Parent and child executed the same fixed typed sequence with bound randomness. Repeating both variants produced 24 deterministic fork traces. All pairs passed lineage, single-target, public-invariance, same-sequence executability, expected state and observation divergence, and exact replay gates. The measured divergence is therefore a trace-level effect of one private-law intervention under fixed actions and noise identity.

7. Agent experimentation with evaluator-complete observability

An endpoint cannot reconstruct how it was reached. ChemWorld therefore keeps terminal commitment, evidence acquisition, continued process investment, resource deployment, failure and outcome trajectory as separate coordinates. Nineteen registered process dimensions remain separate, preserving the experimental structure that a scalar score would discard. The evaluator can inspect complete simulator state and resource consequences while the agent remains restricted to its public task contract.

The complete-agent demonstration used the first protocol-frozen non-reference reaction-distillation world. It ran on 5 August 2026 with OpenAI GPT-5.6-sol at medium reasoning effort through the Codex subscription provider. The fixed one-turn scaffold supplied the public task card, typed tool schemas, resource contract and explicit termination/final-assay requirement. The agent received only this public surface and issued every operation,

including termination and final assay. One uninterrupted session submitted 15 actions under the same model and scaffold. All 15 committed and closed the lifecycle, with zero rollback, right-censoring or public/private leakage.

The environment used 8,158.454 of 10,440 simulated process seconds, four of four instrument uses and 0.00085 of 0.001 L sample. The complete record links each decision to its public observation, hidden simulator consequence, transaction result and resource debit. The 15-step environment trajectory then replayed with zero numerical mismatch, showing that a provider-driven experiment can enter the same auditable record as deterministic use cases.

8. Discussion

8.1 Controlled experimental freedom as a scientific capability

ChemWorld turns world construction into an experimental variable. Declared components compile through explicit compatibility rules; coverage-guided rows extend beyond curated task identities; and reference, generated and non-reference compositions preserve physical, transactional, resource, observation and replay semantics. The result is a construction surface rather than a catalogue of isolated environments.

Six capabilities become available together: qualified new component topologies, continuous condition variation, private-law intervention, exact reset, matched repetition and evaluator-complete process inspection. Researchers can exercise them without booking physical hardware or creating direct wet-laboratory chemical exposure, and can repeat an experiment to its planned denominator under explicit compute and storage budgets. This makes large controlled studies, diagnostic stress tests and counterfactual designs economically accessible at software scale.

The full-census qualification is central to this advantage. Every registered unit carries an exact denominator and every unexpected outcome remains visible. Coherent semantics are therefore established across the declared construction domain before a policy comparison, controlled intervention study or scientific application is layered on top.

8.2 From endpoint benchmarks to controlled counterfactual process analysis

An endpoint reports what was achieved; a process-complete record reveals how evidence, resources, failures and terminal commitment produced it. ChemWorld retains these signals as 19 separate coordinates and binds them to replayable state transitions. This supports questions about measurement strategy, recovery after invalid actions, resource deployment and decision timing that disappear when an experiment is reduced to one score.

Controlled private-law forks add a counterfactual dimension. Parent and child worlds expose the same actions, instruments and resources while one hidden law changes under the same typed sequence and bound randomness. Outcome and observation differences in the reported traces

can therefore be assigned to that simulator intervention. The same construction surface can support future policy-robustness, adaptation, measurement-value, curriculum and failure-recovery studies through separately frozen randomized or matched protocols.

8.3 An extensible bridge to physical experimentation

The released v1 qualification applies to the declared component vocabulary and authored model domains. Those boundaries expose explicit extension points rather than fixed task walls. Future implementations may introduce additional constitutive laws, calibrated unit operations, empirical instrument models and new agents through the same task contract. The complete-agent unit demonstrates that provider-driven action selection already uses this common surface, while comparative agent studies can retain the same environment, resource and replay guarantees.

ChemWorld and physical SDLs therefore form a productive sequence. Software worlds provide cheap repetition, controlled counterfactuals, complete observability and rapid hypothesis narrowing; physical systems provide real-material execution and calibration. The shared experimental logic allows software-scale evidence to focus subsequent laboratory work on the conditions and mechanisms that matter most.

9. Methods

9.1 Construction and compatibility

A composition declaration specifies component roles and parameters plus the public task surface. Normalization is deterministic. Compatibility checks cover dependencies, state ownership, unit agreement, parameter domains, resource feasibility, operation exposure, instrument availability and the existence of a closed lifecycle. A rejected declaration returns structured diagnostics and does not construct an environment.

The task contract is the authoritative public boundary. Runtime validation is restricted to the declared operations and instruments. Private constitutive laws, material identities and hidden simulator state remain evaluator-owned. The 15 registered tasks are mapped into the same component and contract representation used by generated compositions.

9.2 Coverage design

The authoritative qualification protocol was frozen before execution. The frozen object includes eight component patterns, their discrete axes and compatible pairs, continuous bounds, seeds, ordered workflows, pass/failure rules and the first reaction-distillation row selected for complete-agent use. Discrete axes use pairwise covering rows. Continuous axes use seeded Latin hypercube samples inside the authored bounds. Each pattern contains one or two ordered workflows. The denominator is 52 generated cases, including 18 cases across three topologies absent from the reference registry and eight registered-topology




Figure 4. Controlled private-law interventions and complete-agent use of the same instrument surface. **A**, Parent and child worlds share the public contract and action sequence while one private constitutive or material law changes. **B**, Six pairs and 24 deterministic traces pass lineage, public-invariance, divergence and exact-replay gates. **C**, The fixed non-reference reaction–distillation world has an independent 12-step deterministic qualification path and a separate 15-step complete-agent lifecycle. **D**, The agent closes the lifecycle within environment resources through explicit termination and final assay; every state transition and resource event enters the replayable record.

reaction–distillation cases with zero exact task–world identity overlap.

The coverage selection and pass rules are fixed before the reported qualification run and remain invariant across all outcomes. The reader-facing result is the complete frozen census defined by that design.

9.3 Qualification measurements

For each case, the report records normalized construction input, compile diagnostics, public contract, action sequence, schema and transaction status, constitution checks, events, resource preflight and outcome, public observation, termination, trajectory size and exact replay. Missing receipts, non-finite quantities, unexpected commits or roll-backs, leakage, denominator drift and replay mismatch fail the case.

Reference qualification covers 64 task–world units and 1,786 complete recipes. Generated qualification covers

52 compositions. Negative qualification covers 192 invalid probes. Module and interface qualification use 32 and seven units, respectively. Compile mutation uses seven invalid declarations. Counts are exact qualification denominators.

The 32 physical-module probes form an 8×4 design: each of reaction, thermal, phase, separation, crystallization, distillation, continuous flow and electrochemistry contributes one zero-input bounded-runtime probe, one legal low/high probe, one directionality probe and one runtime-constitution probe. Legal low/high checks use numerical reference fixtures for seven modules and a declared conceptual/synthetic fixture for crystallization; directionality checks are declared conceptual/synthetic oracles. Every numerical fixture records its own tolerance. Observation is qualified separately through instrument, sample-accounting, bounded-signal and public/private-boundary checks. The seven compile mutants comprise two missing-dependency cases and one case each for conflicting state

ownership, unit mismatch, invalid parameter domain, life-cycle hole and resource impossibility.

9.4 Process-time envelopes

Environment process time is derived by pattern rather than assigned as an arbitrary common cap. For each pattern,

$$t_{\max} = t_{\text{required stages}} + t_{\text{implicit reserve}} + t_{\text{allowed repeats}}.$$

The required term sums the upper bounds of necessary timed stages. The implicit reserve covers authored quench and transfer operations. Repeat allowance is tied to explicit per-operation repeat limits. The resulting limits are 0 s for phase observation; 3,600 s for reaction-thermal observation; 1,860 s for phase separation; 11,100 s for reaction crystallization; 10,440 s for reaction distillation; 7,200 s for continuous flow; 5,400 s for electrochemistry; and 7,500 s for reaction-phase separation. A proposed action is rejected before runtime if either cumulative process time or a repeat limit would be exceeded.

9.5 Deterministic use cases

The eight cases, seeds and action lists were protocol-frozen before execution. Their total expected census was 89 submitted actions, 88 commits, one rollback and eight final assays. The failure-recovery case specified the failing action and rollback class in advance. All submitted actions were inspected as one deterministic census, isolating instrument qualification from provider variance.

9.6 Controlled forks

Each fork declares a parent, a child, one protocol-frozen private intervention target, an invariant public contract and expected divergence channels. Parent and child execute the same typed action sequence. Gates require lineage validity, exactly one changed private target, invariant public contract, executable sequence on both variants, expected physical and observation divergence and exact replay under deterministic execution.

9.7 Complete-agent non-reference-world protocol

The formal unit is one complete lifecycle on the first protocol-frozen non-reference reaction-distillation composition. The protocol binds one uninterrupted agent session, one model/provider/scaffold configuration and at most 16 submitted actions through the public instrument interface. Termination and final assay must be agent-issued. The action count, tool calls and trajectory records must agree exactly; every action must commit; the lifecycle must contain one termination and exactly one final assay; resources, public boundary and exact replay must pass.

Provider accounting distinguishes one provider session, one logical agent turn, 15 action calls and two read-only calls. The model, provider, reasoning effort, scaffold constraints, public prompt template and execution date are bound before the run. Cumulative input is separated into cached and uncached input; cached context is reused input, not repeated output. The formal limits are 640,000 cumulative input tokens, 192,000 uncached input tokens and

64,000 output tokens. Per-action wait is capped at 600 s, finalization at 300 s and runner reserve at 600 s, giving a conservative method wall limit of 10,500 s. These method resources are separate from the simulated process-time ledger.

9.8 Process readouts

The process profile retains 19 dimensions in five groups: terminal commitment, evidence acquisition, evidence-conditioned action, resource deployment and outcome trajectory. Undefined conditional quantities remain null rather than being set to zero. The resulting coordinate vector preserves process structure for subsequent agent studies and supports dimension-level comparison without a composite score. The released process-coordinate contract supplies each dimension’s numerator, denominator, value range, null condition and boundary-case rule together with the computation binding used by the reports.

9.9 Public boundary and exact replay

The evaluator owns hidden simulator identity and private state. Agent-facing task cards, observations and histories are checked for hidden fields and absolute private paths. Replay reconstructs the bound world from the same normalized contract, runtime, mechanism, observation and scoring identities plus the recorded seeds and interventions. It then resubmits every committed typed action and compares rewards, every public observation, termination and truncation flags, operation types, transaction status, rollback reason, events, state-delta summaries, constitution checks and resource consequences. The reported qualification uses numerical tolerance zero and observed a maximum absolute error of zero. Release bindings define the replay boundary; cross-version or cross-platform reproduction uses the same recorded identities and model definitions. The public release contains the environment evidence needed for this reconstruction while keeping authentication data, private reasoning and unrestricted provider payloads outside the research artifact.

10. Data and code availability

Code, configuration, processed reports, figure source data and release tooling are available in the MIT-licensed ChemWorld repository at github.com/sunyrain/ChemWorld. The tracked materials regenerate the tables and figures and replay released simulator transitions and resource changes. The public submission tag `first-paper-arxiv-v1-2026-08-06` binds the manuscript source, dependency lock, processed evidence, figure data, coverage records and release metadata used for this version. Provider authentication, unrestricted response bodies, private reasoning and hidden evaluator identities are excluded.

The release supports three complementary reproducibility layers. Processed evidence regenerates the reported counts and figures. Released trajectories reconstruct environment transitions, public observations and ledgers. Provider provenance binds the model, configuration and

call accounting for the complete-agent unit. Exact replay refers to the executable world and its complete experimental record.

11. Conclusion

ChemWorld turns simulated chemistry from a fixed task collection into a programmable experimental medium. Reusable components compile into executable worlds; one public contract joins operations, instruments, observations, resources, failures, termination and evaluation; and every committed transition enters a version-bound replayable record. The 15 reference tasks anchor the public surface, while coverage-guided construction expands it to new world identities without changing the runtime contract.

The resulting advantage is controlled experimental freedom at software scale. Researchers can reset worlds exactly, repeat matched experiments without physical consumables or chemical hazard, inspect complete simulator-side consequences and change one private law under an invariant agent interface. Full-census qualification, process-complete use cases, controlled forks and a complete-agent lifecycle show that these capabilities operate together across the declared v1 domain. ChemWorld therefore provides an efficient and extensible substrate for controlled counterfactual agent studies, resource-aware experimentation and focused translation into physical laboratories.

Appendix A. Reader-facing capability map

A reference identity is a registered task identity paired with one of its declared public world seeds. The 15 task identities below expand to 64 task-world units. A generated row is counted as overlapping the reference set only when both its task identity and compiled world identity match a registered unit; the eight protocol-frozen non-reference rows have zero such overlap.

Appendix B. Coverage reconstruction and qualification census

The coverage design below fixes the pattern, seed, continuous domain, workflow count and generated denominator. Discrete factors include component-specific family or profile choices and instrument profiles; the released machine-readable coverage records map every discrete level, compatible pair, continuous stratum and ordered interaction to the rows that cover it. Covered and required counts are identical for all four target classes: 60/60 levels, 180/180 compatible pairs, 212/212 continuous strata and 84/84 ordered interactions.

Appendix C. Instrument-use case library

Appendix D. Process-coordinate dictionary

The 19 process dimensions remain separate and are grouped as follows.

Terminal commitment: closed-lifecycle fraction, assay fraction and discard fraction.

Evidence acquisition: measured-lifecycle fraction,

non-final instrument uses per closed lifecycle and normalized first-measurement position.

Evidence-conditioned action: continued-after-measurement fraction, post-measure process operations per closed lifecycle, threshold-eligible fraction and threshold-decision concordance.

Resource deployment: attempted operations per closed lifecycle, committed operations per closed lifecycle, cost per closed lifecycle and risk per closed lifecycle.

Outcome trajectory: normalized best-discovery position, online incumbent-retention rate, maximum absolute incumbent drawdown, loss-episode recovery rate and terminal-to-best ratio. Mean and best endpoint scores are reported beside this profile and never enter a composite score.

Appendix E. Component model cards and extension points

Each module declares its virtual-instrument domain explicitly. The table gives the runtime formulation, authored domain, qualification oracle and the interface location intended for future alternative or empirically calibrated implementations.

Table 2. Complete reference-task registry. Seed counts define the public world identities exercised for each task.

Reference task identity	Component topology	World seeds	Public operation / instrument summary
electrochemical-conversion	reaction + electrochemistry + observation	5	6 operations; pH, UV-visible, final assay
equilibrium-characterization	reaction + thermal + observation	5	9 operations; pH, UV-visible, final assay
flow-reaction-optimization	reaction + continuous flow + observation	5	7 operations; HPLC, GC, UV-visible, final assay
low-budget-characterization	reaction + thermal + observation	3	9 operations; HPLC, GC, UV-visible, final assay
partition-discovery	reaction + phase + separation + observation	5	9 operations; HPLC, GC, UV-visible, final assay
public-private-generalization	reaction + thermal + observation	5	9 operations; HPLC, GC, UV-visible, final assay
purity-yield-tradeoff	reaction + thermal + phase + separation + observation	5	18 operations; HPLC, GC, UV-visible, final assay
reaction-mechanism-explanation	reaction + thermal + observation	3	9 operations; HPLC, GC, UV-visible, final assay
reaction-optimization-standard	reaction + thermal + observation	5	9 operations; HPLC, GC, UV-visible, final assay
reaction-safety-constrained	reaction + thermal + observation	5	9 operations; HPLC, GC, UV-visible, final assay
reaction-to-assay	reaction + thermal + observation	1	9 operations; HPLC, GC, UV-visible, final assay
reaction-to-crystallization	reaction + thermal + crystallization + observation	5	12 operations; HPLC, final assay
reaction-to-distillation	reaction + thermal + distillation + observation	5	12 operations; HPLC, GC, UV-visible, final assay
reaction-to-purification	reaction + thermal + phase + separation + observation	5	18 operations; HPLC, GC, UV-visible, final assay
tool-agent-planning	reaction + thermal + phase + separation + observation	2	18 operations; HPLC, GC, UV-visible, final assay

Component pattern	Principal state or process	Representative public operations	Representative instruments
Phase + observation	bounded phase/equilibrium state	add solvent, add reagent, measure, terminate	pH, UV-visible, final assay
Reaction + thermal + observation	batch reaction and temperature history	add, heat, quench, sample	HPLC, GC, final assay
Phase + separation + observation	phase formation and transfer	mix, settle, separate, wash, transfer	HPLC, final assay
Reaction + thermal + crystallization + observation	reaction followed by solid formation	heat, seed, cool, filter	HPLC, particle sizing, final assay
Reaction + thermal + distillation + observation	reaction, evaporation and fractionation	heat, quench, evaporate, distil, collect	HPLC, GC, final assay
Reaction + thermal + continuous flow + observation	flow, residence time and conversion	set flow, set temperature, run, sample	HPLC, GC, final assay
Reaction + electrochemistry + observation	potential/current-driven conversion	set potential/current, electrolyse, sample	voltammetry, HPLC, final assay
Reaction + thermal + phase + separation + observation	multistage reaction and purification	react, quench, separate, wash, concentrate, transfer	HPLC, GC, final assay

Table 3. Topology and identity decomposition of the generated block. Topology novelty compares exact component sets with the 15-task reference registry; task-world novelty uses exact registered identities.

Generated group	Cases	Topology relation	Exact task-world relation
phase-observation	6	absent from reference registry	zero overlap implied by component-set difference
phase-separation-observation	6	absent from reference registry	zero overlap implied by component-set difference
reaction-thermal-continuous-flow-observation	6	absent from reference registry	zero overlap implied by component-set difference
reaction-thermal-distillation-observation	8	registered topology	zero exact overlap in the protocol-frozen non-reference block
four remaining generated patterns	26	registered topologies	used for reference-topology coverage; no separate non-reference identity claim

Table 4. Protocol-frozen coverage design. Bounds are inclusive authored domains; “none” denotes a purely discrete design.

Pattern	Seed	Continuous axes and bounds	Workflows	Cases
phase-observation	101	none	1	6
reaction-thermal-observation	102	heat 350–390 K; duration 600–1,800 s	2	6
phase-separation-observation	103	phase 0.010–0.020 L; extractant 0.010–0.025 L; mix 60–300 s; settle 120–600 s	2	6
reaction-thermal-crystallization-observation	104	reaction 350–390 K, 600–1,800 s; seed 0.002–0.010 g; cooling 275–305 K, 900–3,600 s	2	6
reaction-thermal-distillation-observation	105	reaction 350–390 K, 600–1,800 s; evaporation 325–345 K, 300–900 s; distillation 350–390 K, 900–2,400 s; reflux 1.0–3.0; transfer 0.65–0.95	2	8
reaction-thermal-continuous-flow-observation	106	flow 0.5–5.0 mL min ⁻¹ ; residence 60–600 s; temperature 330–390 K	2	6
reaction-electrochemistry-observation	107	potential 0.5–1.8 V; current 25–150 mA; electrolysis 300–1,800 s	2	7
reaction-thermal-phase-separation-observation	108	reaction 350–390 K, 600–1,800 s; phase/extractant 0.010–0.020/0.010–0.025 L; mix 60–300 s; settle 120–600 s; wash 0.003–0.010 L; concentrate 300–900 s; transfer 0.65–0.95	2	7

Qualification unit	Passed	Denominator	Failure classes
Reference task-world units	64	64	0
Complete reference recipes	1,786	1,786	0
Coverage-generated compositions	52	52	0
Protocol-frozen non-reference reaction-distillation compositions	8	8	0
Invalid action probes	192	192	0 unexpected outcomes
Module probes	32	32	0
Cross-module interface paths	7	7	0
Invalid compile mutants	7	7	0 unexpected constructions
Deterministic use cases	8	8	0
Deterministic submitted actions	89	89	0 missing receipts
Controlled fork pairs	6	6	0
Controlled fork traces	24	24	0

Scientific use	Components	What the record demonstrates
Reaction to crystallization	reaction, thermal, crystallization, observation	propagation from reaction through seeding, cooling, filtration and final assay
Resource-limited characterization	phase, observation	measurement choice, sample consumption and explicit stopping under a small budget
Failure and recovery	reaction, thermal, phase, separation, observation	atomic rollback, attempt consequences and continuation from committed state
Controlled private-law fork	one registered component changed privately	invariant public contract with protocol-defined state/observation divergence
Generated reaction to distillation	reaction, thermal, distillation, observation	construction and replay outside the reference task identities; one complete-agent lifecycle closed under the same public contract
Reference library	flow, electrochemistry, distillation, partition, crystallization	breadth of reusable task recipes without a cross-task performance score

Component	Runtime formulation	Principal v1 domain	Qualification oracle	Intended use and extension path
Reaction	stoichiometric mass-action network with Arrhenius temperature dependence	authored reaction families and bounded batch temperature/time	exact amount fixtures, monotonic response, material closure and runtime constitution	reaction-law extension point for future named-reaction calibration
Thermal	dynamic batch heat-release and jacket-energy balance	bounded temperature, duration, vessel pressure and volume	temperature/energy finiteness, bounds, event propagation and ledger reconciliation	thermal-contract extension point for future equipment-specific coefficients
Phase	stability-gated, activity-corrected liquid-liquid equilibrium with TPD-style diagnostics	declared phase identities, volumes and composition ranges	phase/material balance, directional partition response and state identity	activity-model extension point for future compound-specific thermodynamics
Separation	settling, entrainment, wash and transfer coupled to the phase model	bounded mix/settle time, extractant/wash volume and transfer fraction	amount/unit conservation, transfer identity and expected directional response	separation-interface extension point for future hardware-scale transport
Crystallization	van't Hoff solubility with seed, nucleation/growth cohorts, impurity occlusion and CSD summaries	bounded seed mass, cooling temperature and cooling time	material closure, solubility-direction checks, CSD and runtime-constitution receipts	law extension point for future calibrated nucleation and growth models
Distillation	bubble-gated, duty-limited VLE/Fenske fractionation with material and energy ledgers	bounded temperature/time, reflux ratio, fraction count and collected fraction	mass/energy closure, fraction identity, recovery/purity directions and equipment limits	operation-contract extension point for future compound and column models
Continuous flow	geometry-resolved plug-flow reactor with residence time, distributed thermal boundary and pressure drop	bounded flow, residence time and temperature	conversion direction, mass closure, pressure/geometry and solver diagnostics	runtime extension point for future reactor-specific transport or control models
Electrochemistry	Nernst potential, Butler-Volmer kinetics, limiting current, Randles transient and Faraday accounting	bounded potential, current and electrolysis time	charge/material closure, signed work, selectivity and limiting-current checks	electrochemical-law extension point for future material- and cell-specific parameters
Observation	state-coupled synthetic pH, UV-visible, HPLC, GC and final-assay contracts	task-declared instruments, sample and use budgets	instrument availability, sample consumption, finite/bounded signal and non-omniscience checks	instrument-schema extension point for future empirical response models

References

- [1] Daniil A. Boiko, Robert MacKnight, Ben Kline, and Gabe Gomes. Autonomous chemical research with large language models. *Nature*, 624:570–578, 2023. doi: 10.1038/s41586-023-06792-0.
- [2] Andres M. Bran, Sam Cox, Oliver Schilter, Carlo Baldassari, Andrew D. White, and Philippe Schwaller. Augmenting large language models with chemistry tools. *Nature Machine Intelligence*, 6:525–535, 2024. doi: 10.1038/s42256-024-00832-8.
- [3] Nathan J. Szymanski et al. An autonomous laboratory for the accelerated synthesis of inorganic materials. *Nature*, 624:86–91, 2023. doi: 10.1038/s41586-023-06734-w.
- [4] Tianyi Dai, Sriram Vijayakrishnan, Filip T. Szczypiński, et al. Autonomous mobile robots for exploratory synthetic chemistry. *Nature*, 635:890–897, 2024. doi: 10.1038/s41586-024-08173-7.
- [5] Kouros Darvish, Marta Skreta, Yuchi Zhao, et al. OR-GANA: A robotic assistant for automated chemistry experimentation and characterization. *Matter*, 8(2), 2025. doi: 10.1016/j.matt.2024.10.015.
- [6] Tao Song, Man Luo, Xiaolong Zhang, et al. A multiagent-driven robotic AI chemist enabling autonomous chemical research on demand. *Journal of the American Chemical Society*, 147(15):12534–12545, 2025. doi: 10.1021/jacs.4c17738.
- [7] Gihan Panapitiya, Emily Saldanha, Heather Job, et al. AutoLabs: cognitive multi-agent systems with self-correction for autonomous chemical experimentation. *Scientific Reports*, 16:19554, 2026. doi: 10.1038/s41598-026-45593-z.
- [8] Simone Pilon, Elia Savino, Oliver M. Bayley, et al. A flexible and affordable self-driving laboratory for automated reaction optimization. *Nature Synthesis*, 2026. doi: 10.1038/s44160-026-01053-0.
- [9] Aikaterini Vriza, Michael H. Prince, Tao Zhou, Henry Chan, and Mathew J. Cherukara. Operating advanced scientific instruments with AI agents that learn on the job. *npj Computational Materials*, 12:160, 2026. doi: 10.1038/s41524-026-02005-0.
- [10] Kobi C. Felton, Jan G. Rittig, and Alexei A. Lapkin. Summit: Benchmarking machine learning methods for reaction optimisation. *Chemistry–Methods*, 1:116–122, 2021. doi: 10.1002/cmtd.202000051.
- [11] Florian Häse, Matteo Aldeghi, Riley J. Hickman, Loïc M. Roch, Melodie Christensen, Elena Liles, Jason E. Hein, and Alán Aspuru-Guzik. Olympus: a benchmarking framework for noisy optimization and experiment planning. *Machine Learning: Science and Technology*, 2:035021, 2021. doi: 10.1088/2632-2153/abedc8.
- [12] Maximilian Bloor, José Torracca, Ilya Orson Sandoval, et al. PC-Gym: Benchmark environments for process control problems, 2024. Preprint, arXiv:2410.22093.
- [13] Chris Beeler, Sriram Ganapathi Subramanian, Kyle Sprague, et al. ChemGymRL: A customizable interactive framework for reinforcement learning for digital chemistry. *Digital Discovery*, 3:742–758, 2024. doi: 10.1039/D3DD00183K.
- [14] Shreshth A. Malik, Tiarnan Doherty, Panagiotis Tigas, Muhammed Razzak, Stephen J. Roberts, Aron Walsh, and Yarin Gal. MADE: Benchmark environments for closed-loop materials discovery. In *International Conference on Machine Learning*, 2026.
- [15] Nikhil Abhyankar, Sanchit Kabra, Saaketh Desai, and Chandan K. Reddy. LLEMA: Evolutionary search with LLMs for multi-objective materials discovery. In *International Conference on Learning Representations*, 2026. URL <https://openreview.net/forum?id=TIqzhBvCNB>.
- [16] Peter Jansen, Marc-Alexandre Côté, Tushar Khot, Erin Bransom, Bhavana Dalvi Mishra, Bodhisattwa Prasad Majumder, Oyvind Tafjord, and Peter Clark. DiscoveryWorld: A virtual environment for developing and evaluating automated scientific discovery agents. In *Advances in Neural Information Processing Systems*, volume 37, 2024. doi: 10.52202/079017-0324.
- [17] Kanishk Gandhi, Michael Y. Li, Lyle Goodyear, Agam Bhatia, Louise Li, Aditi Bhaskar, Mohammed Zaman, and Noah D. Goodman. BoxingGym: Benchmarking progress in automated experimental design and model discovery, 2025. Preprint, arXiv:2501.01540.
- [18] Haonan Duan, Stephen Zhewen Lu, Caitlin Fiona Harri-gan, Nishkrit Desai, Jiarui Lu, Michał Koziarski, Leonardo Cotta, and Chris J. Maddison. Measuring scientific capabilities of language models with a systems biology dry lab, 2025. Preprint, arXiv:2507.02083.
- [19] Maximilian Nägele and Florian Marquardt. Agentic exploration of physics models. *Physical Review X*, 16:031002, 2026. doi: 10.1103/xnqc-q6nt.
- [20] Tianshi Zheng, Kelvin Kiu-Wai Tam, Newt Hue-Nam K. Nguyen, et al. NewtonBench: Benchmarking generalizable scientific law discovery in llm agents. In *International Conference on Learning Representations*, 2026.
- [21] Rui Li, Zixuan Hu, Wenxi Qu, et al. LabUtopia: High-fidelity simulation and hierarchical benchmark for scientific embodied agents. In *Advances in Neural Information Processing Systems, Datasets and Benchmarks Track*, 2025.
- [22] Yuhan Wu, Zhao Jin, Tao Li, et al. Labimus: A simulation and benchmark for humanoid dexterous manipulation in chemical laboratory, 2026. Preprint.
- [23] David M. Cohen, Siddhartha R. Dalal, Michael L. Fredman, and Gardner C. Patton. The AETG system: An approach to testing based on combinatorial design. *IEEE Transactions on Software Engineering*, 23(7):437–444, 1997. doi: 10.1109/32.605761.
- [24] Luc Moreau and Paolo Missier. PROV-DM: The PROV data model. W3C Recommendation, April 2013. URL <https://www.w3.org/TR/prov-dm/>.