

Algebra

A course for Ph. D. students

Algebra

A course for Ph. D. students

Abhay Soman

University of Hyderabad

September 3, 2025

©2025–2026

This work is licensed under the Creative Commons Attribution-ShareAlike 4.0 International License. To view a copy of this license, visit [CreativeCom-mons.org](https://creativecommons.org)

Contents

1	Associative Algebras	1
1.1	Algebra over a commutative ring	1
1.2	Some examples of algebras	2
1.3	Group algebra	2

Back Matter

Chapter 1

Associative Algebras

We follow “Associative algebra” by R. S. Pierce. We do not claim any originality for the content.

1.1 Algebra over a commutative ring

We begin with a definition of *algebra* over a commutative ring.

Definition 1.1.1 Associative algebra. Let R be a commutative ring with unity 1_R . An R -**algebra** or algebra over R is a unital right R -module A together with R -bilinear map $A \times A \rightarrow A$ denoted by $(x, y) \mapsto xy$ that is associative, i.e., $(xy)z = x(yz)$ for all $x, y, z \in A$, and there is a unity $1_A \in A$ such that $1_R x = x 1_R = x$ for all $x \in A$. \diamond

Observation 1.1.2 The bilinearity of the map $A \times A \rightarrow A$ in [Definition 1.1.1](#) implies the following relation for any $x, y \in A$ and any $r \in R$.

$$(xr)y = x(yr) = (xy)r \quad (1.1.1)$$

We have the following equivalent definition of R -algebra.

Proposition 1.1.3 *Following statements are equivalent.*

1. A is an R -algebra
2. A is a ring and a right R -module such that for any $x, y \in A$ and $r \in R$, we have $(xr)y = x(yr) = (xy)r$.

Proof. Assume that A is an R -algebra. Then A is a ring with the multiplication given by the R -bilinear map $A \times A \rightarrow A$. Indeed, the associativity of the multiplication follows from the definition. The distributivity of the multiplication over addition follows from the bilinearity of the map. The existence of unity is also given in the definition. The relation $(xr)y = x(ry) = (xy)r$ follows from [Observation 1.1.2](#).

Conversely, assume that A is a ring and a right R -module such that for any $x, y \in A$ and $r \in R$, we have $(xr)y = x(yr) = (xy)r$. We consider a map $m: A \times A \rightarrow A$ as the multiplication in ring A . Therefore, $m(x + y, z) = m(x, z) + m(y, z)$ (resp., $m(x, y + z) = m(x, y) + m(x, z)$) for any $x, y, z \in A$ and also there is a unity in ring A . The R -linearity of the map m follows from the relation $(xr)y = x(yr) = (xy)r$, i.e., $m(xr, y) = m(x, yr) = (x, y)r$. \blacksquare

Lemma 1.1.4

1. If A is an R algebra then, there is a ring homomorphism $f: R \rightarrow A$ such

that $f(R)$ is contained in the center of A .

2. Suppose that A is a ring and there is a ring homomorphism $f: R \rightarrow A$ such that $f(R)$ is contained in the center of A . Then, A is an R -algebra with the right R -module structure given by $xr = xf(r)$ for any $x \in A$ and $r \in R$.

Proof. If A is an R -algebra then, we define a map $f: R \rightarrow A$ by $f(r) = 1_A r$ for any $r \in R$. It is easy to see that f is a ring homomorphism and $f(R)$ is contained in the center of A . Conversely, suppose that A is a ring and there is a ring homomorphism $f: R \rightarrow A$ such that $f(R)$ is contained in the center of A . We consider the right R -module structure on A given by $xr = xf(r)$ for any $x \in A$ and $r \in R$. Then, it is easy to see that the relation in Proposition 1.1.3(2) holds. Therefore, by Proposition 1.1.3, we conclude that A is an R -algebra. ■

1.2 Some examples of algebras

Example 1.2.1 Associative ring as \mathbb{Z} -algebra. Every associative ring A can be considered as a \mathbb{Z} -algebra. □

Example 1.2.2 Ring as an algebra over itself. Let R be a commutative ring with unity. Then R can be considered a right R -module using the multiplication in ring R . With this considerations, R is an R -algebra. □

Example 1.2.3 Ring over its center. Suppose that A is a ring (not necessarily commutative), and let $Z(A)$ be the center of A . Then A is a $Z(A)$ -algebra. □

Example 1.2.4 Matrix algebra. Suppose that R is a commutative ring and n is a positive integer. Suppose that A is an R -algebra. Then the ring of $n \times n$ matrices over A , denoted by $M_n(A)$, is an R -algebra. □

Example 1.2.5 Endomorphism algebra. Let R be a commutative ring with unity. Suppose that M is a right R -module. The endomorphism ring $\text{End}_R(M)$ is an R -algebra. □

Example 1.2.6 Polynomial ring as algebra. Let R be a commutative ring with unity. The polynomial ring $R[X]$ is an R -algebra. □

We discuss the example of ‘Group algebra’ in the next section.

1.3 Group algebra

Definition 1.3.1 Group Ring. Let R be a commutative ring with unity and G a group. Consider the following set.

$$RG = \{\phi \in \text{Hom}_{\text{Sets}}(G, R) : \phi(g) = 0 \text{ for almost all } g \in G\}$$

Define ‘addition’ and ‘scalar multiplication’ in RG as follows.

$$(\phi r + \psi s)(g) = \phi(g)r + \psi(g)s$$

The ‘multiplication’ in RG is defined as convolution product.

$$(\phi\psi)(g) = \sum \phi(h)\psi(h^{-1}g) \text{ such that } h \in G, \text{ and } \phi(h)\psi(h^{-1}g) \neq 0$$

◇

Checkpoint 1.3.2 Show that RG is closed under addition, scalar multiplication, and it is an R -module. Furthermore, show that multiplication is bilinear.

We now proceed to check the associativity of multiplication and the existence of unity in RG . We first observe that RG is a free R -module.

Proposition 1.3.3 *Keep notations of Definition 1.3.1 above. Let for $g \in G$, χ_g be the characteristic function of g , i.e., $\chi_g(h) = 0$ if $h \neq g$ and $\chi_g(g) = 1$. Then, set $\{\chi_g : g \in G\}$ is an R -basis of RG . In particular, RG is a free R -module.*

Furthermore, $G \rightarrow RG$ given by $g \mapsto \chi_g$ is a monoid homomorphism. In particular, $\chi_g \chi_h = \chi_{gh}$ and if $e \in G$ is the identity then, χ_e is the unity in RG .

Proof. Suppose that $\sum \chi_{g_i} r_i = 0$ for some $g_i \in G$ and $r_i \in R$. Evaluating at g_j , we get $r_j = 0$. Thus, the set is linearly independent. Now, let $\phi \in RG$. Then, we can write $\phi = \sum \chi_g \phi(g)$ where the sum is over all $g \in G$ such that $\phi(g) \neq 0$. This shows that the set spans RG . Thus, it is an R -basis of RG .

Direct calculation shows that $\chi_g \chi_h = \chi_{gh}$ for all $g, h \in G$.

We show the associativity of multiplication. Let $\phi, \psi, \zeta \in RG$. Then, there exists $r_i, s_j, t_k \in R$ such that $\phi = \sum \chi_{g_i} r_i$, $\psi = \sum \chi_{g_j} s_j$, $\zeta = \sum \chi_{g_k} t_k$. Therefore, we obtain the following.

$$\begin{aligned} (\phi\psi)\zeta &= \left(\left(\sum \chi_{g_i} r_i \right) \left(\sum \chi_{g_j} s_j \right) \right) \sum \chi_{g_k} t_k \\ &= \sum ((\chi_{g_i} \chi_{g_j}) \chi_{g_k}) (r_i s_j t_k) \\ &= \sum \chi_{g_i} (\chi_{g_j} \chi_{g_k}) (r_i s_j t_k) \\ &= \phi(\psi\zeta) \end{aligned}$$

This shows the associativity of multiplication.

For any $\chi_{g_i} a_i \in RG$ we get the following.

$$((\chi_{g_i} a_i) \chi_e)(h) = \sum \chi_{g_i} a_i(x) \chi_e(x^{-1}h)$$

The above equation is nonzero only when $x = g_i$ and $x^{-1}h = e$. This implies that $h = g_i$. Thus, we get $((\chi_{g_i} a_i) \chi_e)(h) = \chi_{g_i} a_i(h)$. Similarly, we can show that $(\chi_e(\chi_{g_i} a_i))(h) = \chi_{g_i} a_i(h)$. This shows that χ_e is the unity in RG . ■

Colophon

This book was authored in PreTeXt.