

Genomic Insights into Extreme Salinity Tolerance in *Salicornia*

Yunchuan Wang¹, Laxman Adhikari¹, Mark A. Tester¹ and Vanessa Melino^{2*}

¹Plant Science Program, Division of Biological and Environmental Sciences and Engineering, King Abdullah University of Science and Technology (KAUST), Thuwal, Saudi Arabia

²Centre for Plant Science, School of Science, University of Newcastle, Callaghan, NSW, Australia.

*Corresponding author: vanessa.melino@newcastle.edu.au



Summary

- Chromosome-scale, annotated genome assemblies were generated for six *Salicornia* species.
- Four distinct subgenomes were identified within the *Salicornia* genus.
- Comparative genomics revealed lineage-specific expansions of the Cupin_1 and CHS gene families
- Global resequencing panel established for both genetic research and breeding applications.
- A database of functionally validated salt stress genes was curated via large-scale literature review using LLM agents.

Introduction





Figure 1. Plants were grown for 6 weeks under NaCl concentrations of 0, 50, 200, 600 mM.

a: Plants 6 weeks after treatment. b: Length of individual plants 6 weeks after treatment. (From Salazar et al 2024)



From Jen Christiansen and Tim Flowers

Saltwater-based agriculture is a sustainable solution to address escalating water scarcity and the growing demand for irrigation water for cropping. Halophytes such as *Salicornia*, a genus in the Amaranthaceae family, is valued for its edible succulent stems and oil-rich seeds, yet its genetic improvement has been constrained by limited genomic resources.

Salicornia plants grow optimally at around 200 mM NaCl, and still grow and complete their life cycle at 600 mM NaCl(Fig 1); these conditions are lethal to most plants. Remarkably, salt promotes the growth of *Salicornia* spp., and a mechanism of tolerance is to sequester the sodium into the vacuole of shoots - referred to as tissue ion tolerance.

Material - Seeds Collection




Figure 2. Geographic distribution of Salicornioideae and morphological characteristics within the *Salicornia* genus
Geographic distribution of *S. bigelovii*, *S. europaea* (France), *S. europaea* (Israel), *S. persica* ssp. *iranica*, *S. sinus-persica*, *S. brachiata* and their diverse morphological characteristics. Distribution data (light blue dots) from GBIF.org

Results




Figure 3. Intergenomic synteny among *Salicornia* lineages.

a: Synteny blocks are connected in gray. The large reciprocal translocations in chromosome three with chromosome eight in *S. persica* ssp. *iranica* are highlighted in light green. Other Large reciprocal translocations are highlighted in pink including. b: Reciprocal translocations revealed by oligo-FISH (*S. europaea* (Israel)) on mitotic metaphase chromosomes of *S. persica* ssp. *iranica* and c: *S. bigelovii*.




Figure 4. Orthogroup-based phylogeny, copy number variation, and gene family expansion in the Amaranthaceae family.

A time-calibrated phylogenetic tree built from single-copy ortholog protein sequences using maximum likelihood. The top axis shows divergence time (MYA). Blue numbers at internal nodes denote inferred gene-family expansions and contractions on the branch leading to each node ($p < 0.05$; CAFÉ). Right columns show per-genome gene counts for the Cupin_1 and chalcone synthase (CHS) families, with bar lengths proportional to copy number. Light blue: species with Cupin_1 family expansions; dark blue: species with CHS family expansions; grey: species without expansions.

Reference

Wang, Y., Adhikari, L., Leal, L. C., Kovalchuk, N., Čížková J, Beránková D, ... Tester, T. & Melino, V. (2025). From Wild Halophyte to Future Crop: Genomic Insights into *Salicornia*. (preprint, DOI: 10.21203/rs.3.rs-7061928/v1)
Salazar, O. R., Chen, K., Melino, V. J., Reddy, M. P., Hřibová, E., Čížková, J., ... & Schmöckel, S. M. (2024). SOS1 tonoplast neo-localization and the RGG protein SALT are important in the extreme salinity tolerance of *Salicornia bigelovii*. *Nature Communications*, 15(1), 4279.

A Literature-Curated Salt Stress Gene Resource Built by Multi-LLM Consensus

PubMed corpus acquisition



Salt-stress + plant query

122,617 abstracts

Topic filtering

KeyBERT + UMAP + HDBSCAN

Haiku 4.5

Noise: 55,644 abstracts

Retained for LLM extraction:

n = 45,639 abstracts

Dual-LLM gene extraction

Sonnet 4.6 → 45,107 calls

Opus 4.7 → 44,800 calls

Agreement: 6,112 abstracts

Disagreement: 1,403 abstracts

Opus adjudication: keep 1,748 | drop 669

Entity normalization

NHX7=SOS1

Alias clusters(gene-level): 7,938

Unique species(normalized): 808

Unique PMIDs with gene: 6,968

Full-text extension

Full text Download:

PMC OA 3,228 + KAUST 2,262

5,490/6,902 papers(79.5%)

PubMed Central (PMC)

Open Access Bulk Mirror

Sonnet 4.6 → 5,486/5,490

Opus 4.7 → 5,444/5,490

Noise rescue


Keyword-density filter on noise

55,644 → 52 candidates → sonnet

+ Dual-LLM gene extraction

+21 rows genes

Database construction



GitHub-based Website

Salt-stress gene table

9,849 rows

7,938 unique gene

808 species

6,968 supporting PMIDs

Acknowledgements

KAUST

Professor Jesse Poland

Laxman Adhikari

Lina Maria Caceres Leal

Nataliya Kovalchuk

Noha Saber

Gabriele Fiene

Yveline Pailles

Institute of Experimental Botany of the Czech Academy of Sciences

Professor Eva Hřibová

Jana Čížková

Denisa Beránková

QR Code

Poster



Scan to Read

LinkedIn



Scan to connect