

# Solutions to "Algebraic Geometry and Arithmetic Curves"

Quentin Asparria

July 21, 2026

## 1 Some topics in commutative algebra

### 1.1 Tensor products

**Exercise 1.1.1.** Let  $\{M_i\}_{i \in I}, \{N_j\}_{j \in J}$  be two families of modules over a ring  $A$ . Then

$$(\oplus_{i \in I} M_i) \otimes_A (\oplus_{j \in J} N_j) \cong \oplus_{(i,j) \in I \times J} (M_i \otimes_A N_j).$$

*Proof.* There is a chain of natural isomorphisms:

$$\begin{aligned} & (\oplus_{i \in I} M_i) \otimes_A (\oplus_{j \in J} N_j) \\ & \cong \oplus_{i \in I} (M_i \otimes_A (\oplus_{j \in J} N_j)) \\ & \cong \oplus_{j \in J} \oplus_{i \in I} (M_i \otimes_A N_j) \\ & \cong \oplus_{(i,j) \in I \times J} (M_i \otimes_A N_j). \end{aligned}$$

□

**Exercise 1.1.2.** There is a unique  $A$ -linear map

$$f : \text{Hom}_A(M, M') \otimes_A \text{Hom}_A(N, N') \rightarrow \text{Hom}_A(M \otimes_A N, M' \otimes_A N')$$

such that  $f(u \otimes v) = u \otimes v$ .

*Proof.* Consider the map  $g : \text{Hom}_A(M, M') \times \text{Hom}_A(N, N') \rightarrow \text{Hom}_A(M \otimes_A N, M' \otimes_A N')$  sending  $(u, v)$  to  $u \otimes v$ . It is clearly bilinear, so factors uniquely through  $\text{Hom}_A(M, M') \otimes_A \text{Hom}_A(N, N')$ . □

**Exercise 1.1.3.** Let  $M, N$  be  $A$ -modules and let  $i : M' \rightarrow M, j : N' \rightarrow N$  be submodules of  $M$  and  $N$ , respectively. Then there is a canonical isomorphism

$$(M/M') \otimes_A (N/N') \cong (M \otimes_A N) / (\text{Im } i_N + \text{Im } j_N).$$

In particular,  $(\mathbf{Z}/n\mathbf{Z}) \otimes_{\mathbf{Z}} (\mathbf{Z}/m\mathbf{Z}) = \mathbf{Z}/l\mathbf{Z}$ , where  $l = \gcd(m, n)$ .

*Proof.* Observe that

$$(M \otimes_A N)/(\text{Im } i_N + \text{Im } j_N) \cong ((M \otimes_A N)/(\text{Im } i_N))/(\text{Im } j_{M/M'}),$$

so Corollary 1.13 gives us a chain of isomorphisms

$$\begin{aligned} & ((M \otimes_A N)/(\text{Im } i_N))/(\text{Im } j_{M/M'}) \\ & \cong ((M/M') \otimes_A N)/(\text{Im } j_{M/M'}) \\ & \cong (M/M') \otimes_A (N/N'). \end{aligned}$$

For the last statement, just recall that  $(m) + (n) = (l)$ . □

**Exercise 1.1.4.** Let  $M, N$  be  $A$ -modules and let  $B, C$  be  $A$ -algebras.

- (a) If  $M$  and  $N$  are finitely generated over  $A$ , then so is  $M \otimes_A N$ .
- (b) If  $B$  and  $C$  are finitely generated over  $A$ , then so is  $B \otimes_A C$ .
- (c) The converses of (a) and (b) are in general false.

*Proof.* Parts (a) and (b) are obvious. For part (c), just notice that  $\mathbf{Q} \otimes_{\mathbf{Z}} \mathbf{Z}/2\mathbf{Z}$  is the zero ring, but  $\mathbf{Q}$  is not finitely generated over  $\mathbf{Z}$ . □

**Exercise 1.1.5.** Let  $\rho : A \rightarrow B$  be a ring homomorphism,  $M$  an  $A$ -module, and  $N$  a  $B$ -module. There is a canonical isomorphism of  $A$ -modules

$$\text{Hom}_A(M, \rho_* N) \cong \text{Hom}_B(\rho^* M, N).$$

*Proof.* Let  $f : \text{Hom}_A(M, \rho_* N) \rightarrow \text{Hom}_B(\rho^* M, N)$  be the map sending  $u$  to the map  $f(u)$  s.t.  $f(u)(m \otimes b) = bu(m)$ . This is well-defined because if  $n \otimes c = m \otimes b$ , we may assume that there is some  $a \in A$  s.t.  $an = m$  and  $\rho(a)b = c$  and so  $f(u)(n \otimes c) = cu(n) = \rho(a)bu(n) = bu(m)$ .

In addition, define  $g : \text{Hom}_B(\rho^* M, N) \rightarrow \text{Hom}_A(M, \rho_* N)$  to be the map s.t.  $g(u)(m) = u(m \otimes 1)$ . This is immediately seen to be well-defined, and we contest that this is an inverse for  $f$ . Indeed,  $f(g(u))(m \otimes b) = bg(u)(m) = bu(m \otimes 1) = u(m \otimes b)$  and  $g(f(u))(m) = f(u)(m \otimes 1) = u(m)$ . □

**Exercise 1.1.6.** Let  $(N_i)_{i \in I}$  be a direct system of modules. Then for any  $A$ -module  $M$ , there exists a canonical isomorphism  $\varinjlim (N_i \otimes_A M) \cong (\varinjlim N_i) \otimes_A M$ .

*Proof.* Tensor-hom.

Just kidding, it follows from the fact that  $\varinjlim (N_i \times M)$  is canonically isomorphic to  $(\varinjlim N_i) \times M$ , and that a bilinear map over the latter must then induce a cocone of bilinear maps. □

**Exercise 1.1.7.** Let  $B$  be an  $A$ -algebra, and let  $M, N$  be  $B$ -modules. Then there exists a canonical surjection  $M \otimes_A N \rightarrow M \otimes_B N$ .

*Proof.* Do I really need to spell it out? □

## 1.2 Flatness

## 1.3 Formal completion

# 2 General properties of schemes

## 2.1 Spectrum of a ring

**Exercise 2.1.1.** For  $k$  a field,  $\text{Spec}(k[[T]])$  is the space  $\{\xi, 0\}$  with generic point  $\xi$  and closed point  $0$ .

*Proof.* It is easy to see that any power series with a non-zero constant term is a unit. Furthermore, if we have  $f = \sum a_n T^n$  with  $a_N = 1$  for some  $N$  and  $a_n = 0$  whenever  $n \leq N$ , then we can pull out a common factor of  $T^N$ . We see that  $f$  is then a unit multiple of  $T^N$  and  $(f) = (T^N)$ . Clearly, this is only prime when  $N = 1$ . It is easy to see that  $k[[T]]$  is an entire ring, so the proposition is immediate.  $\square$

**Exercise 2.1.2.** Let  $\varphi : A \rightarrow B$  be a homomorphism of finitely generated algebras over a field  $k$ . The image of a closed point under  $\text{Spec } \varphi$  is a closed point.

*Proof.* Observe that  $\varphi$  factors as

$$A \rightarrow A/\text{Ker } \varphi \rightarrow \text{Im } \varphi \rightarrow B.$$

Obviously, the first arrow satisfies the conditions, so we may assume  $\varphi$  injective. If  $\text{Spec } \varphi$  takes the maximal ideal  $\mathfrak{m}$  into the ideal  $\mathfrak{p}$ , we get a diagram

$$k \rightarrow A/\mathfrak{p} \xrightarrow{\tilde{\varphi}} B/\mathfrak{m}.$$

By Corollary 1.12,  $B/\mathfrak{m}$  is a finite algebraic extension of  $k$ , so  $B/\mathfrak{m}$  (resp.  $A/\mathfrak{p}$ ) is finite over  $A/\mathfrak{p}$  (resp.  $k$ ). It then suffices to prove that  $A/\mathfrak{p}$  is a field. Any  $a \in A$  has an associated irreducible polynomial  $P = \sum a_i X^i \in k[x]$  such that  $P(a) = 0$ . This polynomial must have a nonzero constant term, so we have

$$-\frac{1}{a_0} \sum_{i>0} a_i a^i = 1,$$

but the polynomial on the left has no constant term and we can pull out a factor of  $a$ .  $\square$