

Energy-Flow Cosmology (EFC) Formal Mathematical Specification

Morten Magnusson

November 14, 2025

Abstract

This document provides the complete mathematical specification of Energy-Flow Cosmology (EFC). It defines the entropy coordinate, the energy-flow field, the structural and dynamical equations, and all related constructs used in the computational implementation found in `src/efc/`.

Contents

1	Notation	2
2	Model Parameters	2
2.1	Entropy Parameters	2
2.2	Energy-Flow Parameters	2
2.3	Expansion Parameters	2
3	EFC-S: Entropy Field and Gradient	2
3.1	Entropy Field	3
3.2	Entropy Gradient	3
4	EFC-D: Energy-Flow Dynamics	3
4.1	Energy-Flow Field	3
4.2	Energy-Flow Potential	3
4.3	Circular Velocity Relation	3
5	EFC-H: Expansion Mapping	4
5.1	Baseline Expansion Law	4
5.2	Generalized Form	4
6	EFC-Structure: Halo Formation and Φ-Equilibria	4
6.1	Structural Equation	4
6.2	Entropy-Flow Coupling	4
7	Observables and Validation Mapping	4
7.1	JWST Predictions	4
7.2	DESI / BAO	5
7.3	SPARC Rotation Curves	5

1 Notation

The Energy-Flow Cosmology (EFC) framework uses the following central quantities:

- $S(\mathbf{x})$: entropy field or entropy coordinate.
- ∇S : entropy gradient, source of structural asymmetry.
- $E_f(\mathbf{x})$: energy-flow field; local density of directional energy flux.
- $\Phi(E_f, S)$: energy-flow potential governing structure.
- $H(E_f, S)$: expansion mapping derived from the flow-strength.
- $v(r)$: circular velocity used in SPARC validation.
- z : cosmological redshift.
- $\rho(\mathbf{x})$: matter/energy distribution (only used for comparison).

Coordinates:

- \mathbf{x} : spatial 3-vector.
- r : radial coordinate for rotation curve modelling.

2 Model Parameters

EFC uses a small set of structural and dynamical parameters. These are conceptually defined here; numerical defaults appear in `src/efc/parameters.py`.

2.1 Entropy Parameters

- S_{\min}, S_{\max} : bounds of the entropy coordinate.
- k_S : scaling factor controlling the strength of entropy gradients.

2.2 Energy-Flow Parameters

- $E_{f,0}$: reference scale for the energy-flow field.
- A_Φ : normalization constant for the energy-flow potential.

2.3 Expansion Parameters

- A_H : normalization constant for the expansion mapping H .
- α : optional exponent controlling non-linearity of the expansion law.

These parameters form the bridge between theory and numerics and determine the scaling of all observables.

3 EFC-S: Entropy Field and Gradient

The EFC-S sector defines the entropy field $S(\mathbf{x})$. This field encodes local structural organization and acts as a non-mass-based driver of curvature-like behavior.

3.1 Entropy Field

The entropy field is treated as a scalar function:

$$S : \mathbb{R}^3 \rightarrow [S_{\min}, S_{\max}]. \quad (1)$$

The field can be interpreted as a measure of information or organizational density in configuration space.

3.2 Entropy Gradient

The entropy gradient is:

$$\nabla S(\mathbf{x}) = \left(\frac{\partial S}{\partial x}, \frac{\partial S}{\partial y}, \frac{\partial S}{\partial z} \right). \quad (2)$$

The scaled physical gradient used in the model is:

$$\mathbf{g}_S = k_S \nabla S. \quad (3)$$

This gradient introduces directional asymmetry which couples to the energy-flow field and affects structure formation.

4 EFC-D: Energy-Flow Dynamics

The energy-flow field $E_f(\mathbf{x})$ represents the directional flow of energy across entropy gradients.

4.1 Energy-Flow Field

We define:

$$E_f : \mathbb{R}^3 \rightarrow \mathbb{R}. \quad (4)$$

Physically, E_f measures how strongly energy is "channeled" along structures created by gradients in S .

4.2 Energy-Flow Potential

EFC defines an effective potential:

$$\Phi(E_f, S) = A_\Phi E_f (1 + S). \quad (5)$$

This is the baseline form used in the Python implementation:

$$\Phi \propto E_f S + E_f. \quad (6)$$

More general forms (not yet implemented) include:

$$\Phi = A_\Phi E_f (1 + S^\beta), \quad (7)$$

where β controls non-linear coupling.

4.3 Circular Velocity Relation

For rotation curves:

$$v(r) = \sqrt{r \frac{\partial \Phi}{\partial r}}. \quad (8)$$

This connects the potential directly to SPARC data.

5 EFC-H: Expansion Mapping

The expansion rate H in EFC is not derived from density as in Λ CDM, but from the interaction of energy-flow and entropy.

5.1 Baseline Expansion Law

The current implementation defines:

$$H(E_f, S) = A_H \sqrt{|E_f|} (1 + S). \quad (9)$$

This links:

- entropy to redshift evolution,
- energy-flow to effective curvature,
- structural conditions to expansion behaviour.

5.2 Generalized Form

A more general version can be written as:

$$H(E_f, S) = A_H |E_f|^{\alpha/2} (1 + S)^\gamma, \quad (10)$$

with tunable exponents α, γ .

6 EFC-Structure: Halo Formation and Φ -Equilibria

Structure formation in EFC is governed by equilibrium conditions in the energy-flow potential.

6.1 Structural Equation

Stable configurations satisfy the condition:

$$\nabla \Phi(E_f, S) = 0. \quad (11)$$

This defines EFC halos.

6.2 Entropy–Flow Coupling

Including the entropy gradient gives:

$$\nabla \Phi + S \nabla S = 0. \quad (12)$$

This yields profiles of the form:

$$\Phi(r) = A_\Phi E_f(r) (1 + S(r)). \quad (13)$$

These profiles predict rotation curves without dark matter and large-scale distribution without a cosmological constant.

7 Observables and Validation Mapping

7.1 JWST Predictions

The structural growth function $G(E_f, S)$ determines the abundance and luminosity of early galaxies. Higher entropy gradients accelerate early structure formation.

7.2 DESI / BAO

The expansion law produces:

$$H(z) = H(E_f(z), S(z)). \quad (14)$$

This maps directly to DESI measurements of the expansion curve.

7.3 SPARC Rotation Curves

Given:

$$\Phi(r) = A_\Phi E_f(r)(1 + S(r)), \quad (15)$$

rotation curves follow:

$$v(r) = \sqrt{r \frac{d\Phi}{dr}}. \quad (16)$$

This is computed numerically in `efc_validation.py`.