

Cardiozen Cardiology Field Brief

Dosing and Competitive Differentiation — Internal Use Only

Overview

Cardiozen is an oral cholesterol-lowering therapy with once-daily dosing, indicated as adjunct to diet for primary hypercholesterolemia. This brief covers the dosing protocol the field is asked about most often and the approved differentiation messaging against the leading external competitor.

Dosing and Titration

Cardiozen is dosed orally once daily. Starting dose is 10 mg for most patients. Titration to 20 mg may be considered after 4 weeks if LDL-C reduction targets are not met. Maximum dose is 40 mg daily.

No dose adjustment is required for mild to moderate renal impairment. In severe hepatic impairment, the drug is contraindicated. Cardiozen may be taken with or without food, but consistency improves adherence. Co-administration with strong CYP3A4 inhibitors requires dose reduction to 5 mg daily.

Differentiation vs. External Competitor

Cardiozen differentiation versus the leading external competitor (referred to here as Product X): once-daily oral dosing versus Product X's twice-daily regimen; demonstrated 52% mean LDL-C reduction at 20 mg in the pivotal trial versus Product X's 41% at the equivalent dose; no required lipid panel monitoring after week 12 per the label, versus Product X's quarterly monitoring requirement; co-pay card with \$0 minimum versus Product X's \$25 minimum.

Reps must only compare on the approved messaging; off-label or unsupported comparisons are not permitted. The competitive grid on the rep portal is updated quarterly by medical affairs.