

Oncorix Field Reference Guide

For Pharmaceutical Sales Representatives — Internal Use Only

Overview

Oncorix is an oncology therapy approved for adult patients with advanced solid tumors. This reference guide consolidates the materials field representatives need most frequently: prior authorization process, dosing administration, the adverse event profile, patient assistance pathways, and approved responses to common field objections.

Reps should never offer medical advice or submit access paperwork on behalf of an office. Always refer clinical questions to medical affairs and direct office staff to the appropriate support program.

Prior Authorization Process

Prior authorization (PA) for Oncorix is required by most commercial and Medicare Part B plans for advanced solid tumor indications. The prescriber must supply: confirmed diagnosis with ICD-10 code and stage, biomarker testing results where indicated (HER2 status, MSI status), prior therapy history including progression on any first-line regimen, and the requested dosing schedule of 50 mg/m squared IV every 3 weeks.

Most payers respond within 5 to 7 business days. If denied, the appeal must include peer-reviewed literature supporting the use; the manufacturer's medical affairs team can supply a letter of medical necessity within 48 hours on request.

Dosing and Administration

Oncorix is administered as an IV infusion at 50 mg/m squared every 3 weeks for adult patients with advanced solid tumors. Premedication with a corticosteroid and antihistamine is required to reduce infusion-related reactions. Infusion duration is 60 minutes for the first cycle, 30 minutes thereafter if no reactions occurred.

Dose reductions for toxicity: first reduction to 40 mg/m squared, second to 30 mg/m squared; further reductions require discontinuation. Treatment continues until disease progression or unacceptable toxicity. No dose adjustment is required for mild renal or hepatic impairment; severe hepatic impairment is a contraindication.

Adverse Event Profile

The most frequently reported Oncorix adverse events from pivotal trial data: peripheral neuropathy (approximately 30% any grade, 8% grade 3 or higher), neutropenia (25% / 12% grade 3 or higher), fatigue (40%), nausea (35%), and infusion-related reactions in the first cycle (15%).

Grade 3 or higher peripheral neuropathy requires dose reduction per the dose-reduction algorithm in the prescribing information; persistent grade 3 neuropathy after dose reduction is a discontinuation criterion. Febrile neutropenia is a medical emergency and must be reported as a serious adverse

event.

Patient Assistance Program

The manufacturer offers a patient assistance program for eligible Oncorix patients without insurance coverage or with high out-of-pocket costs. Eligibility: US residents, household income at or below 500% of the federal poverty level, and documented insurance denial or lack of coverage. Approved patients receive Oncorix at no cost, shipped directly to the treating institution.

The co-pay assistance program for commercially insured patients caps out-of-pocket cost at \$25 per infusion, with an annual maximum of \$25,000. Reps may direct offices to the access support hotline; reps must not enroll patients themselves or handle patient PHI.

Common Objections and Approved Responses

PA is too painful — direct the office to the manufacturer's PA support program, which provides electronic submission and PA specialists who follow up directly with payers; average approval time has dropped to 4 days.

My patients can't afford it — review the co-pay assistance program (\$25 per infusion cap) and the PAP for uninsured patients.

I'm worried about peripheral neuropathy — reference the grade 3 management guide on the rep portal and offer to set up a peer-to-peer call with a medical science liaison.

I prefer once-weekly cisplatin — acknowledge clinician preference; the every-3-week Oncorix schedule reduces patient infusion-day visits by roughly two-thirds, a meaningful patient-experience differentiator for adherence and quality of life. Never disparage a competitor or make claims outside the FDA-approved label.