

Practical 10 : Computation of Eigen Value and
Eigen Vector for dimensionality reduction

i) Find the characteristic equation of the matrix

$$A = \begin{bmatrix} 2 & -3 & 1 \\ 3 & 1 & 3 \\ -5 & 2 & -4 \end{bmatrix}$$

Solⁿ :- Characteristic eqⁿ of A

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} 2-\lambda & -3 & 1 \\ 3 & 1-\lambda & 3 \\ -5 & 2 & -4-\lambda \end{vmatrix} = 0$$

$$= 2-\lambda \begin{vmatrix} 1-\lambda & 3 \\ 2 & -4-\lambda \end{vmatrix} + 3 \begin{vmatrix} 3 & 3 \\ -5 & -4-\lambda \end{vmatrix} + 1 \begin{vmatrix} 3 & 1-\lambda \\ -5 & 2 \end{vmatrix}$$

$$= 2-\lambda [-10 - \lambda + 4\lambda + \lambda^2] + 3 [-3\lambda + 3] + [6 + 5 - 5\lambda]$$

$$= 20 - 2\lambda + 8\lambda + 2\lambda^2 + 10\lambda + \lambda^2 - 4\lambda^2 - \lambda^3 - 9\lambda + 9 + 6 + 5 - 5\lambda$$

$$= -\lambda^3 - \lambda^2 + 2\lambda + 0$$

$$= -(\lambda^3 + \lambda^2 - 2\lambda)$$

ii) Find the eigen values of

$$A = \begin{bmatrix} 1 & -1 & 0 \\ -1 & 2 & 1 \\ 0 & 1 & 1 \end{bmatrix}$$

Solⁿ : $|A - \lambda I| = 0$

$$\begin{vmatrix} 1-\lambda & -1 & 0 \\ -1 & 2-\lambda & 1 \\ 0 & 1 & 1-\lambda \end{vmatrix} = 0$$

$$= 1-\lambda \begin{vmatrix} 2-\lambda & 1 \\ 1 & 1-\lambda \end{vmatrix} + 1 \begin{vmatrix} -1 & 1 \\ 0 & 1-\lambda \end{vmatrix} + 0$$

$$= 1-\lambda [1 - 3\lambda + \lambda^2] - 1 + \lambda$$

$$= 1 - 3\lambda + \lambda^2 - 1 + 3\lambda^2 - \lambda^3 - 1 + \lambda$$

$$= -\lambda^3 + 4\lambda^2 - 3\lambda$$

The eigen values for the given matrix are

$$\lambda = 3, \lambda = 1, \lambda = 0$$

iii) Find the eigen value and eigen Vector of the matrix

$$A = \begin{bmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ -1 & -2 & 0 \end{bmatrix}$$

Solⁿ: characteristic eqⁿ of A

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} -2-\lambda & 2 & -3 \\ 2 & 1-\lambda & -6 \\ -1 & -2 & -\lambda \end{vmatrix} = 0$$

$$= -2-\lambda \begin{vmatrix} 1-\lambda & -6 \\ -2 & -\lambda \end{vmatrix} - 2 \begin{vmatrix} 2 & -6 \\ 1 & -\lambda \end{vmatrix} - 3 \begin{vmatrix} 2 & 1-\lambda \\ -1 & -2 \end{vmatrix}$$

$$= -2-\lambda [-\lambda + \lambda^2 - 12] - 2(-2\lambda - 6) - 3(-4 + 1 - \lambda)$$

$$= +2\lambda - 2\lambda^2 + 24 + \lambda^2 - \lambda^3 + 12\lambda + 4\lambda + 12 + 12 - 3 + 3\lambda$$

$$= -\lambda^3 - \lambda^2 + 21\lambda + 45$$

eigen values

$$\lambda = 5, \quad \lambda = -3$$

for $\lambda = 5$

$$\begin{vmatrix} -2-\lambda & 2 & -3 \\ 2 & 1-\lambda & -6 \\ -1 & -2 & -\lambda \end{vmatrix} = \begin{vmatrix} -7 & 2 & -3 \\ 2 & -4 & -6 \\ 1 & -2 & -5 \end{vmatrix}$$

$$-7x + 2y - 3z = 0 \quad \text{--- ①}$$

$$2x - 4y - 6z = 0 \quad \text{--- ②}$$

$$1x - 2y - 5z = 0 \quad \text{--- ③}$$

using Cramer's rule ① & ②

$$\begin{vmatrix} x & -y \\ 2 & -3 \\ -4 & -6 \end{vmatrix} = \begin{vmatrix} -7 & -3 \\ 2 & -6 \end{vmatrix} = \begin{vmatrix} 2 & -7 \\ 2 & -4 \end{vmatrix}$$

$$x = -24$$

$$y = -48$$

$$z = 24$$

for $\lambda = 5$

$$\begin{bmatrix} -1 \\ -2 \\ 1 \end{bmatrix}$$

for $\lambda = -3$

$$\begin{vmatrix} -2-\lambda & 2 & -3 \\ 2 & 1-\lambda & -6 \\ -1 & -2 & -\lambda \end{vmatrix} = \begin{vmatrix} 1 & 2 & -3 \\ 2 & 4 & -6 \\ -1 & -2 & 3 \end{vmatrix}$$

$$1x + 2y - 3z = 0 \quad \text{--- (1)}$$

$$2x + 4y - 6z = 0 \quad \text{--- (2)}$$

$$-1x - 2y + 3z = 0 \quad \text{--- (3)}$$

using Cramer's rule (1) & (2)

$$\begin{vmatrix} x & -y \\ 2 & -6 \end{vmatrix} = \begin{vmatrix} 1 & -3 \\ 2 & -6 \end{vmatrix} = \begin{vmatrix} z & 2 \\ 1 & 2 \\ 2 & 4 \end{vmatrix}$$

$$x = 0$$

$$y = 0$$

$$z = 0$$

for $\lambda = -3$

$$\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

The eigen values for the given matrix is 5 and -3

and eigen vectors are $\begin{bmatrix} -1 \\ -2 \\ 1 \end{bmatrix}$ & $\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ respectively

iv) Find all the eigen values and eigen vector for the following matrix A

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 1 & 5 & 1 \\ 3 & 1 & 1 \end{bmatrix}$$

Solⁿ: Characteristic eqⁿ of A

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} 1-\lambda & 1 & 3 \\ 1 & 5-\lambda & 1 \\ 3 & 1 & 1-\lambda \end{vmatrix}$$

$$= (1-\lambda) \begin{vmatrix} 5-\lambda & 1 \\ 1 & 1-\lambda \end{vmatrix} - 1 \begin{vmatrix} 1 & 1 \\ 3 & 1-\lambda \end{vmatrix} + 3 \begin{vmatrix} 1 & 5-\lambda \\ 3 & 1 \end{vmatrix}$$

$$\begin{aligned}
 &= 1 - \lambda(4 - 5\lambda - \lambda + \lambda^2) - (1 - \lambda + -3) + 3(1 - 15 + 3\lambda) \\
 &= 4 - 5\lambda - \lambda + \lambda^2 - 4\lambda + 5\lambda^2 + \lambda^2 - \lambda^3 - 1 + \lambda + 3 + 3 - 45 + 9\lambda \\
 &= -\lambda + 7\lambda^2 - 36
 \end{aligned}$$

eigen values

$$\lambda = -2, \lambda = 6, \lambda = 3$$

for $\lambda = -2$

$$\begin{vmatrix} 1-\lambda & 1 & 3 \\ 1 & 5-\lambda & 1 \\ 3 & 1 & 1-\lambda \end{vmatrix} = \begin{vmatrix} 3 & 1 & 3 \\ 1 & 7 & 1 \\ 3 & 1 & 2 \end{vmatrix}$$

$$3x + y + 3z = 0 \dots \textcircled{1}$$

$$1x + 7y + 1z = 0 \dots \textcircled{2}$$

$$3x + 1y + 2z = 0 \dots \textcircled{3}$$

using cramer's rule on eqⁿ ① & ②

$$\begin{vmatrix} x & -y & z \\ 1 & 3 & 3 \\ 7 & 1 & 1 \end{vmatrix} = \begin{vmatrix} 3 & 3 \\ 1 & 1 \end{vmatrix} = \begin{vmatrix} 3 & 1 \\ 1 & 7 \end{vmatrix}$$

$$x = -20, -y = 0, z = +20$$

$$\text{for } \lambda = -2 \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$$

for $\lambda = 6$

$$\begin{vmatrix} -5 & 1 & 3 \\ 1 & -1 & 1 \\ 3 & 1 & -5 \end{vmatrix} = \begin{aligned} &-5x + y + 3z = 0 \dots \textcircled{1} \\ &x - y - z = 0 \dots \textcircled{2} \\ &3x + y - 5z = 0 \dots \textcircled{3} \end{aligned}$$

using cramer's rule on eqⁿ ① & ②

$$\begin{vmatrix} x & -y & z \\ 1 & 3 & 3 \\ -1 & -1 & 1 \end{vmatrix} = \begin{vmatrix} -5 & 3 \\ 1 & -1 \end{vmatrix} = \begin{vmatrix} -5 & 1 \\ 1 & -1 \end{vmatrix}$$

$$x = 4, +y = +8, z = 4$$

$$\text{for } \lambda = 6 \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$$

Practical 11 : Calculate Maxima and Minima for the given function

i) Find the absolute extreme points of a function

$$f(x) = 2x^3 - 3x^2 + 5 \text{ in the interval } [-2, 2]$$

Solⁿ: $f(x) = 2x^3 - 3x^2 + 5$

$$f'(x) = 6x^2 - 6x$$

$$f'(x) = 0$$

$$\therefore 6x^2 - 6x = 0$$

$$6x(x-1) = 0$$

$$6x = 0 \quad \text{or} \quad x = 1$$

$$x = 0$$

for $x = 0$

$$f(x) = 2x^3 - 3x^2 + 5$$

$$f(0) = 0 - 0 + 5$$

$$f(0) = 5$$

for $x = 1$

$$f(x) = 2(1)^3 - 3(1)^2 + 5$$

$$f(1) = 4$$

Checking endpoints of interval

$$x = -2$$

$$f(-2) = 2(-2)^3 - 3(-2)^2 + 5$$

$$f(-2) = -23$$

$$x = 2$$

$$f(2) = 2(2)^3 - 3(2)^2 + 5$$

$$f(2) = 9$$

The minimum value is -23 at $x = -2$ &

maximum value is 9 at $x = 2$

ii) Find the absolute maxima and minima of function $f(x) = 2\sin x + \cos x$ in the interval $[-1, 1]$

Solⁿ: $f(x) = 2\sin x + \cos x$

$$f'(x) = 2\cos x - \sin x$$

$$f'(x) = 0$$

$$2\cos x - \sin x = 0$$

$$2\cos x = \sin x$$

$$2 = \tan x$$

$$x = \tan^{-1}(2)$$

$$x \approx 1.1071$$

for $x \approx 1.1071$

$$f(x) = 2\sin x + \cos x$$

$$f(1.1071) = 2\sin(1.1071) + \cos(1.1071)$$

$$= 2(0.8940) + 0.4472$$

$$= 1.788 + 0.4472$$

$$= 2.2352$$

Checking endpoints of interval

for $x = -1$

$$f(-1) = 2\sin(-1) + \cos(-1)$$

$$= 2(-0.8415) + 0.5403$$

$$= -1.683 + 0.5403$$

$$= -1.1427$$

for $x = 1$

$$f(1) = 2\sin(1) + \cos(1)$$

$$= 2(0.8415) + 0.5403$$

$$= 1.683 + 0.5403$$

$$= 2.2233$$

The minimum value is -1.1427 at $x = -1$ &

maximum value is 2.2352 at $x \approx 1.1071$

iii) Find the absolute extreme points of function

$$f(x) = \log x \text{ in the interval } [-2, 2]$$

Solⁿ: $f(x) = \log x$

$$f'(x) = \frac{1}{x}$$

$$f'(x) = 0$$

$$\frac{1}{x} = 0$$

conclusion:

Thus, there are no critical points in the given interval $[-1, 1]$

\therefore No solution for x , because $\frac{1}{x}$ is ^{never} ~~negati~~ equal to zero for any value of x .

iv) Find the applied maxima & minima of

$$f(x) = x \log x$$

Solⁿ:- $f(x) = x \log x$

$$f'(x) = x \cdot 1 \cdot (\log x) + x \cdot \left(\frac{1}{x}\right)$$
$$= \log x + 1$$

$$f''(x) = \log x + 1$$

$$= \frac{1}{x}$$

$$f'(x) = 0$$

$$\log x + 1 = 0$$

$$\log x = -1$$

$$x = e^{-1}$$

$$\dots (\log_a x = y : a^y = x)$$

$$x = 0.3678$$

for $x = e^{-1}$ (i.e. $x = \frac{1}{e}$)

$$f''(x) = \frac{1}{e^{-1}}$$
$$= 2.7182 > 0$$

$$f''(x) > 0$$

$$f(x) = x \log x$$

$$= e^{-1} \log_e e^{-1}$$

$$= -0.3678$$

The function $f(x)$ has minimum value -0.3678 at $x = e^{-1}$

The function has no maximum value

v) Find applied maxima & minima of

$$f(x) = x^2 + \frac{16}{x}$$

Solⁿ:-

$$f'(x) = 2x + \left(-\frac{16}{x^2}\right)$$

$$= 2x - \frac{16}{x^2}$$

$$f''(x) = 2 - \left(-\frac{16 \times 2}{x^3}\right)$$

$$\dots \left(\because \frac{1}{x^n} = \frac{-n}{x^{n+1}}\right)$$

$$= 2 + \frac{32}{x^3}$$

$$f'(x) = 0$$

$$2x - \frac{16}{x^2} = 0$$

$$2x = \frac{16}{x^2} \quad \text{or } x=0$$

$$2x^3 = 16$$

$$x^3 = 8$$

$$x = 2$$

$$\text{for } x = 2$$

$$f''(x) = 2 + \frac{32}{x^3}$$
$$= 2 + \frac{32}{8}$$

$$f''(x) = 6$$

$$f''(x) > 0$$

$\therefore f(x)$ attains minimum at $x=2$

$$f(x) = x^2 + \frac{16}{x}$$

$$= 4 + \frac{16}{2}$$

$$= 12$$

\therefore The function $f(x)$ has minimum value 12 at $x=2$

The function has no maximum value.

Practical 12: Impute an example based on Rolle's
and Mean value Theorem

a) Verify Rolle's theorem

i) $f(x) = \sin x + \cos x + 7$ at interval $x \in [0, 2\pi]$

Solⁿ :- The given $f(x)$ is a polynomial function

hence it is continuous at $[0, 2\pi]$ and differentiable at $(0, 2\pi)$

$$f(0) = \sin x + \cos x + 7$$

$$= \sin 0 + \cos 0 + 7$$

$$= 0 + 1 + 7$$

$$f(0) = 8$$

$$f(2\pi) = \sin x + \cos x + 7$$

$$= \sin 2\pi + \cos 2\pi + 7$$

$$= 0 + 1 + 7$$

$$f(2\pi) = 8$$

$$\therefore f(0) = f(2\pi)$$

Thus the function $f(x)$ satisfy all the condition of Rolle's theorem.

There exist $c \in (1, 4)$ such that $f'(c) = 0$

$$f'(x) = \sin x + \cos x + 7$$

$$f(c) = \sin c + \cos c + 7$$

$$f'(c) = \cos c - \sin c + 0$$

$$f'(c) = 0$$

$$\cos c - \sin c = 0$$

$$\cos c = \sin c$$

they are equal at $c = \frac{\pi}{4}$

$$\therefore c = \frac{\pi}{4} \in [0, 2\pi]$$

Hence the Rolle's theorem is verified.

$$\text{ii) } f(x) = \sin \frac{x}{2} \quad ; \quad x \in [0, 2\pi]$$

Solⁿ:- The given function $f(x)$ is a polynomial function
hence it is continuous at $[0, 2\pi]$ and differentiable on $[0, 2\pi]$

$$f(0) = \sin \frac{x}{2}$$

$$= \sin \frac{0}{2}$$

$$f(0) = 0$$

$$f(2\pi) = \sin \frac{x}{2}$$

$$= \sin \frac{2\pi}{2}$$

$$f(2\pi) = 0$$

$$f(0) = f(2\pi)$$

Thus the function satisfy all the condition of Rolle's theorem
there exist $c \in (0, 2\pi)$ such that $f'(c) = 0$

$$f(c) = \sin \frac{c}{2}$$

$$f'(c) = \cos \frac{c}{2}$$

$$f'(c) = 0$$

$$\cos \frac{c}{2} = 0$$

$$(\because \cos \frac{\pi}{2} = 0)$$

$$\therefore c = \pi \in [0, 2\pi]$$

Hence the Rolle's theorem is verified

b) Verify Lagrange's Mean Value Theorem

$$i) f(x) = x^2 - 3x - 1; x \in \left[-\frac{11}{7}, \frac{13}{7}\right]$$

Solⁿ:- The given function $f(x)$ is a polynomial function
it is continuous at $\left[-\frac{11}{7}, \frac{13}{7}\right]$ and differentiable at $\left[-\frac{11}{7}, \frac{13}{7}\right]$

$$f(a) = f\left(-\frac{11}{7}\right) = x^2 - 3x - 1 \\ = \left(-\frac{11}{7}\right)^2 - 3\left(-\frac{11}{7}\right) - 1$$

$$f(a) = 6.1836$$

$$f(b) = f\left(\frac{13}{7}\right) = x^2 - 3x - 1 \\ = \left(\frac{13}{7}\right)^2 - 3\left(\frac{13}{7}\right) - 1$$

$$f(b) = -3.1224$$

$$f(c) = c^2 - 3c - 1$$

$$f'(c) = 2c - 3$$

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$2c - 3 = \frac{(-3.1225) - 6.1836}{\left(\frac{13}{7} - \left(-\frac{11}{7}\right)\right)}$$

$$2c - 3 = \frac{-9.306}{24/7}$$

$$14c - 21 = \frac{-9.306}{24}$$

$$14c - 21 = -0.38775$$

$$14c = -0.38775 + 21$$

$$c = \frac{20.61225}{14}$$

$$c = 1.4723 \in \left[-\frac{11}{7}, \frac{13}{7}\right]$$

$$\text{ii) } f(x) = 2x - x^2 \quad ; \quad x \in [0, 1]$$

Solⁿ:- The given function $f(x)$ is a polynomial function
it's continuous at $[0, 1]$ and differentiable at $(0, 1)$

$$f(0) = 2x - x^2 \\ = 2(0) - 0$$

$$f(a) = f(0) = 0$$

$$f(b) = f(1) = 2x - x^2 \\ = 2 - 1^2$$

$$f(b) = f(1) = 1$$

$$f(c) = 2c - c^2$$

$$f'(c) = 2 - 2c$$

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$2 - 2c = \frac{1 - 0}{1 - 0}$$

$$2 - 2c = 1$$

$$-2c = 1 - 2$$

$$2c = 1$$

$$c = \frac{1}{2} \in [0, 1]$$

Practical 13: Dot product and Norm of the Vector

i) Consider the Two vector $u = (1, 2, -1)$ & $v = (0, 1, 1)$

a) determine the dot product of u & v .

$$u \cdot v = (1 \cdot 0) + (2 \cdot 1) + (-1 \cdot 1)$$

$$= 0 + 2 - 1$$

$$u \cdot v = 1$$

b) determine the Norm of 'v'.

$$||x||^2 = \sqrt{x_1^2 + x_2^2 + x_3^2}$$

$$= \sqrt{0^2 + 1^2 + 1^2}$$

$$= \sqrt{2}$$

c) determine the norm of 'u'

$$||x||^2 = \sqrt{x_1^2 + x_2^2 + x_3^2}$$

$$= \sqrt{1^2 + 2^2 + (-1)^2}$$

$$= \sqrt{1 + 4 + 1}$$

$$= \sqrt{6}$$

ii) determine the angle between $a = (3, 2)$ and $b = (1, 7)$

$$a \cdot b = (3 \times 1) + (2 \times 7)$$

$$a \cdot b = 3 + 14$$

$$a \cdot b = 17$$

$$|a| = \sqrt{3^2 + 2^2}$$

$$= \sqrt{13}$$

$$|b| = \sqrt{1^2 + 7^2}$$

$$= \sqrt{50}$$

$$\theta = \cos^{-1} \left(\frac{a \cdot b}{|a| \cdot |b|} \right)$$

$$= \cos^{-1} \left(\frac{17}{\sqrt{13} \cdot \sqrt{50}} \right)$$

$$= \cos^{-1} (0.667)$$

$$\theta = 48.16^\circ$$

Practical 14 : Compute Inverse of Matrix by adjoint Method

i) Find the adjoint of Matrix

$$a) A = \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix}$$

$$\text{Sol}^n:- A_{11} = (-1)^2 \cdot a_{22} = 2$$

$$A_{12} = (-1)^3 \cdot a_{12} = -2$$

$$A_{21} = (-1)^3 \cdot a_{21} = 0$$

$$A_{22} = (-1)^4 \cdot a_{11} = 1$$

$$\text{adj}(A) = \begin{bmatrix} 2 & -2 \\ 0 & 1 \end{bmatrix}$$

$$b) A = \begin{bmatrix} 2 & -3 & 1 \\ 3 & 1 & 3 \\ -5 & 2 & -4 \end{bmatrix}$$

$$\text{Sol}^n:- C_{11} = (-1)^2 [(1 \times -4) - (3 \times 2)] = -10$$

$$C_{12} = (-1)^3 [(3 \times -4) - (5 \times 3)] = -3$$

$$C_{13} = (-1)^4 [(3 \times 2) - (1 \times -5)] = 11$$

$$C_{21} = (-1)^3 [(-3 \times -4) - (2 \times 1)] = -10$$

$$C_{22} = (-1)^4 [(2 \times -4) - (1 \times -5)] = -3$$

$$C_{23} = (-1)^5 [(2 \times 2) - (-5 \times -3)] = 11$$

$$C_{31} = (-1)^4 [(-3 \times 3) - 1] = -10$$

$$C_{32} = (-1)^5 [(2 \times 3) - (3)] = -3$$

$$C_{33} = (-1)^6 [(2) - (3 \times -3)] = 11$$

$$\text{Cofactor } A = \begin{bmatrix} -10 & -3 & 11 \\ -10 & -3 & 11 \\ -10 & -3 & 11 \end{bmatrix}$$

$$\text{adj}(A) = A^T$$

$$\text{adj}(A) = \begin{bmatrix} -10 & -10 & -10 \\ -3 & -3 & -3 \\ 11 & 11 & 11 \end{bmatrix}$$

ii) Find the inverse of a matrix by adjoint method

$$A = \begin{bmatrix} 9 & 2 & 1 \\ 5 & -1 & 6 \\ 4 & 0 & -2 \end{bmatrix}$$

Solⁿ:- Finding the Cofactors

$$C_{11} = (-1)^2 (2-0) = 2$$

$$C_{12} = (-1)^3 (-10-24) = 34$$

$$C_{13} = (-1)^4 (0+4) = 4$$

$$C_{21} = (-1)^3 (-4+0) = 4$$

$$C_{22} = (-1)^4 (-18-4) = -22$$

$$C_{23} = (-1)^5 (0-8) = 8$$

$$C_{31} = (-1)^4 (12+1) = 13$$

$$C_{32} = (-1)^5 (54-5) = -49$$

$$C_{33} = (-1)^6 (-9-10) = -19$$

$$\text{co-factor matrix of } A = \begin{bmatrix} 2 & 34 & 4 \\ 4 & -22 & 8 \\ 13 & -49 & -19 \end{bmatrix}$$

$$\text{adj}(A) = A^T$$

$$\text{adj}(A) = \begin{bmatrix} 2 & 4 & 13 \\ 34 & -22 & -49 \\ 4 & 8 & -19 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 9 & 2 & 1 \\ 5 & -1 & 6 \\ 4 & 0 & -2 \end{vmatrix} = 9(+2) - 2(10-24) + 1(4) \\ = +18 + 28 + 4 \\ = 49$$

$$\therefore |A| = 49$$

$$A^{-1} = \text{adj}(A) \cdot \frac{1}{|A|}$$

$$= \frac{1}{49} \begin{bmatrix} 2 & 4 & 13 \\ 34 & -22 & -49 \\ 4 & 8 & -19 \end{bmatrix}$$