

Uncalibrated Stereo



Shree K. Nayar

Columbia University



Topic: Uncalibrated Stereo, Module: Reconstruction II

First Principles of Computer Vision


Simple (Calibrated) Stereo





Simple (Calibrated) Stereo



Simple (Calibrated) Stereo




Simple (Calibrated) Stereo



Simple (Calibrated) Stereo

$$u_l = f_x \frac{x}{z} + o_x$$

$$v_l = f_y \frac{y}{z} + o_y$$



$$u_r = f_x \frac{x - b}{z} + o_x$$

$$v_r = f_y \frac{y}{z} + o_y$$


f_x, f_y, b, o_x, o_y are
in pixel units.



Simple (Calibrated) Stereo

$$u_l = f_x \frac{x}{z} + o_x$$

$$v_l = f_y \frac{y}{z} + o_y$$



$$u_r = f_x \frac{x - b}{z} + o_x$$

$$v_r = f_y \frac{y}{z} + o_y$$


f_x, f_y, b, o_x, o_y are
in pixel units.



Simple (Calibrated) Stereo

$$u_l = f_x \frac{x}{z} + o_x$$

$$v_l = f_y \frac{y}{z} + o_y$$



$$u_r = f_x \frac{x - b}{z} + o_x$$

$$v_r = f_y \frac{y}{z} + o_y$$

f_x, f_y, b, o_x, o_y are
in pixel units.



Depth and Disparity


Solving for (x, y, z) :

$$x = \frac{b(u_l - o_x)}{(u_l - u_r)}$$

$$y = \frac{bf_x(v_l - o_y)}{f_y(u_l - u_r)}$$

$$z = \frac{bf_x}{(u_l - u_r)}$$

where $(u_l - u_r)$ is called the **Disparity**.



Uncalibrated Stereo

Method to estimate 3D structure of a static scene from two arbitrary views.



Uncalibrated Stereo

Method to estimate 3D structure of a static scene from two arbitrary views.



Uncalibrated Stereo

Method to estimate 3D structure of a static scene from two arbitrary views.



Uncalibrated Stereo

Method to estimate 3D structure of a static scene from two arbitrary views.

Topics:

- (1) Problem of Uncalibrated Stereo



Uncalibrated Stereo

Method to estimate 3D structure of a static scene from two arbitrary views.

Topics:

- (1) Problem of Uncalibrated Stereo
- (2) Epipolar Geometry



Uncalibrated Stereo

Method to estimate 3D structure of a static scene from two arbitrary views.

Topics:

- (1) Problem of Uncalibrated Stereo
- (2) Epipolar Geometry




Uncalibrated Stereo

Method to estimate 3D structure of a static scene from two arbitrary views.

Topics:

- (1) Problem of Uncalibrated Stereo
- (2) Epipolar Geometry
- (3) Estimating Fundamental Matrix



Uncalibrated Stereo

Method to estimate 3D structure of a static scene from two arbitrary views.

Topics:

- (1) Problem of Uncalibrated Stereo
- (2) Epipolar Geometry
- (3) Estimating Fundamental Matrix




Uncalibrated Stereo

Method to estimate 3D structure of a static scene from two arbitrary views.

Topics:

- (1) Problem of Uncalibrated Stereo
- (2) Epipolar Geometry
- (3) Estimating Fundamental Matrix
- (4) Finding Dense Correspondences



Uncalibrated Stereo

Method to estimate 3D structure of a static scene from two arbitrary views.

Topics:

- (1) Problem of Uncalibrated Stereo
- (2) Epipolar Geometry
- (3) Estimating Fundamental Matrix
- (4) Finding Dense Correspondences
- (5) Computing Depth



Uncalibrated Stereo

Method to estimate 3D structure of a static scene from two arbitrary views.

Topics:

- (1) Problem of Uncalibrated Stereo
- (2) Epipolar Geometry
- (3) Estimating Fundamental Matrix
- (4) Finding Dense Correspondences
- (5) Computing Depth
- (6) Stereopsis: Stereo in Nature

