

Lecture 2: Sets of Numbers, Propositional Logic

<http://book.imt-decal.org>, Ch. 1.3, 1.4

Introduction to Mathematical Thinking


September 12th, 2018

Suraj Rampure

Announcements

- Midterm date changed to **Wednesday, October 10th** (because of CS 61A midterm). No conflicting dates with 61A/61B/16A/16B.
- Homework 2 will be released after lecture, and is due Monday at 6:30PM.
- Last week's lecture video is up. Today's should be up by soon (done with half of it), along with a walkthrough of the homework problems.
- Office hours are official: Tuesday 2-3PM and Thursday 5-6PM, both in Cory 299.

$$2 \neq -2$$



Recap from last lecture

We studied three types of functions.

- **Injections (one-to-one):** No two elements in the domain map to the same element in the codomain

$$f(x_1) = f(x_2) \implies x_1 = x_2$$

- **Surjections (onto):** Every element in the codomain is mapped to by some element in the domain

all possible outputs { codomain = range } set of all actual outputs

- **Bijections:** Both injective and surjective

1.3: Sets of Numbers

We've seen the following sets of numbers before:

- Natural numbers
- Whole numbers
- Integers
- Rational numbers
- Irrational numbers
- Real numbers
- Complex numbers

We want to determine the relative sizes of each of these sets.

$|S|$: # of unique elements in S

$$|\{1, 2, 2, 8\}| = 3 \quad |\{1, 2, 3, 4, 5, \dots, 3\}| = ? \cdot ? \cdot ?$$

Let's extend our definition of cardinality to support sets with infinitely many elements:

Definition: Cardinality

We say two sets have the same cardinality if and only if there exists a bijection between the two sets.

$$\begin{array}{ccc} A = \{a_1, a_2, a_3\} \\ \downarrow \quad \downarrow \quad \downarrow \\ B = \{b_1, b_2, b_3\} \end{array}$$

this definition holds
for finite sets
as well

Natural Numbers

Definition: Natural numbers

The **natural numbers** (also known as the counting numbers), denoted by \mathbb{N} , are the most primitive numbers; ones that occur trivially in nature that can be used to count a (non-zero) number of things.

$$\mathbb{N} = \{1, 2, 3, 4, \dots\}$$

$$\setminus \mathbb{N} \in \mathbb{N}$$


Whole Numbers

Definition: Whole numbers

The set of **whole numbers**, denoted by \mathbb{N}_0 , is the union of the set of counting numbers with the number 0.

$$\mathbb{N}_0 = \{0, 1, 2, 3, 4, \dots\} = \{0\} \cup \mathbb{N}$$

Is there a bijection between the natural numbers and whole numbers?



yes bijection

$$f(x) = x - 1$$

Injective

$$f(a) = f(b) \implies a = b$$

$$a - 1 = b - 1 \implies a = b$$

$$|\mathbb{N}| = |\mathbb{N}_0|$$

Surjective

$$f: \mathbb{C} + 1 \mapsto \mathbb{C}$$

every whole number

eventually appears

$$f: \mathbb{A} \rightarrow \mathbb{B}$$

$$f: x \mapsto x^2$$

$$\mathbb{N} = \{1, 2, 3, 4, \dots\}$$

\mathbb{N}	S
1	s_1
2	s_2
3	s_3
\vdots	\vdots

Definition: Countably infinite

We say set S is **countably infinite** if and only if there exists a bijection from the natural numbers (or any other countable set) to S . If such a bijection does not exist, we say S is **uncountably infinite**.

One way to think of this is to give each number a waiting number in an infinitely long line!

there exists
an ordering of the set

$$\begin{aligned} f: \mathbb{N} &\rightarrow S \\ f: \mathbb{N}_0 &\rightarrow S \\ \text{OR } f: S &\rightarrow \mathbb{N}, \\ &S \rightarrow \mathbb{N}_0 \end{aligned}$$

$$\mathbb{N} \subset \mathbb{N}_0 \subset \mathbb{Z}$$

Integers

Definition: Integers

The set of integers, denoted by \mathbb{Z} , is the union of the whole numbers with their negatives


$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

Are the integers countably infinite, or uncountably infinite?

yes!

\mathbb{N}_0

\mathbb{Z}



$$f(x) = \begin{cases} -\frac{x}{2} & x \text{ is even} \\ \frac{x+1}{2} & x \text{ is odd} \end{cases}$$

eventually will get to any integer I want $\xrightarrow{\text{BIJECTION}}$

Rational Numbers

$$\mathbb{N} \subset \mathbb{N}_0 \subset \mathbb{Z} \subset \mathbb{Q}$$

Definition: Rational numbers



The set of rational numbers, denoted by \mathbb{Q} , is the set of all possible combinations of one integer divided by another, with the latter integer being non-zero.

$$\mathbb{Q} = \left\{ \frac{p}{q} : p, q \in \mathbb{Z}, q \neq 0 \right\}$$

rational numbers

are
DENSE

Are the rational numbers countably infinite, or uncountably infinite?



$$\frac{1}{5}, \frac{1}{6}, \dots, \frac{1}{10^{10}}, \dots$$

$$1) f: \underbrace{\mathbb{N}}_{\sim} \rightarrow \underbrace{\mathbb{Q}}_{\sim}$$

$$f: x \mapsto x$$

This is an injection!

$$2) f: \mathbb{Q} \rightarrow \mathbb{N} \quad : \text{next slide}$$

$$A = \{a_1, a_2, a_3\}$$

$$B = \{b_1, b_2, b_3, b_4\}$$

$$\text{since } |A| \leq |B|,$$


we can map each
element in A to
a diff. element in B

position

$$(a, b) \rightarrow \frac{b}{a}$$

Injection from $\mathbb{Q} \rightarrow \mathbb{N}$:

\mathbb{Q}	\mathbb{N}
0	1
1	2
-1	3
-1/2	4
1/2	5
	6
	⋮




injection
 $\mathbb{N} \rightarrow \mathbb{Q}$

injection
 $\mathbb{Q} \rightarrow \mathbb{N}$

\therefore
bijection
 $\mathbb{N} \rightarrow \mathbb{Q}$

\therefore
 \mathbb{Q} countably
infinite



Definition: Real numbers

The set of real numbers, denoted by \mathbb{R} , is the set of all possible distances from 0 on a number line

$$\mathbb{R} = \{3, \pi, -\sqrt{63}, 0.1224, \frac{2}{3}, \dots\}$$

Definition: Irrational numbers

The set of irrational numbers, denoted by $\mathbb{R} - \mathbb{Q}$, is the set of real numbers that are not rational. That is, they are real numbers that cannot be written as an integer divided by another integer.

$$\mathbb{R} - \mathbb{Q} = \{\pi, -e, \sqrt{5}, \dots\}$$

Cantor's Diagonalization

Real Numbers


Rational Numbers

Integers

Whole Numbers

Natural
Numbers

Irrational
Numbers



1.4: Propositional Logic

$2 + 2$ not a
proposition

Definition: Proposition

A proposition is a statement that has a definitive value - either true or false.

Are the following statements propositions?

- ✓ • "13 is prime"
- ✓ • " x is prime" \rightarrow $P(x)$ proposition that depends on x
- ✓ • "it is 93 degrees outside right now"
- ✗ • "LeBron James is the greatest basketball player of all time"

I think it's true but not a
proposition

Logical Operators

Lots of parallels to set theory!

1. **Conjunction:** $A \wedge B$, read " A and B "

similar to intersection

$$A \wedge B = \{ x : x \in A \wedge x \in B \}$$

2. **Disjunction:** $A \vee B$, read " A or B "

similar to union

$$A \vee B = \{ x : x \in A \vee x \in B \}$$

3. **Negation:** $\neg A$, read "not A "

similar to complement

$$A^c = \{ x : \neg (x \in A) \}$$

We can use conjunctions, disjunctions and negations to create more complicated logical statements.

$$U(x) : x < 100$$

$$E(x) : x \text{ is even}$$

$$P(x) : x \text{ is prime}$$

$$\underbrace{U(x)}_{x < 100} \quad \wedge \quad \underbrace{(E(x) \vee P(x))}_{x \text{ is even or prime}}$$

Implications

$P \Rightarrow Q$, read " P implies Q ", says that if P is true, then Q must be true; if P is false, it says nothing about Q (Q could either be true or false).

P

For example, if all we know about today's date is that it's Christmas, we also know that the current month is December. However, if we don't know that it's Christmas, then it may or may not be December.

Truth Tables

We claim that $P \rightarrow Q \equiv \neg P \vee Q$.

"equivalent"
→ they have the same value

P	Q	$P \rightarrow Q$	$\neg P \vee Q$
True	True	True ✓	True
True	False	False ✓	False
False	True	True ✓	True
False	False	True ✓	True

This suffices as a proof!