



Research Article

Evaluating parametric and non-parametric quantile delta mapping for CMIP6 precipitation bias correction: independent validation in coastal Thailand

Abstract: Bias correction of climate-model precipitation is often assessed in the calibration period, which can overstate transferability. This study compared annual and month-wise parametric and non-parametric quantile delta mapping (QDM) for daily rainfall from five CMIP6 models at 12 stations in Prachuap Khiri Khan, Thailand. Transfer functions were fitted in 1981–2000 and frozen before independent validation in 2001–2014. Parametric candidates were selected by maximum-likelihood Akaike information criterion (AIC); robustness tests used small-sample AIC (AICc), wet-day thresholds of 0.1, 0.5, and 1.0 mm day⁻¹, and alternative upper-tail caps. At the primary 1.0-mm threshold and five-times cap, parametric monthly QDM reduced mean monthly RMSE from 116.20 to 105.06 mm and increased mean Kling-Gupta efficiency (KGE) from 0.229 to 0.337; non-parametric monthly QDM yielded 109.45 mm and 0.329. Their paired median KGE difference was 0.0002, and the ranking reversed without the cap. AICc reproduced every AIC selection. Non-parametric monthly KGE remained 0.329–0.332 across wet-day thresholds, whereas parametric monthly KGE decreased to 0.217 at 0.1 mm. Non-parametric annual QDM gave the lowest mean absolute 99th-percentile bias (24.79% versus 52.98% raw). The 60 model-station pairs share model grid cells and are descriptive paired evaluations, not independent spatial replicates. Month-wise calibration is supported for monthly water-balance applications, but parametric gains remain conditional on occurrence and tail safeguards; empirical monthly QDM is the more assumption-lean default.

Keywords: bias correction; quantile delta mapping; CMIP6; independent validation; coastal Thailand

1. Introduction

Daily rainfall controls water-resources planning, flood and drought risk, agricultural operations, and climate adaptation in Southeast Asia. CMIP6 simulations provide physically based climate information, but precipitation biases remain because of coarse resolution and imperfect representations of convection, topography, and regional circulation [1, 2]. These limitations are especially consequential in Prachuap Khiri Khan, where monsoon seasonality, a narrow coastal plain, and the Tenasserim Range generate strongly intermittent and skewed rainfall.

Quantile mapping is widely used to align simulated and observed distributions [3–6]. Among these methods, Quantile Delta Mapping (QDM) [7] was specifically developed to preserve modeled relative changes at each quantile, a property recognized as essential for retaining long-term climate-change signals during bias adjustment [8]. Method performance nevertheless depends on reference uncertainty, spatial mismatch, dry-day treatment, and the statistical representation of extremes [9, 10]. Recent global and multi-region studies likewise show that the preferred method varies with climate, terrain, variable, and impact metric [10–12]. Direct holdout comparisons of parametric and empirical QDM, including occurrence and tail sensitivity, remain limited for narrow tropical coastal provinces.

Three unresolved issues motivate the present comparison. First, evaluation in the calibration period measures construction-dependent agreement and can overstate out-of-sample performance. Second, univariate quantile mapping adjusts marginal distributions but cannot reconstruct missing event timing or wet-spell persistence [13]. Third, finite samples make upper-tail behavior sensitive to extrapolation and parameterization [8, 14]. These issues are particularly relevant when monthly stratification improves seasonality but reduces the wet-day sample available for fitting.

This study therefore compares four multiplicative QDM variants: non-parametric annual, non-parametric monthly, parametric annual, and parametric monthly. All observation-dependent quantities are fitted in 1981–

2000, frozen, and applied to 2001–2014. The primary endpoint is validation-period monthly Kling-Gupta efficiency (KGE); secondary endpoints characterize distributional fidelity, upper-tail bias, seasonal structure, and temporal correspondence. The objective is to determine which conclusions are transferable, which depend on numerical safeguards, and which uses of the corrected rainfall remain defensible.

2. Materials and Methods

2.1 Study area, observations, and CMIP6 simulations

Located in coastal Thailand, Prachuap Khiri Khan is a narrow province situated between the Gulf of Thailand and the Tenasserim Range. The local climate exhibits distinct spatial variations, heavily influenced by its complex topography and seasonal monsoons that generate highly intermittent and skewed rainfall patterns across the region [15]. To capture this spatial variability, the analysis utilized daily rainfall records from a 12-station inventory distributed throughout the province (Figure 1; Table 1). The observational dataset covers a continuous, common period from 1 January 1981 through 31 December 2014 (12,418 days).




Figure 1. Location of Prachuap Khiri Khan Province and the 12 rainfall stations used in the analysis.

The elevation surface in Figure 1 is provided only as regional topographic context. Station elevations in Table 1 are reproduced from the supplied station metadata; no DEM-derived elevations were substituted where station metadata were unavailable.

All stations had complete daily records. Declared missing codes were masked before negative-value handling. Negative values between -0.1 and 0 mm day^{-1} were set to zero, and values below -0.1 mm day^{-1} were treated as missing; no such corrections were required in the observation file used here. Annual totals were screened with the Pettitt change-point test at $\alpha = 0.05$ [16]. No station rejected the null hypothesis ($p = 0.059\text{--}1.000$), which is evidence of no detected change point rather than proof of homogeneity. No homogenization was applied.

The CMIP6 input comprised daily precipitation (pr) from the historical experiment for ACCESS-ESM1-5, CESM2, CanESM5, EC-Earth3, and MIROC6. The source, experiment, member, grid, variable, frequency, and date range are encoded in the analysis-ready filenames and summarized in Table 2; model descriptions follow the corresponding publications [17–21]. The EC-Earth3 gr identifier is retained exactly as recorded in the analysis-ready filename. These identifiers were cross-checked against the federated Earth System Grid Federation (ESGF) catalogue. Because ESGF datasets can be replicated across data nodes, the complete CMIP6 facets are used as the stable source description rather than a transient mirror endpoint. Values were already in mm day^{-1} ; the workflow checks units and would multiply flux units in $\text{kg m}^{-2} \text{s}^{-1}$ by 86,400. The station-series files contain two or three distinct model series across the 12 gauges, consistent with nearest native-grid extraction and shared grid cells. A machine-readable manifest records the exact derivative filenames, byte sizes, and SHA-256 checksums used in the analysis.

Standard-calendar models and observations contain 7,305 calibration and 5,113 validation days. CESM2 and CanESM5 use a 365-day calendar, with 7,300 and 5,110 days. Analysis used dates common to each model and the observations; 29 February was omitted for 365-day models and was not interpolated. The 60 model-station combinations are therefore paired evaluation units, not 60 independent spatial experiments, because stations can share model grid cells and the models share components and development lineages.

Table 1. Rain-gauge station characteristics for 1981–2014.

Station	Lat. (°N)	Long. (°E)	Elev. (m)	Miss. (%)	Annual rain. (mm)	SDII (mm day ⁻¹)	Max. daily rain. (mm day ⁻¹)	Lag-1 ACF
500001	12.37	99.95	9.1	0.0	984	12.0	170.0	0.21
500002	11.18	99.55	n/a	0.0	1,376	11.1	155.0	0.22
500003	12.02	99.90	n/a	0.0	1,100	13.1	181.3	0.19
500004	11.47	99.65	n/a	0.0	1,106	11.4	152.0	0.21
500005	11.60	99.68	5.0	0.0	1,147	6.4	143.5	0.41
500006	11.62	99.67	23.0	0.0	1,203	6.5	201.1	0.35
500007	11.90	99.78	21.3	0.0	1,134	6.2	176.6	0.35
500008	12.53	99.55	130.3	0.0	1,206	5.1	68.0	0.66
500009	12.22	99.87	14.4	0.0	1,177	5.2	54.2	0.63
500201	11.83	99.83	n/a	0.0	1,108	12.1	296.9	0.25
500202	12.59	99.96	n/a	0.0	944	11.8	298.5	0.26
500301	12.58	99.73	101.8	0.0	1,065	11.4	226.0	0.25

Note: n/a indicates that elevation metadata were absent from the supplied station file. SDII is mean rainfall on wet days ($\geq 1.0 \text{ mm day}^{-1}$), and ACF is lag-1 autocorrelation.

Table 2. CMIP6 historical simulations and metadata recovered from the analysis-ready files.

Model	Institution	Variant	Grid	Cal.	Days	Series	Ref.
ACCESS-ESM1-5	CSIRO	r1i1p1f1	gn	Std.	12,418	2	[21]
CESM2	NCAR	r1i1p1f1	gn	365-day	12,410	3	[18]

Model	Institution	Variant	Grid	Cal.	Days	Series	Ref.
CanESM5	CCCma	r1i1p1f1	gn	365-day	12,410	2	[17]
EC-Earth3	EC-Earth Consortium	r1i1p1f1	gr	Std.	12,418	3	[19]
MIROC6	MIROC Consortium	r1i1p1f1	gn	Std.	12,418	2	[20]

Note: Grid identifiers (gn/gr) are retained as recorded in the analysis-ready files. Distinct series is the number of unique daily station-series signatures across 12 gauges and indicates shared native grid cells; it is not a formal effective sample size.

2.2 Split-sample protocol and wet-day adaptation

Calibration covered 1981–2000 and independent validation covered 2001–2014. Every observed threshold, empirical distribution, and parametric calibration distribution was estimated from calibration data only and then frozen. Validation observations were used solely to calculate evaluation metrics.

Wet-day occurrence and positive intensity were corrected separately. A day was wet when observed rainfall was at least 1.0 mm day⁻¹, consistent with the Expert Team on Climate Change Detection and Indices convention used for wet-day precipitation indices [22]. For each model, station, and calibration group, p_{dry} was the fraction of observed calibration days below 1.0 mm day⁻¹. The model threshold was the empirical p_{dry} quantile of all model calibration values. Values at or below that threshold were set to zero; with ties, this rule ensures that the adapted model wet-day count does not exceed the observed count. The threshold was reused unchanged in validation. Annual variants used one threshold per model-station pair, whereas monthly variants used calendar-month-specific thresholds. Throughout this paper, “monthly QDM” denotes calendar-month calibration with the prespecified annual fallback for blocks containing fewer than 30 wet days. The full workflow was also rerun at 0.1 and 0.5 mm day⁻¹ to quantify threshold dependence.

A minimum of 30 wet days was required for a monthly transfer function. Of 720 model-station-month blocks, 100 dry-season blocks did not meet this threshold. These blocks were not discarded or set to zero; they used the corresponding annual transfer function fitted from calibration data. This fallback was declared uniformly and recorded in the diagnostics.

2.3 Multiplicative QDM

For a model precipitation value x at time t in target period p , the multiplicative Quantile Delta Mapping (QDM) procedure was applied following Cannon et al. [7].

$$\tau(t) = F_{m,p}[x_{m,p}(t)] \quad (1)$$

$$\Delta(t) = \frac{x_{m,p}(t)}{F_{m,h}^{-1}[\tau(t)]} \quad (2)$$

$$\hat{x}(t) = F_{o,h}^{-1}[\tau(t)] \times \Delta(t) \quad (3)$$

Here, $F_{m,p}$ is the model cumulative distribution function in the target period. The terms $F_{m,h}^{-1}$ and $F_{o,h}^{-1}$ represent the model and observed calibration-period quantile functions (inverse cumulative distributions), respectively. The variable τ is the non-exceedance probability, Δ represents the modeled relative-change ratio, and $\hat{x}$ is the final bias-corrected rainfall. Because the target-period probability $\tau(t)$ is calculated using model values only, it does not violate the independent observation holdout. Retaining the relative-change ratio Δ is the fundamental characteristic that distinguishes QDM from ordinary quantile mapping methods.

To prevent division by very small historical quantiles, the ratio Δ was bounded a priori to the interval [0.2,5.0]. For the primary parametric analysis, the corrected rainfall was capped at five times the maximum observed wet-day rainfall in the corresponding calibration block. Both the ratio bounds and the upper cap act strictly as numerical safeguards, not as hydrological return-level thresholds. The cap was applied after the mathematical transformation. Sensitivity runs were also conducted using a stricter factor of 1.5 and a

completely uncapped scenario; all lower-bound clipping and upper-bound capping events were systematically counted for evaluation.

2.4 Parametric and non-parametric variants

For the non-parametric variants, the empirical cumulative distribution functions (eCDF) were estimated using the plotting position formula:

$$F(x_i) = \frac{i-0.5}{n} \quad (4)$$

where i is the rank of the data point in ascending order and n is the total sample size of wet days. Linear interpolation was applied between these plotting positions to estimate intermediate quantiles. Annual variants pooled all months to form a single distribution, whereas monthly variants fitted the empirical distribution for each calendar month separately.

For the parametric variants, the maximum likelihood estimation (MLE) method fitted five theoretical probability distributions to the wet-day intensities (x), including gamma, generalized extreme value (GEV), Weibull minimum, lognormal, and Pearson type III.

To prevent generating precipitation values below zero, the location parameters for the gamma, Weibull minimum, and lognormal distributions were strictly fixed at zero, reducing them to their two-parameter forms. The probability density functions (PDFs) for these distributions are defined in Table 3.

Table 3. Theoretical probability distributions and parameters for modeling wet-day precipitation intensities

Distribution (Parameter Form)	Probability Density Function (PDF)	Parameters & Conditions
Gamma (2-parameter)	$f(x) = \frac{x^{k-1} e^{-x/\theta}}{\theta^k \Gamma(k)}$	k : shape parameter θ : scale parameter $\Gamma(\cdot)$: gamma function Note: Location parameter is strictly fixed at zero to prevent negative values.
Weibull Minimum (2-parameter)	$f(x) = \frac{c}{\lambda} \left(\frac{x}{\lambda}\right)^{c-1} \exp\left[-\left(\frac{x}{\lambda}\right)^c\right]$	c : shape parameter λ : scale parameter Note: Location parameter is strictly fixed at zero to prevent negative values.
Lognormal (2-parameter)	$f(x) = \frac{1}{x\sigma\sqrt{2\pi}} \exp\left[-\frac{(\ln x - \mu)^2}{2\sigma^2}\right]$	σ : shape parameter (standard deviation of the log) μ : scale parameter (mean of the log) Note: Location parameter is strictly fixed at zero to prevent negative values.
Generalized Extreme Value (GEV) (3-parameter)	$f(x) = \frac{1}{\alpha} \left[1 + \kappa \left(\frac{x - \xi}{\alpha}\right)\right]^{\frac{1}{\kappa}-1} \exp\left\{-\left[1 + \kappa \left(\frac{x - \xi}{\alpha}\right)\right]^{\frac{1}{\kappa}}\right\}$	ξ : freely estimated location parameter α : scale parameter κ : shape parameter Condition: The function is defined for $1 + \kappa(x - \xi)/\alpha > 0$.
Pearson Type III (3-parameter)	$f(x) = \frac{1}{ \beta \Gamma(\alpha)} \left \frac{x - \xi}{\beta}\right ^{\alpha-1} \exp\left(-\frac{x - \xi}{\beta}\right)$	ξ : freely estimated location parameter β : scale parameter

Distribution (Parameter Form)	Probability Density Function (PDF)	Parameters & Conditions
		α : shape parameter Condition: Formulation allows for both positive and negative skewness in precipitation data.

The optimal probability distribution for each calibration block was deterministically selected based on the minimum Akaike Information Criterion (AIC), which penalizes model complexity to prevent overfitting. The AIC is defined as:

$$AIC = 2k - 2 \ln(L) \quad (5)$$

where k is the number of freely estimated parameters and L represents the maximized value of the likelihood function for the fitted distribution. Candidates resulting in non-finite parameters or non-finite likelihoods were strictly disqualified.

To account for small-sample sensitivity during the month-wise calibration, the distribution selection and the complete correction workflow were repeated using the corrected Akaike Information Criterion (AIC_c):

$$AIC_c = AIC + \frac{2k(k+1)}{n-k-1} \quad (6)$$

where n is the sample size. The Kolmogorov-Smirnov statistic was stored as a goodness-of-fit diagnostic for all combinations. It should be noted that the GEV distribution was included strictly as a flexible, whole-distribution candidate, rather than asserting that daily wet-day rainfall constitutes a block-maximum extreme value sample. All computational procedures and maximum likelihood optimizations were executed using Python 3.12.3 and the SciPy 1.17.1 library [23].

2.5 Evaluation endpoints and uncertainty

The primary evaluation endpoint was the monthly Kling-Gupta efficiency (KGE) [24], selected because it explicitly combines linear correlation, variability, and mean bias into a single objective metric. The KGE is mathematically defined as:

$$KGE = 1 - \sqrt{(r-1)^2 + (\alpha-1)^2 + (\beta-1)^2} \quad (7)$$

where r is the Pearson correlation coefficient between the simulated (S) and observed (O) monthly rainfall totals, $\alpha = \sigma_S/\sigma_O$ represents the ratio of their standard deviations (variability), and $\beta = \mu_S/\mu_O$ represents the ratio of their means (bias).

Secondary endpoints included monthly root mean square error (RMSE), mean absolute error (MAE), percent bias (PBIAS), absolute relative bias of the 99th wet-day percentile (q99 bias), the Kolmogorov-Smirnov statistic (D) for wet-day distributions, monthly Pearson correlation (r), and correlation between the 12 monthly climatological means. These metrics are defined in Equations (8)–(14).

$$RMSE = \sqrt{\frac{1}{N} \sum_{i=1}^N (S_i - O_i)^2} \quad (8)$$

$$|q_{99} \text{ bias}| = \left| \frac{q_{99,S} - q_{99,O}}{q_{99,O}} \right| \times 100\% \quad (9)$$

$$D = \sup_x |F_S(x) - F_O(x)| \quad (10)$$

Here, N is the number of paired monthly records; S_i and O_i are simulated and observed monthly totals; and q_{99} is the 99th percentile of wet-day intensity. In Equation (10), D is the supremum absolute difference between the simulated and observed empirical cumulative distribution functions of wet-day precipitation. The climatology correlation is calculated from the 12 month-specific means, not from independent observations. KGE and correlations are better at larger values, whereas the remaining endpoints are better at smaller values. Direction-based rankings are labeled descriptive and do not imply statistical superiority.

$$PBIAS = 100 \times \frac{\bar{S} - \bar{O}}{\bar{O}} \quad (11)$$

$$\text{MAE} = \frac{1}{N} \sum_{i=1}^N |S_i - O_i| \quad (12)$$

$$r = \frac{\sum_{i=1}^N (S_i - \bar{S})(O_i - \bar{O})}{\sqrt{\left(\sum_{i=1}^N (S_i - \bar{S})^2\right) \left(\sum_{i=1}^N (O_i - \bar{O})^2\right)}} \quad (13)$$

$$r_{\text{clim}} = \text{corr}(\{\bar{S}_1, \dots, \bar{S}_{12}\}, \{\bar{O}_1, \dots, \bar{O}_{12}\}) \quad (14)$$

The PBIAS and MAE equations use the same paired monthly records as RMSE; r is the Pearson correlation of monthly totals; and the climatology correlation is calculated from the 12 simulated and observed monthly climatological means.

Uncertainty in corrected-minus-raw differences was estimated separately for each model-station combination using 2,000 paired year-block bootstrap replicates with a fixed random seed (seed 42). Validation monthly totals were structured as a 14-year by 12-month matrix. Fourteen years were sampled with replacement in each replicate, and the exact same year indices were applied simultaneously to observations, raw simulations, and all corrected variants to preserve temporal correlation. The 95% interval was extracted from the 2.5th and 97.5th percentiles of the bootstrap distribution. These intervals and favorable-point counts are descriptive pair-specific summaries; no network-wide p-value or multiple-testing claim is made because the 60 combinations are spatially and structurally dependent. The fit-freeze-apply sequence is summarized in Figure 2.




Figure 2. Fit-freeze-apply framework. Observation-dependent transfer functions were estimated in 1981–2000 and applied without re-estimation in 2001–2014; uncertainty used paired year-block bootstrap resampling.

3. Results and Discussion

3.1 Observed rainfall and model context

Mean annual observed rainfall ranged from 944 to 1,376 mm among stations, and maximum daily rainfall ranged from 54.2 to 298.5 mm day⁻¹. Calibration and validation retained the same broad bimodal seasonal cycle, but several stations changed wet-day frequency between periods (Figure 3). This non-stationarity makes the holdout test more demanding than same-period evaluation.

Table 1 and Figure 3 show two distinct observational structures. Stations 500005–500009 have lower SDII (5.1–6.5 mm day⁻¹) and substantially higher wet-day frequencies than the other stations, while 500008 and 500009 also have lower recorded daily maxima (68.0 and 54.2 mm, respectively). These contrasts may reflect genuine gauge-regime differences or data-handling practices, but the available metadata do not permit

attribution. All stations were therefore retained, and station-level comparisons are interpreted descriptively rather than as independent spatial evidence.

The five model files represented only two or three distinct native-grid series across the 12 stations (Table 2). Consequently, averaging across all 60 combinations would overstate spatial replication. Results are therefore reported as paired model-station distributions and counts, with this dependence carried into interpretation.




Figure 3. Observed rainfall in calibration and validation: (a) monthly climatology with interquartile ribbons across stations, (b) annual rainfall, and (c) paired station-level wet-day frequencies. Colored connecting lines show the direction of change; these summaries are descriptive and motivate the independent holdout test.

3.2 Independent validation performance

Month-wise calibration produced the clearest descriptive gains (Table 4; Figure 4). P monthly had the most favorable mean primary endpoint, increasing monthly KGE from 0.229 to 0.337, and the lowest monthly RMSE, decreasing it from 116.20 to 105.06 mm. NP monthly was close, with KGE 0.329 and RMSE 109.45 mm. Both monthly methods increased monthly-climatology correlation to at least 0.864 and produced a favorable point change for that metric in 59 of 60 paired evaluations.

The paired bootstrap did not support a broad parametric advantage over the empirical monthly method (Table 5). For P monthly minus NP monthly, the median combination-level KGE difference was 0.0002, with 30 combinations favoring each method. The median RMSE difference was -1.45 mm, with P monthly lower in 36 combinations and NP monthly lower in 24. Thus, the aggregate ranking is driven by heterogeneous station-model responses rather than uniform dominance.

Table 4 reports arithmetic means of metric levels, whereas Table 5 reports medians of pair-specific corrected-minus-raw changes. The two summaries therefore need not coincide when pairwise changes are skewed or contain influential values.

Upper-tail performance followed a different ordering (Figure 5). NP annual reduced mean absolute q99 bias from 52.98% to 24.79%, compared with 27.37% for NP monthly and 32.54% for P monthly. Pooling months provides a larger empirical wet-day sample and therefore more stable upper quantiles, whereas month-wise fitting directly resolves seasonality at the cost of smaller samples. No method was best on every endpoint.

Annual variants were weaker for monthly performance: NP annual increased mean RMSE to 124.74 mm and reduced mean KGE to 0.203, while P annual yielded 123.72 mm and 0.152. Distributional correction alone therefore does not guarantee better month-to-month agreement. This is consistent with the limitation that univariate mapping cannot reconstruct event sequencing absent from free-running climate simulations [13].

Station-aggregated responses were heterogeneous (Figure 6). Mean corrected-minus-raw KGE was positive at 9 of 12 stations for NP monthly and 11 of 12 for P monthly. Stations 500008 and 500009 showed the largest favorable KGE changes (approximately 0.29–0.31), whereas station 500002 decreased under both methods. The station-by-model display is descriptive: the small network, missing elevation metadata, and shared model cells do not support a causal geographic interpretation.

Table 4. Independent-validation means across paired model-station evaluations.

Metric	Raw	NP annual	NP monthly	P annual	P monthly	Favorable mean (descriptive)
Mean absolute PBIAS (%)	19.99	13.47	13.81	22.13	15.48	NP annual
Kolmogorov-Smirnov D	0.201	0.106	0.101	0.141	0.098	P monthly
Mean absolute q99 bias (%)	52.98	24.79	27.37	44.86	32.54	NP annual
Monthly RMSE (mm)	116.20	124.74	109.45	123.72	105.06	P monthly
Monthly MAE (mm)	78.15	81.06	70.47	80.46	67.40	P monthly
Monthly KGE	0.229	0.203	0.329	0.152	0.337	P monthly
Monthly correlation r	0.342	0.328	0.425	0.342	0.428	P monthly
Monthly climatology r	0.643	0.629	0.880	0.644	0.864	NP monthly

Note: PBIAS and q99 bias are summarized as absolute values. The final column is a direction-based descriptive ranking and does not imply statistical significance; the 60 pairs are not independent spatial replicates.

Table 5. Paired year-block bootstrap summary for corrected-minus-raw monthly metrics.

Method	Met.	Raw mean	Corrected mean	Median delta	Median 95% interval	Favorable point change	CI F/A/O
NP annual	RMSE	116.20	124.74	+8.75	[+3.33, +17.16]	22/60	19/32/9
NP annual	KGE	0.229	0.203	-0.048	[-0.126, +0.057]	25/60	14/20/26
NP monthly	RMSE	116.20	109.45	-5.57	[-24.68, +10.13]	38/60	24/5/31
NP monthly	KGE	0.229	0.329	+0.071	[-0.078, +0.264]	41/60	20/0/40
P annual	RMSE	116.20	123.72	+0.73	[-3.00, +4.67]	29/60	23/27/10
P annual	KGE	0.229	0.152	-0.006	[-0.118, +0.056]	27/60	16/19/25
P monthly	RMSE	116.20	105.06	-11.07	[-26.99, +2.54]	44/60	27/1/32
P monthly	KGE	0.229	0.337	+0.095	[-0.041, +0.261]	45/60	21/0/39

Note: Interval bounds are medians of 60 pair-specific bootstrap limits, not a confidence interval for a network-wide effect. For RMSE, negative delta is favorable; for KGE, positive delta is favorable. CI F/A/O denotes favorable, adverse, and overlapping-zero intervals; counts are descriptive because pairs are dependent.




Figure 4. Validation distributions for 60 model-station combinations per variant. Boxes show the interquartile range, whiskers show the 10th–90th percentiles, diamonds show means, center lines show medians, and dashed vertical lines show the raw mean in each panel; these are not bootstrap distributions.




Figure 5. Percentage of 60 paired model-station evaluations with a favorable point change relative to raw CMIP6 simulations. Values above 50% indicate a favorable change for most pairs; percentages are descriptive and pairs are not independent spatial replicates.




Figure 6. Station-level response of monthly QDM. (a) Heatmap of corrected-minus-raw KGE for NP monthly QDM by station and CMIP6 model. (b) Paired station means for non-parametric and parametric monthly QDM. Values are descriptive because gauges share model grid cells; station patterns are not interpreted as causal geographic effects.

3.3 Parametric selection, fallback, and tail sensitivity

Maximum-likelihood fitting changed the distribution-selection pattern relative to the earlier method-of-moments implementation. Pearson type III was selected for 55 of 60 observed annual fits and 610 of 620 eligible observed monthly fits. Model-side fits were more diverse: Pearson type III accounted for 496 monthly fits, generalized extreme value for 106, lognormal for 12, gamma for 3, and Weibull minimum for 3 (Table 6). All 620 eligible monthly blocks fitted successfully.

One hundred monthly blocks were ineligible because at least one calibration wet-day sample contained fewer than 30 values. These blocks clustered in dry-season months and used the annual fallback. This distinction is essential: the structural total remains 720, selection counts sum to the 620 eligible monthly fits, and fallback counts account for the remaining 100.

Tail safeguards affected the parametric ranking (Table 7). With the primary factor of 5.0, P monthly KGE was 0.337 versus 0.329 for NP monthly; without a cap, P monthly KGE decreased to 0.327. A stricter 1.5 factor capped 171 P-monthly validation values and improved mean RMSE to 101.91 mm, illustrating that aggressive truncation can improve central error metrics while imposing a strong assumption on extremes. The cap sensitivity therefore precludes a universal claim that parametric monthly QDM is superior.

The q99 metric changed little across the three parametric cap settings because most capped values were above the station-level 99th percentile. This does not make the cap inconsequential: a small number of altered maxima can materially affect return-level, IDF, or flood applications. Recent work similarly emphasizes that tail handling can behave differently inside and outside calibration and must be evaluated explicitly [8, 14].

AICc produced exactly the same selections and validation metrics as AIC: observed fits selected Pearson type III in 55 of 60 annual and 610 of 620 eligible monthly blocks, while model-side monthly counts remained 496 Pearson type III, 106 generalized extreme value, 12 lognormal, 3 gamma, and 3 Weibull. Because AICc reproduced the selections and validation metrics in this analysis, the selection pattern was not sensitive to the small-sample correction. This does not establish a physical rainfall law; Pearson type III retained greater shape and location flexibility than several constrained candidates. Because family-specific boundary and parameter-

shape diagnostics were not prespecified, the selection frequencies are descriptive and should not be interpreted as evidence that Pearson type III uniquely represents regional rainfall physics.

Wet-day threshold sensitivity was method-dependent (Figure 7). NP monthly remained stable across 0.1, 0.5, and 1.0 mm day⁻¹ (KGE 0.332, 0.330, and 0.329; RMSE 109.01, 109.26, and 109.45 mm). P monthly changed more strongly: KGE was 0.217, 0.327, and 0.337, while absolute q99 bias was 49.36%, 45.87%, and 32.54%. Thus, the 1.0-mm convention is defensible and gives the most favorable tested parametric result, but the parametric ranking is conditional on how trace rainfall is classified.

Table 6. Parametric fit accounting and numerical safeguards in validation.

Method	Eligible/structural	Observed selection	Model selection	Annual fallback	Fit failures	Capped	Clipped events
P annual	60/60	P3 55; LN 5	P3 45; GEV 15	0	0	3	247
P monthly	620/720	P3 610; GEV 5; LN 5	P3 496; GEV 106; LN 12; Gamma 3; Weibull 3	100	0	8	229

Note: P3 = Pearson type III; GEV = generalized extreme value; LN = lognormal. Monthly selection counts sum to 620 eligible blocks. The 100 ineligible blocks used annual calibration-only transfer functions. Capped values count validation outputs truncated above the selected cap. Clipped delta events are days on which the delta ratio was constrained at the lower or upper bound [0.2, 5.0]; event counts are not numbers of fitted blocks.

Table 7. Sensitivity of parametric validation metrics to the upper-tail cap factor.

Cap factor	P annual RMSE	P annual KGE	P monthly RMSE	P monthly KGE	P monthly q99 bias (%)	P annual capped	P monthly capped
1.5	122.01	0.167	101.91	0.355	32.60	35	171
5.0 (primary)	123.72	0.152	105.06	0.337	32.54	3	8
None	126.64	0.125	106.46	0.327	32.54	0	0

Note: The cap is factor times the maximum observed wet-day rainfall in the relevant calibration block and is a numerical safeguard rather than a return-level threshold. None denotes the uncapped sensitivity run. This cap-sensitivity experiment applies only to the parametric variants; non-parametric results are not included in Table 7.




Figure 7. Sensitivity of monthly QDM validation metrics to the observed wet-day threshold. Points and lines are means across 60 paired model-station evaluations; translucent ribbons show the interquartile range across those pairs, and the dashed line shows the raw mean. Lower is favorable for RMSE and absolute q99 bias, whereas higher is favorable for KGE. Lines aid comparison and do not imply continuous interpolation.

3.4 Implications and limitations

The model-specific raw results illuminate why a single correction function was insufficient. ACCESS-ESM1-5 and MIROC6 overestimated wet-day frequency by 0.227 and 0.232, respectively, and had positive mean PBIAS of 18.7% and 25.3%. EC-Earth3 instead had a negative mean PBIAS of -15.9% but the lowest raw monthly RMSE (92.30 mm) among the five models. Thus, similar aggregate errors arose from different combinations of event frequency, intensity, and seasonal phase. This heterogeneity agrees with Southeast Asian CMIP6 evaluations showing that rainfall skill reflects both monsoon circulation and model-specific parameterization, and that good large-scale circulation does not guarantee local rainfall skill [25, 26].

A physically plausible interpretation follows from the province's narrow coastal geometry. Moist southwesterly flow encounters the Tenasserim Range along the western boundary, while Gulf moisture and the northeast monsoon contribute to late-year rainfall on the eastern coastal plain [15]. The station-series files show that several gauges share identical native-grid extractions, so the effective spatial resolution remains coarse relative to the province's mountain-coast transition. The especially large wet-frequency excess in ACCESS-ESM1-5 and MIROC6 is consistent with overly frequent resolved or parameterized rainfall, whereas EC-Earth3's negative volume bias suggests a different compensation between frequency and intensity. Grid size alone cannot explain this ordering because model physics, circulation, and convection schemes also differ. These are process-informed interpretations, not causal attribution, because circulation and moisture-flux diagnostics were not analyzed.

The most stable result is the advantage of month-wise calibration for seasonal and monthly metrics. Annual pooling combines dry-season occurrence, southwest-monsoon rainfall, and the late-year coastal maximum in one transfer function. Month-wise QDM instead estimates separate wet-day thresholds and intensity distributions for each phase of the seasonal cycle, which better respects the changing atmospheric regime without claiming to simulate its dynamics. Both monthly methods consequently improved RMSE, KGE, monthly correlation, and climatology more consistently than annual calibration. The countervailing cost is a roughly twelve-fold reduction in sample size per transfer function, which explains the dry-season fallback and contributes to upper-tail uncertainty.

Method choice should follow application needs. NP monthly is an assumption-lean default for seasonal reservoir-inflow and monthly drought or water-balance screening; it should not be interpreted as a substitute for event-scale irrigation scheduling. NP annual is preferable when marginal q99 fidelity is the priority, although q99 is not a return-level estimator. P monthly can reduce monthly error further, but its advantage is conditional on distribution, wet-day threshold, and tail constraint. Fixed caps are unsuitable for design-storm, IDF, or flood-frequency estimation; such uses require uncapped sensitivity and dedicated extreme-value analysis. None of the variants restores observed daily event timing, spatial coherence, or wet-spell persistence.

Four limitations remain. The study covers one province, 12 gauges, five historical runs, and a 14-year validation window. Shared native grid cells reduce spatial replication, and the analysis-ready CMIP6 products are station-matched derivatives rather than original NetCDF files with per-file tracking identifiers. The metadata summary and checksum manifest identify the exact derivative files used; independent recreation would additionally require repeating the spatial extraction from the corresponding ESGF datasets. No future SSP experiment was analyzed, so preservation of projected quantile change was implemented by the QDM formulation but not empirically evaluated. Future work should compare QDM directly with ordinary quantile mapping and scaled distribution mapping under the same holdout, test moving-window seasonal fits and explicit extreme-value tails, analyse circulation and spatial dependence, and evaluate regional or dynamically downscaled ensembles across future scenarios.

4. Conclusions

Independent temporal validation indicates that month-wise QDM gives the most favorable descriptive pattern for monthly rainfall performance in Prachuap Khiri Khan. P monthly had the highest mean KGE and lowest mean RMSE under the five-times cap, but its KGE advantage over NP monthly was negligible in paired comparisons and reversed without the cap. AICc reproduced AIC selections, whereas lowering the wet-day threshold to 0.1 mm reduced P-monthly KGE to 0.217 while NP monthly remained stable. Seasonal

stratification, rather than a universally superior distribution family, is therefore the principal transferable finding.

NP annual gave the lowest q99 bias, whereas monthly methods best represented seasonal structure. For monthly water-balance applications, NP monthly is the conservative default; NP annual can support upper-quantile screening; and P monthly requires documented occurrence, fallback, clipping, and cap sensitivity. Fixed-cap outputs should not be used directly for design extremes. The 60 paired evaluations share grid cells and model lineages and are not independent spatial replicates, so network-wide inferential superiority is not claimed. The fit-freeze-apply protocol and explicit sensitivities provide a stronger basis for further regional and future-scenario evaluation than same-period fitting alone.

Acknowledgements: The author acknowledges the World Climate Research Programme (WCRP), which coordinated and promoted CMIP6 through its Working Group on Coupled Modelling; the participating climate modelling groups for producing and making their model output available; and the Earth System Grid Federation (ESGF) for archiving and providing access to CMIP6 data. The author also acknowledges the responsible Thai data custodians who provided access to the rainfall records used in this study.

Author Contributions: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Data curation, Visualization, Writing – original draft, and Writing – review and editing were performed by the author. The author has read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: The rainfall observations used in this study were supplied through the Hydro-Informatics Institute/Royal Irrigation Department project archive and remain subject to the terms of the responsible Thai data custodians. Access requests should be directed to those custodians. CMIP6 datasets are discoverable through ESGF using the source, experiment, member, grid, variable, frequency, and date-range metadata summarized in Table 2. The complete reproducibility package, including the analysis code, input-file checksum manifest, derived metrics, diagnostics, sensitivity outputs, and figure-generation scripts, is available from the corresponding author upon reasonable request.

Conflicts of Interest: The author declares no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Eyring, V.; Bony, S.; Meehl, G.A.; Senior, C.A.; Stevens, B.; Stouffer, R.J.; Taylor, K.E. Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) Experimental Design and Organization. *Geosci. Model Dev.* 2016, 9, 1937–1958. <https://doi.org/10.5194/gmd-9-1937-2016>.
2. Tebaldi, C.; Knutti, R. The Use of the Multi-Model Ensemble in Probabilistic Climate Projections. *Philos. Trans. R. Soc. A* 2007, 365, 2053–2075. <https://doi.org/10.1098/rsta.2007.2076>.
3. Maraun, D. Bias Correction, Quantile Mapping, and Downscaling: Revisiting the Inflation Issue. *J. Clim.* 2013, 26, 2137–2143. <https://doi.org/10.1175/JCLI-D-12-00821.1>.
4. Themeßl, M.J.; Gobiet, A.; Leuprecht, A. Empirical-Statistical Downscaling and Error Correction of Daily Precipitation from Regional Climate Models. *Int. J. Climatol.* 2011, 31, 1530–1544. <https://doi.org/10.1002/joc.2168>.
5. Gudmundsson, L.; Bremnes, J.B.; Haugen, J.E.; Engen-Skaugen, T. Technical Note: Downscaling RCM Precipitation to the Station Scale Using Statistical Transformations. *Hydrol. Earth Syst. Sci.* 2012, 16, 3383–3390. <https://doi.org/10.5194/hess-16-3383-2012>.
6. Piani, C.; Haerter, J.O.; Coppola, E. Statistical Bias Correction for Daily Precipitation in Regional Climate Models over Europe. *Theor. Appl. Climatol.* 2010, 99, 187–192. <https://doi.org/10.1007/s00704-009-0134-9>.
7. Cannon, A.J.; Sobie, S.R.; Murdock, T.Q. Bias Correction of GCM Precipitation by Quantile Mapping: How Well Do Methods Preserve Changes in Quantiles and Extremes? *J. Clim.* 2015, 28, 6938–6959. <https://doi.org/10.1175/JCLI-D-14-00754.1>.
8. Padulano, R.; Gomez-Mogollon, L.A.; Napolitano, L.; Rianna, G. Quantile-Based Bias-Correction of Extreme Rainfall: Pros and Cons of Popular Methods for Climate Signal Preservation. *J. Hydrol.* 2025, 653, 132814. <https://doi.org/10.1016/j.jhydrol.2025.132814>.

9. Casanueva, A.; Herrera, S.; Iturbide, M.; Lange, S.; Jury, M.; Dosio, A.; Maraun, D.; Gutiérrez, J.M. Testing Bias Adjustment Methods for Regional Climate Change Applications under Observational Uncertainty and Resolution Mismatch. *Atmos. Sci. Lett.* 2020, 21, e978. <https://doi.org/10.1002/asl.978>.
10. Spuler, F.R.; Wessel, J.B.; Comyn-Platt, E.; Varndell, J.; Cagnazzo, C. ibicus: A New Open-Source Python Package and Comprehensive Interface for Statistical Bias Adjustment and Evaluation in Climate Modelling (v1.0.1). *Geosci. Model Dev.* 2024, 17, 1249–1269. <https://doi.org/10.5194/gmd-17-1249-2024>.
11. Song, Y.H.; Chung, E.-S. Intercomparison of Bias Correction Methods for Precipitation of Multiple GCMs across Six Continents. *Geosci. Model Dev.* 2025, 18, 8017–8045. <https://doi.org/10.5194/gmd-18-8017-2025>.
12. Gergel, D.R.; Malevich, S.B.; McCusker, K.E.; Tenezakis, E.; Delgado, M.T.; Fish, M.A.; Kopp, R.E. Global Downscaled Projections for Climate Impacts Research: Preserving Quantile Trends for Modeling Future Climate Impacts. *Geosci. Model Dev.* 2024, 17, 191–227. <https://doi.org/10.5194/gmd-17-191-2024>.
13. Potter, N.J.; Chiew, F.H.S.; Charles, S.P.; Fu, G.; Zheng, H.; Zhang, L. Bias in Dynamically Downscaled Rainfall Characteristics for Hydroclimatic Projections. *Hydrol. Earth Syst. Sci.* 2020, 24, 2963–2979. <https://doi.org/10.5194/hess-24-2963-2020>.
14. Berg, P.; Bosshard, T.; Bozhinova, D.; Bärting, L.; Löw, J.; Nilsson, C.; Strandberg, G.; Södling, J.; Thuresson, J.; Wilcke, R.; et al. Robust Handling of Extremes in Quantile Mapping - Murder Your Darlings. *Geosci. Model Dev.* 2024, 17, 8173–8179. <https://doi.org/10.5194/gmd-17-8173-2024>.
15. Pannak, M.; Sittichok, K.; Theprasit, C.; Chompuchan, C. Spatial Exceedance Probability Mapping of Monthly Rainfall Using Gridded Precipitation Products in an Orographically Complex Monsoon Basin, Western Thailand. *Hydrology* 2026, 13, 155. <https://doi.org/10.3390/hydrology13060155>.
16. Pettitt, A.N. A Non-Parametric Approach to the Change-Point Problem. *Appl. Stat.* 1979, 28, 126–135. <https://doi.org/10.2307/2346729>.
17. Swart, N.C.; Cole, J.N.S.; Kharin, V.V.; Lazare, M.; Scinocca, J.F.; Gillett, N.P.; Anstey, J.; Arora, V.; Christian, J.R.; Jiao, Y.; et al. The Canadian Earth System Model Version 5 (CanESM5.0.3). *Geosci. Model Dev.* 2019, 12, 4823–4873. <https://doi.org/10.5194/gmd-12-4823-2019>.
18. Danabasoglu, G.; Lamarque, J.-F.; Bacmeister, J.; Bailey, D.A.; DuVivier, A.K.; Edwards, J.; Emmons, L.K.; Fasullo, J.; Garcia, R.; Gettelman, A.; et al. The Community Earth System Model Version 2 (CESM2). *J. Adv. Model Earth Syst.* 2020, 12, e2019MS001916. <https://doi.org/10.1029/2019MS001916>.
19. Döscher, R.; Acosta, M.; Alessandri, A.; Anthoni, P.; Arneth, A.; Arsouze, T.; Bergmann, T.; Bernadello, R.; Boussetta, S.; Caron, L.P.; et al. The EC-Earth3 Earth System Model for the Coupled Model Intercomparison Project 6. *Geosci. Model Dev.* 2022, 15, 2973–3020. <https://doi.org/10.5194/gmd-15-2973-2022>.
20. Tatebe, H.; Ogura, T.; Nitta, T.; Komuro, Y.; Ogochi, K.; Takemura, T.; Sudo, K.; Sekiguchi, M.; Abe, M.; Saito, F.; et al. Description and Basic Evaluation of Simulated Mean State, Internal Variability, and Climate Sensitivity in MIROC6. *Geosci. Model Dev.* 2019, 12, 2727–2765. <https://doi.org/10.5194/gmd-12-2727-2019>.
21. Ziehn, T.; Chamberlain, M.A.; Law, R.M.; Lenton, A.; Bodman, R.W.; Dix, M.; Stevens, L.; Wang, Y.-P.; Srbinovsky, J. The Australian Earth System Model: ACCESS-ESM1.5. *J. South. Hemisph. Earth Syst. Sci.* 2020, 70, 193–214. <https://doi.org/10.1071/ES19035>.
22. Zhang, X.; Alexander, L.; Hegerl, G.C.; Jones, P.D.; Tank, A.K.; Peterson, T.C.; Trewin, B.; Zwiers, F.W. Indices for Monitoring Changes in Extremes Based on Daily Temperature and Precipitation Data. *Wiley Interdiscip. Rev. Clim. Change* 2011, 2, 851–870. <https://doi.org/10.1002/wcc.147>.
23. Virtanen, P.; Gommers, R.; Oliphant, T.E.; Haberland, M.; Reddy, T.; Cournapeau, D.; Burovski, E.; Peterson, P.; Weckesser, W.; Bright, J.; et al. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nat. Methods* 2020, 17, 261–272. <https://doi.org/10.1038/s41592-019-0686-2>.
24. Gupta, H.V.; Kling, H.; Yilmaz, K.K.; Martinez, G.F. Decomposition of the Mean Squared Error and NSE Performance Criteria: Implications for Improving Hydrological Modelling. *J. Hydrol.* 2009, 377, 80–91. <https://doi.org/10.1016/j.jhydrol.2009.08.003>.
25. Khadka, D.; Babel, M.S.; Abatan, A.A.; Collins, M. An Evaluation of CMIP5 and CMIP6 Climate Models in Simulating Summer Rainfall in the Southeast Asian Monsoon Domain. *Int. J. Climatol.* 2022, 42, 1181–1202. <https://doi.org/10.1002/joc.7296>.
26. Liu, S.; Raghavan, S.V.; Ona, B.J.; Nguyen, N.S. Bias Evaluation in Rainfall over Southeast Asia in CMIP6 Models. *J. Hydrol.* 2023, 621, 129593. <https://doi.org/10.1016/j.jhydrol.2023.129593>.