

ENSO-Conditioned Seasonal Rainfall Extremes and Signal Preservation after Cross-Fitted Quantile Delta Mapping over Uttaradit, Thailand

Surasit Punyawansiri^{1,*}

¹Office of Water Management and Hydrology, Royal Irrigation Department, Dusit, Bangkok 10300, Thailand

*Corresponding author: Surasit.ku@ku.th

Abstract

Bias adjustment is commonly assumed to retain physically meaningful climate variability, but this assumption is rarely tested for El Niño–Southern Oscillation (ENSO) signals at the daily station scale. We quantified ENSO-conditioned rainfall over Uttaradit, northern Thailand, and tested whether blocked cross-fitted quantile delta mapping (QDM) preserved the historical ENSO response of seven CMIP6 models. Daily rainfall at 13 gauges and exact-member model simulations were analysed for 1981–2014. Observed seasons were classified from a frozen NOAA CPC ERSSTv6 Oceanic Niño Index; every model was classified from its own detrended Niño-3.4 sea-surface-temperature anomaly. Persistent episodes required at least five overlapping three-month windows, and seasons were assigned only by strict majority. Four pre-specified endpoints were seasonal precipitation total (PRCPTOT), wet-day frequency, maximum one-day rainfall (Rx1day), and maximum consecutive dry days (CDD). Inference resampled management-season climate years rather than days or stations. Observations suggested drier El Niño conditions, including rainy-season PRCPTOT -7.2% (95% CI -18.8 to 4.4), rainy-season Rx1day -24.3% (-40.1 to 3.0), and hot/dry-season PRCPTOT -28.7% (-55.5 to 6.1), while La Niña increased hot/dry wet-day frequency by 33.0% (0.0 to 53.1). None of the 16 observed phase–metric tests survived Benjamini–Hochberg adjustment (minimum $q=0.433$), and no neutral-centred asymmetry test was supported. Across model-ensemble responses, QDM preserved only 2 of 16 signals, attenuated 4, amplified 3, reversed 1, and left 6 indeterminate because the raw response was near zero. The hot/dry El Niño PRCPTOT response reversed from -10.4% raw to $+14.4\%$ after QDM, compared with -28.7% observed. Thus, marginal bias correction did not guarantee preservation of ENSO-conditioned rainfall occurrence, extremes, or persistence. Event-level uncertainty and explicit signal-preservation diagnostics should accompany bias-adjusted hydroclimate applications.

Keywords: bias adjustment; CMIP6; El Niño–Southern Oscillation; rainfall extremes; quantile delta mapping; Thailand

1. Introduction

El Niño–Southern Oscillation (ENSO) reorganises tropical circulation and affects seasonal rainfall across Southeast Asia. Thailand commonly experiences rainfall deficits during El Niño and wetter conditions during La

32 Niña, although the magnitude and timing depend on region and season [1–4]. Recent national analyses based
33 on gridded reanalysis confirm this broad contrast, but local water management depends on daily station
34 rainfall, dry-spell persistence, and short-duration extremes that monthly national averages cannot resolve [4].
35 Northern Thailand is especially sensitive because water supply, flood control, and rain-fed agriculture share a
36 strongly seasonal monsoon resource.

37 Global climate models provide physically consistent ENSO and rainfall simulations, yet their coarse grids and
38 precipitation biases limit direct station-scale use [5]. Quantile mapping is therefore widely used to align
39 simulated and observed marginal distributions [6,7]. Quantile delta mapping (QDM) was designed to preserve
40 simulated changes in quantiles while correcting historical bias [6]. That property, however, concerns
41 distributional change between periods; it does not ensure that a circulation-conditioned contrast such as El
42 Niño minus Neutral will survive adjustment. The risk is greater for wet-day occurrence and sequence-dependent
43 metrics because a univariate mapping changes magnitudes but does not reconstruct the model's temporal
44 ordering [7,8]. In-sample calibration can further conceal instability.

45 Three gaps motivate this study. First, observational ENSO analyses in Thailand have generally used national or
46 regional monthly data rather than a dense daily gauge network [1–4]. Second, model evaluation often assigns
47 observed ENSO years to all simulations. Free-running historical models do not reproduce the observed
48 chronology, so their rainfall responses must be conditioned on ENSO diagnosed from each model's own sea-
49 surface temperatures. Third, few bias-adjustment studies evaluate whether an ENSO-conditioned response is
50 preserved out of sample, and fewer still respect the event-level sampling unit.

51 We address these gaps for Uttaradit Province using 13 rain gauges, seven CMIP6 models, and a fully blocked
52 cross-fitted QDM experiment over 1981–2014. The objectives were to: (i) estimate observed El Niño and La
53 Niña responses relative to Neutral for four management-relevant rainfall endpoints; (ii) compare raw and QDM-
54 adjusted model-ensemble responses using model-specific ENSO classifications; and (iii) distinguish the
55 directional phase contrast from true neutral-centred asymmetry. The central hypothesis was not that QDM
56 would reproduce the observed ENSO response, but that its preservation behaviour could be measured
57 transparently.

58 2. Materials and Methods

59 2.1 Study area and data

60 Uttaradit lies in monsoon-dominated northern Thailand. Thirteen daily rainfall gauges (station codes 351001–
61 351012 and 351201) span 17.23–18.02°N, 100.05–101.07°E and 54.6–427.9 m above mean sea level. The
62 analysis period was 1 January 1981–31 December 2014. The rainy management season comprised May–
63 October and was labelled by calendar year. The hot/dry season comprised November–April and was labelled by
64 its November start year; consequently, complete seasons extended from 1981/82 through 2013/14.

65 Daily historical precipitation was used from seven CMIP6 models: ACCESS-ESM1-5, CanESM5, CESM2, EC-Earth3,
66 FGOALS-g3, MIROC6, and MRI-ESM2-0 [5,9–15]. One ensemble member was retained per model and matched
67 exactly between precipitation and sea-surface-temperature data. Gauge extraction can duplicate a coarse grid
68 cell; therefore, station series sharing an identical raw precipitation time series were collapsed to a unique grid
69 signature before model aggregation. Models contributed one value each to the ensemble regardless of native
70 resolution.

71 Observed ENSO status came from the NOAA Climate Prediction Center historical Oceanic Niño Index (ONI) table
72 based on ERSSTv6, frozen on 1 September 2026 [16]. For each GCM, monthly tos from the exact historical
73 member was area-weighted over Niño-3.4 (5°S–5°N, 170°W–120°W). Anomalies used the model's 1981–2010
74 monthly climatology, a centred linear detrend, and a centred three-month running mean.

75 2.2 Quality control and common calendar

76 Observed zero rainfall was screened on the native Gregorian calendar before leap-day removal. A zero month
77 was flagged as probable missing only when at least two pre-specified indicators supported the diagnosis,
78 including neighbour disagreement, an anomalously long zero run, or an annual rainfall deficit. Only probable-
79 missing values were set to missing; confirmed dry and uncertain zeros were retained. This classified 43 station-
80 months as probable missing and removed 1,308 daily values. All series were then placed on a common 365-day
81 calendar by dropping 29 February without inserting synthetic days. A station-season entered a metric only if its
82 expected daily coverage was complete after quality control.

83 2.3 ENSO episodes and management-season classification

84 El Niño and La Niña episodes required ONI or model Niño-3.4 anomalies of at least +0.5°C or at most –0.5°C,
85 respectively, for five or more overlapping three-month windows [16]. For each management season, all
86 overlapping persistent windows were counted. A season was El Niño or La Niña only when that phase had a

87 strict majority; Neutral required no persistent episode. Ties, mixed-phase seasons, and seasons lacking a strict
88 majority were labelled transition/unclassified and excluded from phase contrasts. This same algorithm was
89 applied to observations and each model, while the underlying index remained source-specific.

90 2.4 Blocked cross-fitted QDM

91 Multiplicative QDM with wet-day frequency adaptation was fitted separately for each station/grid signature and
92 calendar month [6]. Seven climate-year blocks were held out in turn: 1981–1985, 1986–1990, 1991–1995,
93 1996–2000, 2001–2005, 2006–2010, and 2011–2014. Observations from the target block were never used in its
94 mapping. Wet-day quantiles used the ≥ 1 mm threshold, Hazen mid-ranks, linear interpolation, and constant-
95 ratio extrapolation. At least 30 wet days were required for calibration. If a single month was sparse, only
96 calibration samples were pooled from that month and its two adjacent months; the corrected target remained
97 in its original month. This pre-specified fallback was used in 574 of 7,644 fits (7.5%).

98 2.5 Rainfall endpoints and hierarchy

99 Four primary endpoints were fixed before analysis: PRCPTOT, the total from days with rainfall ≥ 1 mm; wet-day
100 frequency, the percentage of season days ≥ 1 mm; Rx1day, the seasonal maximum daily rainfall; and CDD, the
101 longest run of days below 1 mm [17]. Secondary indices were retained only in the supplementary workbook.
102 Aggregation followed the data-generating hierarchy: daily values $\rightarrow$ management-season climate year $\rightarrow$ station
103 $\rightarrow$ unique raw-grid signature $\rightarrow$ model $\rightarrow$ ensemble. Within each source, the phase response was

$$104 R_p = 100(M_p - M_N) / M_N,$$

105 where (M_p) and (M_N) are phase and Neutral means within a station or unique grid signature, followed by
106 regional and ensemble medians. At least three phase seasons and three Neutral seasons were required. The
107 observed analysis used the median station response. Model responses first collapsed duplicated grid signatures
108 and then used the median across models.

109 2.6 Inference, asymmetry, and preservation

110 Uncertainty was estimated with 5,000 bootstrap repetitions and 95% percentile intervals. Permutation p-values
111 used 4,999 label permutations. Resampling and permutation occurred at the management-season climate-year
112 level; model ensembles resampled events independently within model. The 16 phase-by-season-by-metric tests
113 were adjusted by the Benjamini–Hochberg procedure. Effect sizes and intervals are emphasised because phase
114 samples were small.

Two quantities were kept separate. The phase contrast, $\backslash(A_{\{La\}}-A_{\{El\}}\backslash)$, describes the difference between La Niña and El Niño responses. True neutral-centred asymmetry, $\backslash(A_{\{La\}}+A_{\{El\}}\backslash)$, tests whether equal and opposite responses cancel around Neutral. A large phase contrast alone is not evidence of asymmetry.

QDM signal preservation was classified from paired-model ensemble responses. A raw response with magnitude <5 percentage points was indeterminate. Otherwise, a sign reversal was reversed; a magnitude ratio $|QDM/raw|$ of 0.8–1.2 was preserved, <0.8 was attenuated, and >1.2 was amplified. Directional agreement with observations and paired-model sign retention were reported separately.

All processing was scripted in Python. Acceptance gates verified frozen ENSO provenance, calendar completeness, source-specific classification, seven-model/13-gauge coverage, absence of target-fold leakage, paired phase responses, separation of contrast and asymmetry, and event-level inference.

3. Results and Discussion

3.1 ENSO samples and observed response

The observed rainy-season sample comprised 6 El Niño, 8 Neutral, 7 La Niña, and 13 transition/unclassified seasons; the corresponding hot/dry counts were 8, 8, 12, and 5 (Figure 1; Table 1). Model-specific classification produced appreciably different sample counts, as expected for free-running simulations. One ACCESS-ESM1-5 rainy-season La Niña response failed the minimum sample gate, leaving six paired models for that contrast; other ensemble contrasts retained seven models.

Observed point estimates were broadly consistent with rainfall suppression during El Niño but remained uncertain (Figure 2; Table 2). Rainy-season El Niño responses were -7.2% for PRCPTOT and -24.3% for Rx1day, while CDD increased by 17.9% . Hot/dry El Niño PRCPTOT was -28.7% and Rx1day was -27.2% . La Niña increased hot/dry wet-day frequency by 33.0% , whose unadjusted permutation p-value was 0.0278, but the multiple-testing adjusted q-value was 0.433. All other observed intervals also included zero, and none of the 16 observed primary tests survived adjustment (minimum $q=0.433$). These results align directionally with national studies [1–4] but show that local daily responses are less certain than a simple wet-La Niña/dry-El Niño narrative suggests.

The wide intervals are not a computational defect: after persistent-episode classification and transition exclusion, only 6–12 observed event seasons were available per phase. Treating stations or daily values as independent replicates would have generated artificially narrow intervals. The event-level design therefore gives a more conservative and hydrologically interpretable uncertainty statement.

3.2 Raw versus cross-fitted QDM ENSO signals

Bias adjustment substantially altered several ENSO responses (Figure 3; Table 2). Of 16 model-ensemble phase-metric signals, only rainy PRCPTOT under El Niño and rainy CDD under El Niño met the preservation criterion. Four signals were attenuated, three amplified, one reversed, and six were indeterminate because the raw response was within ± 5 percentage points.

The clearest failure occurred for hot/dry El Niño PRCPTOT: the raw ensemble response of -10.4% agreed in sign with the observed -28.7% , whereas QDM produced $+14.4\%$. Model directional agreement with observations fell from 0.86 to 0.43. For rainy El Niño Rx1day, the raw -12.7% response was attenuated to -5.7% , both smaller than the observed -24.3% . Under rainy La Niña, both raw and QDM ensemble responses for PRCPTOT were negative (-10.0% and -15.0%), opposite to the observed $+2.7\%$; only one of six contributing models agreed with the observed direction.

These changes are compatible with the mechanics of univariate QDM. The mapping is fitted to calendar-month marginal distributions without conditioning on ENSO. Cross-fitting prevents the held-out observed events from directly shaping their correction, but does not impose a physical relationship between Niño-3.4 variability and local rainfall. Frequency adaptation and month-specific quantiles can also change event composites when ENSO phases sample different parts of the precipitation distribution. Sequence-dependent CDD is especially vulnerable because rank-based correction does not reorder wet and dry days [7,8].

3.3 Phase contrast and neutral-centred asymmetry

Observed La Niña-minus-El Niño phase contrasts were sizeable for some hot/dry metrics: $+46.0$ percentage points for PRCPTOT and $+43.8$ points for wet-day frequency (Figure 4). These contrasts indicate separation between phase composites, but their neutral-centred sums were -15.1 and $+15.5$ points, respectively. Across all eight observed season-metric combinations, true asymmetry q-values were ≥ 0.781 . The data therefore did not support a claim that El Niño and La Niña responses were unequal around Neutral.

QDM could change the asymmetry quantity even when paired-model signs appeared stable. For hot/dry PRCPTOT, neutral-centred asymmetry shifted from -8.3% raw to $+52.7\%$ after adjustment. This result should not be interpreted as evidence that QDM created a physical nonlinear ENSO response; rather, it demonstrates that a marginal correction can materially modify a conditioned composite. The distinction between phase contrast and asymmetry avoids a common interpretive error.

172 3.4 Implications, strengths, and limitations

173 The practical implication is that bias-adjusted rainfall should not automatically be treated as reliable for ENSO-
174 conditioned planning. A correction may improve the unconditional distribution yet distort the sign or magnitude
175 of the seasonal teleconnection used for reservoir operation or drought preparedness. A minimum workflow
176 should therefore report raw, corrected, and observed phase responses together; preserve the model-specific
177 ENSO chronology; and test occurrence, intensity, and persistence endpoints separately.

178 The study's main strengths are source-specific ENSO diagnosis, blocked cross-fitting, explicit raw-grid
179 deduplication, a fixed hierarchy, event-level resampling, and a reproducible acceptance audit. The analysis also
180 avoids inferring asymmetry from a phase difference alone. Limitations remain. Thirty-four years yield few
181 persistent events, so the study is powered for effect-size estimation rather than definitive detection. Each GCM
182 contributes one historical member; internal variability is therefore not separated from structural model
183 differences. The seven-model ensemble is not an independent random sample of all plausible models. The
184 gauge network is confined to one province, and QDM is univariate and precipitation-only. Finally, the observed
185 zero-screening rules reduce obvious missing-data contamination but cannot recover unrecorded rainfall.
186 Future work should add initial-condition ensembles, circulation diagnostics, and multivariate or weather-state-
187 conditioned corrections. Pre-registered comparisons with occurrence models or stochastic weather generators
188 would clarify whether sequence-dependent ENSO signals can be improved without sacrificing marginal
189 performance. Longer records could also distinguish season timing and ENSO intensity while retaining event-
190 level inference.

191 4. Conclusions

192 Daily observations over Uttaradit showed directionally drier El Niño seasons and a wetter hot/dry La Niña wet-
193 day response, but all primary phase effects were uncertain after multiplicity adjustment. Blocked cross-fitted
194 QDM did not reliably preserve model ENSO signals: only 2 of 16 ensemble responses were preserved, while one
195 reversed and most others were attenuated, amplified, or indeterminate. No neutral-centred asymmetry was
196 supported. These findings do not imply that QDM is generally unsuitable; they show that marginal distributional
197 correction and teleconnection preservation are distinct validation targets. ENSO-sensitive applications should
198 retain source-specific event chronology, use out-of-sample correction, resample at the event level, and audit
199 conditioned signals before operational interpretation.

200 Acknowledgements
 201 The author thanks the Royal Irrigation Department and the Thai Meteorological Department for access to the
 202 rain-gauge records, the World Climate Research Programme for coordinating CMIP6, and the modelling groups
 203 for producing and making their output available.

204 Author Contributions
 205 S. Punyawansiri: Conceptualization, methodology, software, formal analysis, data curation, validation,
 206 visualization, writing—original draft, and writing—review and editing.

207 Data and Code Availability
 208 The frozen analysis package contains configuration, source manifests with checksums, tests, the executable
 209 notebook, numerical outputs, figures, and supplementary tables and is available from the corresponding author
 210 upon reasonable request. Redistribution of station observations remains subject to the data providers' terms.
 211 CMIP6 model data are available through the Earth System Grid Federation; the NOAA CPC ONI source and
 212 capture metadata are recorded in the package.

213 Ethical Approval
 214 Not applicable. The study used meteorological observations and publicly archived climate-model output and
 215 involved no human participants or animals.

216 Declaration of Generative AI and AI-Assisted Technologies
 217 During preparation, the author used a generative AI assistant for code review, statistical verification, figure and
 218 document generation, and language editing. The author reviewed the analyses and text and takes responsibility
 219 for the submitted content.

220 Conflict of Interest Statement
 221 The author declares that there is no conflict of interest.

222 References
 223 [1] Singhrattana N., Rajagopalan B., Kumar K.K. and Clark M., Interannual and interdecadal variability of Thailand
 224 summer monsoon season. *J. Climate*, 2005; **18**: 1697–1708. DOI 10.1175/JCLI3364.1.
 225 [2] Limsakul A. and Singhruck P., Long-term trends and variability of total and extreme precipitation in Thailand.
 226 *Atmos. Res.*, 2016; **169**: 301–317. DOI 10.1016/j.atmosres.2015.10.015.
 227 [3] Räsänen T.A., Lindgren V., Guillaume J.H.A., Buckley B.M. and Kummu M., On the spatial and temporal
 228 variability of ENSO precipitation and drought teleconnection in mainland Southeast Asia. *Clim. Past*, 2016; **12**:
 229 1889–1905. DOI 10.5194/cp-12-1889-2016.

230 [4] Chanalert W., Wongsai N., Wongsai S. and Komontree P., Seasonal rainfall variability in Thailand: Long-term
231 trends and regional patterns from ERA5 data. *Chiang Mai J. Sci.*, 2026; **53**(4): e2026065. DOI
232 10.12982/CMJS.2026.065.

233 [5] Eyring V., Bony S., Meehl G.A., Senior C.A., Stevens B., Stouffer R.J. and Taylor K.E., Overview of the Coupled
234 Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geosci. Model Dev.*,
235 2016; **9**: 1937–1958. DOI 10.5194/gmd-9-1937-2016.

236 [6] Cannon A.J., Sobie S.R. and Murdock T.Q., Bias correction of GCM precipitation by quantile mapping: How
237 well do methods preserve changes in quantiles and extremes? *J. Climate*, 2015; **28**: 6938–6959. DOI
238 10.1175/JCLI-D-14-00754.1.

239 [7] Maraun D., Bias correcting climate change simulations: A critical review. *Curr. Clim. Change Rep.*, 2016; **2**:
240 211–220. DOI 10.1007/s40641-016-0050-x.

241 [8] Addor N., Rohrer M., Furrer R. and Seibert J., Propagation of biases in climate models from the synoptic to
242 the regional scale: Implications for bias adjustment. *J. Geophys. Res. Atmos.*, 2016; **121**: 2075–2089. DOI
243 10.1002/2015JD024040.

244 [9] Ziehn T., Chamberlain M.A., Law R.M., Lenton A., Bodman R.W., Dix M., et al., The Australian Earth System
245 Model: ACCESS-ESM1.5. *J. South. Hemisph. Earth Syst. Sci.*, 2020; **70**: 193–214. DOI 10.1071/ES19035.

246 [10] Swart N.C., Cole J.N.S., Kharin V.V., Lazare M., Scinocca J.F., Gillett N.P., et al., The Canadian Earth System
247 Model version 5 (CanESM5.0.3). *Geosci. Model Dev.*, 2019; **12**: 4823–4873. DOI 10.5194/gmd-12-4823-2019.

248 [11] Danabasoglu G., Lamarque J.-F., Bacmeister J., Bailey D.A., DuVivier A.K., Edwards J., et al., The Community
249 Earth System Model version 2 (CESM2). *J. Adv. Model Earth Syst.*, 2020; **12**: e2019MS001916. DOI
250 10.1029/2019MS001916.

251 [12] Döscher R., Acosta M., Alessandri A., Anthoni P., Arneth A., Arsouze T., et al., The EC-Earth3 Earth system
252 model for the Coupled Model Intercomparison Project 6. *Geosci. Model Dev.*, 2022; **15**: 2973–3020. DOI
253 10.5194/gmd-15-2973-2022.

254 [13] Li L., Yu Y., Tang Y., Lin P., Xie J., Song M., et al., The Flexible Global Ocean–Atmosphere–Land System
255 Model grid-point version 3 (FGOALS-g3): Description and evaluation. *J. Adv. Model Earth Syst.*, 2020; **12**:
256 e2019MS002012. DOI 10.1029/2019MS002012.

257 [14] Tatebe H., Ogura T., Nitta T., Komuro Y., Ogochi K., Takemura T., et al., Description and basic evaluation of
 258 simulated mean state, internal variability, and climate sensitivity in MIROC6. *Geosci. Model Dev.*, 2019; **12**:
 259 2727–2765. DOI 10.5194/gmd-12-2727-2019.

260 [15] Yukimoto S., Kawai H., Koshiro T., Oshima N., Yoshida K., Urakawa S., et al., The Meteorological Research
 261 Institute Earth System Model version 2.0, MRI-ESM2.0. *J. Meteorol. Soc. Jpn.*, 2019; **97**: 931–965. DOI
 262 10.2151/jmsj.2019-051.

263 [16] National Oceanic and Atmospheric Administration Climate Prediction Center, Cold & Warm Episodes by
 264 Season: Historical Oceanic Niño Index, ERSSTv6; Available at:
 265 https://www.cpc.ncep.noaa.gov/products/analysis_monitoring/enso/oni/v6/ (accessed 1 September 2026).

266 [17] Zhang X., Alexander L., Hegerl G.C., Jones P., Tank A.K., Peterson T.C., Trewin B. and Zwiers F.W., Indices for
 267 monitoring changes in extremes based on daily temperature and precipitation data. *WIREs Clim. Change*, 2011;
 268 **2**: 851–870. DOI 10.1002/wcc.147.

269

270 **Tables**
271

Table 1. ENSO management-season sample sizes. Model ranges use each GCM's exact-member Niño-3.4 classification.

Season	Complete seasons	Observed El Niño	Model range	Observed Neutral	Model range	Observed La Niña	Model range	Transition/unclassified
Rainy (May–Oct)	34	6	4–12	8	3–13	7	2–12	13
Hot/dry (Nov–Apr)	33	8	6–14	8	4–12	12	5–15	5

272
273

274
275

Table 2. Observed, raw-model, and cross-fitted-QDM ensemble response relative to Neutral (%; 95% bootstrap CI). BH q is shown for the observed response.

Season	Metric	Phase	Observed response [95% CI]	BH q	Raw ensemble	QDM ensemble
Rainy	PRCPTOT	El Niño	-7.2 [-18.8, 4.4]	0.471	-12.2 [-23.8, 1.0]	-14.5 [-25.1, -1.4]
Rainy	PRCPTOT	La Niña	2.7 [-10.2, 17.3]	0.790	-10.0 [-20.0, 3.0]	-15.0 [-26.3, -1.1]
Rainy	Wet-day frequency	El Niño	-0.1 [-12.3, 2.8]	0.980	-4.4 [-10.8, 3.2]	-7.0 [-12.4, -1.7]
Rainy	Wet-day frequency	La Niña	-2.4 [-9.6, 6.3]	0.647	-6.8 [-11.8, 0.3]	-4.7 [-8.2, -0.8]
Rainy	Rx1day	El Niño	-24.3 [-40.1, 3.0]	0.433	-12.7 [-32.3, 2.7]	-5.7 [-35.4, 9.7]
Rainy	Rx1day	La Niña	0.5 [-26.8, 35.8]	0.980	-12.9 [-28.2, 4.6]	-28.3 [-41.9, 0.5]
Rainy	CDD	El Niño	17.9 [-6.7, 41.0]	0.433	13.1 [-2.0, 41.0]	11.9 [-4.1, 25.3]
Rainy	CDD	La Niña	-5.8 [-19.0, 14.3]	0.711	16.2 [4.0, 45.6]	3.4 [-5.1, 21.9]
Hot/dry	PRCPTOT	El Niño	-28.7 [-55.5, 6.1]	0.433	-10.4 [-27.8, 7.2]	14.4 [-28.6, 42.9]
Hot/dry	PRCPTOT	La Niña	13.3 [-15.6, 58.0]	0.709	-0.4 [-13.9, 29.4]	20.3 [-16.6, 68.7]
Hot/dry	Wet-day frequency	El Niño	-10.4 [-34.7, 18.1]	0.711	-4.8 [-14.0, 2.5]	-1.7 [-17.6, 11.7]
Hot/dry	Wet-day frequency	La Niña	33.0 [0.0, 53.1]	0.433	4.7 [-6.3, 12.6]	11.5 [-2.2, 32.3]
Hot/dry	Rx1day	El Niño	-27.2 [-48.2, 4.2]	0.433	-17.1 [-27.6, 18.3]	-2.1 [-28.5, 62.6]
Hot/dry	Rx1day	La Niña	-12.6 [-29.1, 29.3]	0.681	6.2 [-21.6, 35.0]	23.0 [-24.3, 80.7]
Hot/dry	CDD	El Niño	-9.6 [-27.9, 15.5]	0.711	4.8 [-11.2, 25.3]	13.3 [-2.8, 29.3]
Hot/dry	CDD	La Niña	-18.5 [-36.9, 9.2]	0.433	-3.9 [-17.4, 6.2]	0.6 [-14.8, 17.9]

276
277

278
279

Table 3. Cross-fitted QDM signal-preservation audit. Values in parentheses are QDM minus raw ensemble responses (percentage points); asymmetry is the neutral-centred sum.

Season	Metric	El Niño category (shift)	La Niña category (shift)	Observed asymmetry	Raw asymmetry	QDM asymmetry
Rainy	PRCPTOT	Preserved (−2.3)	Amplified (−5.0)	−3.5	−23.4	−28.0
Rainy	Wet-day frequency	Indeterminate (−2.6)	Attenuated (+2.1)	−6.1	−12.2	−10.6
Rainy	Rx1day	Attenuated (+7.1)	Amplified (−15.4)	−23.0	−26.4	−34.6
Rainy	CDD	Preserved (−1.1)	Attenuated (−12.8)	8.8	22.8	7.6
Hot/dry	PRCPTOT	Reversed (+24.8)	Indeterminate (+20.7)	−15.1	−8.3	52.7
Hot/dry	Wet-day frequency	Indeterminate (+3.1)	Indeterminate (+6.7)	15.5	0.4	8.8
Hot/dry	Rx1day	Attenuated (+15.0)	Amplified (+16.8)	−23.9	15.7	20.9
Hot/dry	CDD	Indeterminate (+8.4)	Indeterminate (+4.4)	−24.4	1.0	6.4

280
281




Figure 1. Frozen observed ONI record and management-season ENSO classification. Persistent episodes require five overlapping three-month windows at $|ONI| \geq 0.5^\circ\text{C}$. Transition/unclassified seasons are excluded from phase composites.




Figure 2. Observed El Niño and La Niña responses relative to Neutral for four primary endpoints. Points are median station responses; bars are event-level 95% bootstrap intervals.





Figure 3. Observed, raw-model, and blocked cross-fitted-QDM ensemble responses relative to Neutral. Model ensembles use exact-member Niño-3.4 classifications and equal model weight.



R reversed; At attenuated; P preserved; Am amplified; NZ raw response <5%

Figure 4. QDM preservation category and neutral-centred ENSO asymmetry. The phase contrast and true asymmetry are reported as separate estimands.