

# Assignment 11

Surbhi Agarwal

**Abstract**—This document illustrates linear transformation matrices with respect to a set of linearly independent eigenvectors.

Hence, matrix of  $\mathbf{T} - \mathbf{S}$  with respect to  $\mathbf{B}$  can be represented as

$$[\mathbf{T} - \mathbf{S}]_B = \begin{pmatrix} \lambda_1 - \alpha & 0 & \dots & 0 \\ 0 & \lambda_2 - \alpha & \dots & 0 \\ \vdots & \ddots & & \\ 0 & \dots & 0 & \lambda_n - \alpha \end{pmatrix} \quad (2.0.7)$$

## 1 PROBLEM

Let  $\mathbf{S} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be given by  $\mathbf{S}(\mathbf{v}) = \alpha\mathbf{v}$ , for a fixed  $\alpha \in \mathbb{R}, \alpha \neq 0$ . Let  $\mathbf{T} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a linear transformation such that  $\mathbf{B} = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  is a set of linearly independent eigenvectors of  $\mathbf{T}$ . Then

- 1) The matrix of  $\mathbf{T}$  with respect to  $\mathbf{B}$  is diagonal
- 2) The matrix of  $(\mathbf{T} - \mathbf{S})$  with respect to  $\mathbf{B}$  is diagonal
- 3) The matrix of  $\mathbf{T}$  with respect to  $\mathbf{B}$  is not necessarily diagonal, but is upper triangular
- 4) The matrix of  $\mathbf{T}$  with respect to  $\mathbf{B}$  is diagonal but the matrix of  $(\mathbf{T} - \mathbf{S})$  with respect to  $\mathbf{B}$  is not diagonal.

## 2 SOLUTION

Given that  $\mathbf{T} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a linear transformation and  $\mathbf{B}$  represents a set of linearly independent eigenvectors of  $\mathbf{T}$  given as follows

$$\mathbf{B} = \{\mathbf{v}_1, \dots, \mathbf{v}_n\} \quad (2.0.1)$$

So,

$$\mathbf{T}(\mathbf{v}_i) = \mathbf{A}\mathbf{v}_i = \lambda_i\mathbf{v}_i \quad (2.0.2)$$

where  $\lambda_i$  represents the eigenvalue corresponding to  $\mathbf{v}_i$ . Hence, the matrix  $\mathbf{T}$  with respect to  $\mathbf{B}$  can be represented as

$$[\mathbf{T}]_B = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \ddots & & \\ 0 & \dots & 0 & \lambda_n \end{pmatrix} \quad (2.0.3)$$

And,

$$(\mathbf{T} - \mathbf{S})\mathbf{v}_i = \mathbf{T}(\mathbf{v}_i) - \mathbf{S}(\mathbf{v}_i) \quad (2.0.4)$$

$$= \lambda_i\mathbf{v}_i - \alpha\mathbf{v}_i \quad (2.0.5)$$

$$= (\lambda_i - \alpha)\mathbf{v}_i \quad (2.0.6)$$

1. The matrix of $\mathbf{T}$ w.r.t to $\mathbf{B}$ is diagonal	True, as seen from (2.0.3)
2. The matrix of $(\mathbf{T} - \mathbf{S})$ w.r.t $\mathbf{B}$ is diagonal	True, as seen from (2.0.7)
3. The matrix of $\mathbf{T}$ with respect to $\mathbf{B}$ is not necessarily diagonal but is upper triangular	False, as already proved $[\mathbf{T}]_B$ is diagonal
4. The matrix of $\mathbf{T}$ with respect to $\mathbf{B}$ is diagonal but the matrix of $(\mathbf{T} - \mathbf{S})$ with respect to $\mathbf{B}$ is not diagonal	False, as already proved $[\mathbf{T} - \mathbf{S}]_B$ is diagonal

TABLE 1: Verifying the given options

## 3 EXAMPLE

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  where

$$\mathbf{T}(x) = \mathbf{A}\mathbf{x} = \begin{pmatrix} 4 & -2 \\ 3 & -3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \quad (3.0.1)$$

Here, the eigenvalues of the above transformation matrix are  $\lambda_1 = 3, \lambda_2 = -2$ . And the corresponding eigenvectors are  $\mathbf{v}_1 = \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \mathbf{v}_2 = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$ . Thus,

$$\mathbf{B} = \{\mathbf{v}_1, \mathbf{v}_2\} \quad (3.0.2)$$

Now,

$$\mathbf{T}(\mathbf{v}_1) = \mathbf{A}\mathbf{v}_1 \quad (3.0.3)$$

$$= \begin{pmatrix} 4 & -2 \\ 3 & -3 \end{pmatrix} \begin{pmatrix} 2 \\ 1 \end{pmatrix} \quad (3.0.4)$$

$$= \begin{pmatrix} 6 \\ 3 \end{pmatrix} \quad (3.0.5)$$

$$= 3 \begin{pmatrix} 2 \\ 1 \end{pmatrix} \quad (3.0.6)$$

$$= \lambda_1 \mathbf{v}_1 \quad (3.0.7)$$

And,

$$\mathbf{T}(\mathbf{v}_2) = \mathbf{A}\mathbf{v}_2 \quad (3.0.8)$$

$$= \begin{pmatrix} 4 & -2 \\ 3 & -3 \end{pmatrix} \begin{pmatrix} 1 \\ 3 \end{pmatrix} \quad (3.0.9)$$

$$= \begin{pmatrix} -2 \\ -6 \end{pmatrix} \quad (3.0.10)$$

$$= -2 \begin{pmatrix} 1 \\ 3 \end{pmatrix} \quad (3.0.11)$$

$$= \lambda_2 \mathbf{v}_2 \quad (3.0.12)$$

$$(3.0.13)$$

For any vector  $\mathbf{v} \in \mathbb{R}^2$ ,  $\mathbf{v} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2$

$$[\mathbf{v}]_B = \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} \quad (3.0.14)$$

$$\mathbf{T}(\mathbf{v}) = \mathbf{T}(c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2) \quad (3.0.15)$$

$$= c_1 \mathbf{T}(\mathbf{v}_1) + c_2 \mathbf{T}(\mathbf{v}_2) \quad (3.0.16)$$

$$= c_1 \lambda_1 \mathbf{v}_1 + c_2 \lambda_2 \mathbf{v}_2 \quad (3.0.17)$$

$$[\mathbf{T}(\mathbf{v})]_B = \begin{pmatrix} \lambda_1 c_1 \\ \lambda_2 c_2 \end{pmatrix} \quad (3.0.18)$$

$$= \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} \quad (3.0.19)$$

$$= \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} [\mathbf{v}]_B \quad (3.0.20)$$

$$= \begin{pmatrix} 3 & 0 \\ 0 & -2 \end{pmatrix} [\mathbf{v}]_B \mathbf{S}(\mathbf{v}) = \alpha \mathbf{v}, \alpha \neq 0 \quad (3.0.21)$$

$$= \alpha(c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2) \quad (3.0.22)$$

$$= \alpha c_1 \mathbf{v}_1 + \alpha c_2 \mathbf{v}_2 \quad (3.0.23)$$

$$(3.0.24)$$

$$[\mathbf{S}(\mathbf{v})]_B = \begin{pmatrix} \alpha c_1 \\ \alpha c_2 \end{pmatrix} \quad (3.0.25)$$

$$[(\mathbf{T} - \mathbf{S})(\mathbf{v})]_B = \begin{pmatrix} \lambda_1 c_1 - \alpha c_1 \\ \lambda_2 c_2 - \alpha c_2 \end{pmatrix} \quad (3.0.26)$$

$$= \begin{pmatrix} \lambda_1 - \alpha & 0 \\ 0 & \lambda_2 - \alpha \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \end{pmatrix} \quad (3.0.27)$$

$$= \begin{pmatrix} \lambda_1 - \alpha & 0 \\ 0 & \lambda_2 - \alpha \end{pmatrix} [\mathbf{v}]_B \quad (3.0.28)$$

$$= \begin{pmatrix} 3 - \alpha & 0 \\ 0 & -2 - \alpha \end{pmatrix} [\mathbf{v}]_B \quad (3.0.29)$$

Hence, shown from (3.0.21) and (3.0.29) that the matrix of  $\mathbf{T}$  and of  $\mathbf{T} - \mathbf{S}$  w.r.t to  $\mathbf{B}$  is diagonal.