

Thoracic Spine MRI

ACG: A-0058 (AC)

[Link to Codes](#)

This MCG care guideline may overlap CMS guidance located in CMS statutes and regulations that are not currently part of MCG's Medicare Compliance content. For Medicare Advantage beneficiaries, before referring to MCG care guidelines, users should first follow all guidance related to this topic located in CMS National Coverage Determinations (NCDs), Local Coverage Determinations (LCDs), and other statutes and regulations. Content used in MCG care guidelines may be used to supplement but does not replace, modify, or supersede existing Medicare regulations or applicable NCDs or LCDs.

- Clinical Indications
- Evidence Summary
 - Background
 - Criteria
 - Rationale
 - Related CMS Coverage Guidance
- References
- Footnotes
- Codes

Clinical Indications

- Thoracic spine MRI may be indicated for **1 or more** of the following(1)(2)(3)(4)(5)(6):
 - ☐ Cancer or neoplasm evaluation, staging, or surveillance needed, as indicated by **1 or more** of the following(21)(22)(23)(24)(25)(26):[\[1\]](#)
 - Bone neoplasm (benign or malignant) involving spine, known or suspected (eg, chondrosarcoma, chordoma, Ewing sarcoma family of tumors, giant cell tumor of bone, osteosarcoma, metastatic tumors), as indicated by **1 or more** of the following(33)(34)(35):
 - Initial evaluation or staging(36)
 - Monitoring disease progression or response to treatment
 - Post-treatment surveillance
 - Histiocytic or dendritic cell neoplasm[A] with involvement of vertebrae, known or suspected(37)(40)(41)
 - Localized midline thoracic back pain and **1 or more** of the following(42):
 - Bone scan positive(36)(43)(44)
 - Pain occurs mainly at night.(45)
 - Persistent back pain in patient older than 50 years(46)
 - Personal history or concurrent diagnosis of malignancy(27)(28)(29)(47)(48)
 - Weakness, rapidly progressing(46)
 - Weight loss, unexplained(46)
 - Myelopathy signs or symptoms (eg, motor weakness, bowel or bladder dysfunction)(27)(28)(29)(49)(50)
 - Primary central nervous system tumor (eg, adult-type or pediatric-type diffuse glioma, ependymoma, medulloblastoma, other central nervous system embryonal tumor, primary central nervous system lymphoma, meningioma), and need for **1 or more** of the following(34)(51)(52)(53)(54)(55):
 - Brain tumor, and need for evaluation for spinal involvement
 - Initial evaluation, staging, or treatment planning
 - Monitoring disease progression or response to treatment
 - Post-treatment surveillance
 - Primary non-central nervous system tumor, and need for diagnosis, staging, or monitoring of known or suspected spinal cord, spine, or leptomeningeal metastasis, as indicated by **1 or more** of the following(34)(51)(54)(55)(56):
 - Known spinal cord, spine, or leptomeningeal metastasis, and need for monitoring disease progression or response to treatment
 - Post-treatment surveillance

- Suspected spinal cord, spine, or leptomeningeal metastasis (eg, due to myelopathy, pain, finding on other imaging), and need for evaluation, staging, or treatment planning
 - Screening for central nervous system tumor in patient with high-risk genetic disorder or syndrome (eg, neurofibromatosis type 2, von Hippel-Lindau syndrome)(55)(57)(58)(59)(60)
- Cerebrospinal fluid leak, suspected(21)(61)(62):[N](#)
- ☐ Congenital spinal conditions, suspected, as indicated by **1 or more** of the following(21)(63)(64)(65):[N](#)
 - Diastematomyelia (ie, split cord malformation)(66)
 - Dorsal dermal sinus(67)
 - Hemangioma(68)(69)(70)(71)
 - Hydromyelia
 - Intradural lipoma
 - Myelocele
 - Myelomeningocele(66)
 - Split notochord syndrome(67)
- ☐ Infection, known or suspected (eg, vertebral osteomyelitis, disk space infection, epidural abscess), as indicated by **ALL** of the following(6)(21)(72)(73)(74)(75):[N](#)
 - Risk factors for spinal infection, as indicated by **1 or more** of the following:
 - Bone scan or plain x-ray suggestive for infection(82)
 - C-reactive protein or erythrocyte sedimentation rate elevated(83)(84)(85)
 - Fever(86)(87)
 - History of intravenous drug abuse, other source of infection, or recent invasive procedure(88)
 - Immunosuppression(89)
 - Positive blood culture(83)(90)
 - Recent history of spinal surgery(91)
 - Tuberculosis, concurrent or suspected(81)(89)(92)(93)
 - Suspected spinal infection (eg, localized midline back pain, limp, refusal to walk)(83)(86)
- ☐ Inflammatory or demyelinating process, suspected (eg, clinically isolated syndrome, multiple sclerosis, other demyelinating disease), as indicated by **1 or more** of the following(21)(94)(95):[N](#)
 - Multiple sclerosis, suspected, as indicated by **1 or more** of the following(96)(97)(100)(106):
 - Ascending numbness or tingling (eg, from foot to trunk)(99)
 - Brown-Sequard syndrome(99)
 - Conditions mimicking multiple sclerosis (eg, Sjogren syndrome, systemic lupus erythematosus, antiphospholipid syndrome) cannot yet be excluded.(50)(107)(108)
 - MRI of brain nondiagnostic for multiple sclerosis(109)
 - Signs or symptoms of myelopathy or myelitis(87)(110)(111)
 - Transverse myelitis in spinal cord, suspected (idiopathic or in conjunction with multiple sclerosis), as indicated by **1 or more** of the following(89)(112)(113)(114):
 - Bilateral signs or symptoms of involvement of appropriate level of spinal cord
 - Clearly defined sensory level
 - Sudden onset of sensory, motor, and autonomic dysfunction attributable to appropriate level of spinal cord
- ☐ [N](#) Pain localized to thoracic back or radicular in nature, subacute or chronic, as indicated by **1 or more** of the following(115)(116):[N](#)
 - Adult and **ALL** of the following:
 - Failure to improve after 6 or more weeks of nonoperative treatment, as indicated by **1 or more** of the following(74)(87)(117)(118)(119)(120)(121):
 - Activity modification(122)
 - Exercise(123)
 - Pharmacotherapy (eg, analgesics, anti-inflammatory medications)(124)(125)
 - Physical therapy(125)(126)(127)
 - Patient being considered for invasive treatment (eg, epidural steroids, surgery)
 - Significant interference with daily function
 - Child or adolescent and **ALL** of the following(6):
 - Age younger than 18 years
 - Concern for serious underlying pathology (eg, malignancy, infection, inflammatory process), as indicated by **1 or more** of the following:
 - Associated signs suggestive of cancer (eg, easy bruising, abdominal mass, lymphadenopathy)
 - Fever
 - Gait abnormalities or refusal to walk or crawl
 - Morning stiffness
 - Neurologic deficit
 - Nocturnal pain or constant pain
 - Pain lasting more than 4 weeks or worsening pain

- Unintentional weight loss
- ☐ Postoperative spinal complications, suspected, as indicated by **1 or more** of the following(128)(129)(130):[N](#)
 - Postoperative hemorrhage or hematoma(91)
 - Pseudomeningocele
 - Significant new neurologic findings (eg, spinal cord impingement)
 - Significant new pain symptoms (eg, recurrent or new radicular symptoms, delayed pain at fusion site)
 - Spondylodiscitis
 - Sterile arachnoiditis
- ☐ Scoliosis, as indicated by **1 or more** of the following(21)(131)(132)(133):[N](#)
 - Congenital scoliosis
 - Early-onset scoliosis (age 9 years or younger)
 - Neurofibromatosis
 - Presurgical planning for adolescent idiopathic scoliosis to assess possible neural axis malformation, as indicated by **1 or more** of the following(131)(132)(136)(137):
 - Abnormal neurologic findings on clinical examination(138)
 - Age at first visit 10 years or younger
 - Kyphosis at curve apex
 - Left-sided thoracic curvature
 - Male gender
 - Pain, moderate to severe(7)(138)
 - Rapid curve progression (ie, more than 1 degree per month)
 - Short segment curve (ie, less than 6 vertebral segments)
 - Thoracic kyphosis 30 degrees or greater
 - Vertebral abnormalities (eg, hemivertebrae, block vertebrae) detected on x-ray
 - Spinal cord compression, myelopathy, or cauda equina syndrome, suspected (eg, positive Babinski sign, depressed ankle deep tendon reflexes, gait abnormality, hyperreflexia, sensory or motor deficits)(21)(29)(89)(139)(140)(141)[N](#)
 - Spinal cord infarction or ischemia, suspected (eg, sudden-onset back pain, sensory or motor deficits, urinary retention)(142)[N](#)
 - Spinal cord stimulator placement planning needed, and no MRI of region within 12 months prior to placement(147)(148)(149)[N](#)
- ☐ Spinal dysraphism, suspected, as indicated by **1 or more** of the following(66)(153)(154)(155)(156)(157)(158):[N](#)
 - For infant, as indicated by **1 or more** of the following(64)(160)(161)(162):
 - Absent reflexes
 - Aplasia cutis congenita
 - Decreased spontaneous leg movement
 - Decreased urinary stream
 - Deviation of gluteal cleft
 - Foot asymmetry(67)
 - Leg atrophy
 - Midline lumbosacral skin lesions (eg, atypical dimple, dermal sinus, lipoma, tail)(7)(67)
 - Unilateral lumbosacral hypertrichosis
 - For toddler: developmental delay in walking, abnormal gait
 - For older child: spasticity, hyperreflexia, bowel or bladder disturbance, including incontinence(67)
- ☐ Spinal stenosis of thoracic spine, suspected, as indicated by **ALL** of the following(21)(163):[N](#)
 - Patient being considered for invasive treatment(165)(166)
 - Progressive or disabling symptoms of thoracic spinal stenosis, as indicated by **1 or more** of the following(164):
 - Hyperactive reflexes
 - Muscle weakness
 - Sensory loss
 - Spasticity(167)
 - Spondylolysis, known or suspected (eg, plain radiography-negative early spondylolysis, radiculopathy in patient with spondylolysis)(168)(169)(170)[N](#)
 - Stereotactic spine radiotherapy treatment planning(172)(173)(174)(175)(176)[N](#)
- ☐ Syringomyelia in thoracic spine, suspected, as indicated by **1 or more** of the following(21)(179)(180)(181):[N](#)
 - Muscle wasting in appropriate thoracic spine dermatomes
 - Sensory loss in appropriate thoracic spine dermatomes(182)
 - Weakness in appropriate thoracic spine dermatomes
- ☐ Tethered cord, suspected, as indicated by **1 or more** of the following(64)(66)(153)(156)(183)(184):[N](#)
 - Anorectal malformation(160)(185)(186)
 - Cutaneous manifestations of occult spina bifida (eg, nevus, lipoma, tufts of hair, hemangioma, dimple overlying spine, asymmetric gluteal cleft, dermal sinus tract)(160)(161)(187)
 - Gait abnormality or difficulty
 - Urinary dribbling or lack of bladder control(67)(86)
 - Urodynamic tests abnormal

- ☐ Trauma or fracture of thoracic spine, known or suspected, as indicated by **1 or more** of the following(188)(189)(190)(191)(192):
 - Epidural hematoma, suspected(199)(203)
 - Fall from height greater than 10 feet (3.1 meters)(204)
 - Ligamentous injury, suspected(205)(206)(207)(208)
 - Malignancy, known or suspected(202)(209)(210)
 - Neurologic symptoms associated with traumatic mechanism of injury, as indicated by **1 or more** of the following(21)(204)(211)(212)(213)(214):
 - Ejection from motorcycle(205)(215)
 - High-speed motor vehicle accident (greater than 50 mph (80.5 kph))(205)(215)
 - Intoxicated state(216)
 - Minimal trauma in osteoporotic patient with suspected fragility fracture
 - Motor vehicle accident, with acute hyperflexion of spine over seat belt(217)
 - Osteoporosis, known or suspected(202)(218)
 - Pediatric patient(195)(219)(220)(221)(222)
 - Results of plain x-ray or CT scan nondiagnostic or show abnormal spine motion or burst fracture(193)(194)(201)(223)
 - Traumatic disk herniation, suspected(206)(224)
- ☐ Tuberculosis of spine, known or suspected, as indicated by **1 or more** of the following(81)(225)(226):
 - Diagnostic evaluation of suspected spinal tuberculosis(229)
 - Preparation for surgical management
- ☐ Repeat evaluation of specific area or structure with same imaging modality, and **ALL** of the following:
 - Clinical need for repeat imaging, as indicated by **1 or more** of the following:
 - Change in clinical status (eg, worsening symptoms or new associated symptoms), and findings may impact treatment
 - Need for interval reassessment that may impact treatment plan
 - Need for re-imaging either prior to or after performance of invasive procedure
 - Prior imaging results of specific area or structure with same imaging modality documented and available for comparison

Evidence Summary

Background

Some gadolinium-based contrast agents used for diagnostic and treatment guidance with MRI or MRA have been associated with nephrogenic systemic fibrosis in patients with chronic kidney disease or acute kidney injury.(7)(8)(9)(10)(11) **(EG 2)** Gadolinium deposition in the brain has been reported by multiple research groups; however, the clinical significance of this finding is unclear.(7)(8)(12)(13) **(EG 2)** Specialty society guidelines recommend weighing the risks vs benefits when using gadolinium-based contrast agents for MRI.(14) **(EG 2)** MRI-based techniques may be contraindicated in patients with ferromagnetic intracranial aneurysm clips and certain types of ferromagnetic foreign bodies and cochlear implants.(15) **(EG 2)** Although these implants have traditionally included cardiac pacemakers and implantable cardioverter-defibrillators, multiple trials have validated the safe and effective use of pacemaker systems designed for MRI use, and a guideline supported by multiple specialty societies states that many patients with implantable cardiac devices, including those not originally intended for MRI scanning, can safely undergo MRI scanning under defined clinical indications.(8)(16)(17)(18)(19)(20) **(EG 1)**

Criteria

The evidence for the clinical indications found in this guideline includes 186 published peer reviewed articles, 12 specialty society or other evidence-based guidelines, and 13 book sections.

For cancer or neoplasm evaluation or staging, evidence demonstrates at least moderate certainty of at least moderate net benefit. **(RG A1)** Whole-spine MRI is the preferred imaging modality for suspected metastatic disease to the spine in patients with known primary malignancy; metastatic epidural spinal cord compression is seen in 5% to 10% of patients with cancer.(27)(28)(29) **(EG 2)** Spinal cord tumor in children may present with persistent localized back pain that is worse at night.(6)(30)(31) **(EG 2)** Additional findings in children with spinal cord tumor may include burning pain with paresthesia, gait disturbance, paraspinal muscle spasm, progressive motor weakness, progressive scoliosis, and sensory deficits.(32) **(EG 2)**

For cerebrospinal fluid leak, evidence demonstrates a net benefit, but of less than moderate certainty, and may consist of a consensus opinion of experts, case studies, and common standard care. **(RG A2)** A specialty society guideline notes that MRI of the spine may be useful to discover the location of a cerebrospinal fluid leak in patients with intracranial hypotension.(21) **(EG 2)**

For congenital spinal conditions, evidence demonstrates a net benefit, but of less than moderate certainty, and may consist of a consensus opinion of experts, case studies, and common standard care. **(RG A2)** A specialty society guideline indicates that MRI is an appropriate imaging modality for assessment of congenital spinal conditions.(21) **(EG 2)**

For infection of the spine, evidence demonstrates at least moderate certainty of at least moderate net benefit. **(RG A1)** MRI is a preferred option for diagnosis of suspected spinal infection and inflammation, including spondylitis, epidural abscess, osteomyelitis, and spinal tuberculosis.(76)(77)(78)(79)(80) **(EG 2)** Because spinal tuberculosis can present in noncontiguous vertebral bodies, patients with known spinal tuberculosis should undergo MRI imaging of the entire spine.(81) **(EG 2)**

For inflammatory or demyelinating processes, evidence demonstrates a net benefit, but of less than moderate certainty, and may consist of a consensus opinion of experts, case studies, and common standard care. **(RG A2)** Diagnostic criteria have been developed for inflammatory or demyelinating processes, such as clinically isolated syndrome or multiple sclerosis (most commonly manifested as unilateral optic neuritis), brainstem syndrome, or partial myelitis, based on the demonstration of central nervous system lesions disseminated in space and disseminated in time on MRI imaging of the brain and, if appropriate, on spinal cord imaging.(96)(97)(98)(99)(100)(101) **(EG 2)** Spinal cord involvement is seen on MRI imaging in 80% to 90% of patients with multiple sclerosis; asymptomatic spinal cord lesions may be present in up to 40% of patients with clinically isolated syndrome, with lesions occurring more commonly in the cervical spine than in the thoracic spine.(102)(103)(104)(105) **(EG 2)** An expert consensus panel recommends spinal cord imaging for patients with suspected demyelinating disease who present with signs or symptoms of spinal cord involvement or if the number of lesions on brain MRI is insufficient to meet established criteria for multiple sclerosis.(96) **(EG 2)**

For pain localized to the thoracic back or radicular in nature, evidence demonstrates at least moderate certainty of at least moderate net benefit. **(RG A1)** Review articles consider MRI to be an appropriate initial imaging modality for pain that significantly interferes with daily function and for which invasive treatment is being considered.(115)(116) **(EG 2)** For children and adolescents with back or neck pain, a specialty society guideline notes that, while imaging is not generally indicated for pain that is self-limited or not associated with signs or symptoms of an underlying disorder, MRI is appropriate for evaluating pain associated with potential red flags (eg, associated gait abnormalities, unintentional weight loss, persistence over more than 4 weeks).(6) **(EG 2)**

For postoperative spinal complications, evidence demonstrates a net benefit, but of less than moderate certainty, and may consist of a consensus opinion of experts, case studies, and common standard care. **(RG A2)** Review articles indicate that MRI is effective at demonstrating common complications after spinal surgery.(128)(129)(130) **(EG 2)**

For scoliosis, evidence demonstrates a net benefit, but of less than moderate certainty, and may consist of a consensus opinion of experts, case studies, and common standard care. **(RG A2)** Review articles conclude that MRI is not routinely indicated for pain in the absence of abnormal neurologic examination findings or an atypical curve, such as a left-sided curve (levoscoliosis), a short segment, or kyphosis near the curve apex.(132)(134)(135)(136) **(EG 2)** However, it is useful preoperatively in evaluating the neural axis.(137) **(EG 2)**

For spinal cord compression, evidence demonstrates at least moderate certainty of at least moderate net benefit. **(RG A1)** MRI is the preferred imaging modality for detection and confirmation of malignant spinal cord compression.(49) **(EG 2)** Literature reviews of acute transverse myelitis recommend urgent neuroimaging with MRI to rule out cord compression; however, if MRI is not available or is contraindicated, CT myelography is indicated.(95)(112) **(EG 2)**

For spinal cord infarction or ischemia, evidence demonstrates a net benefit, but of less than moderate certainty, and may consist of a consensus opinion of experts, case studies, and common standard care. **(RG A2)** MRI of the spine is the preferred diagnostic procedure to detect spinal cord ischemia. The clinical manifestations may evolve over minutes to hours, vary depending on the vascular territory involved, may be bilateral or unilateral, and may present transiently (also called spinal transient ischemic attacks).(142)(143)(144) **(EG 2)** Etiologies may vary (eg, aortic dissection, atherosclerosis, coagulopathy, embolization, mechanical compression, systemic hypoperfusion, vasculitis).(142)(144)(145)(146) **(EG 2)** For patients experiencing spinal infarct or ischemia, timely identification facilitates reperfusion intervention, which may improve outcomes.(143) **(EG 2)**

For spinal cord stimulator placement planning, evidence demonstrates a net benefit, but of less than moderate certainty, and may consist of a consensus opinion of experts, case studies, and common standard care. **(RG A2)** Preoperative MRI of the spine is recommended within 12 months prior to the spinal cord stimulator placement to evaluate potential pathology that could alter the diagnosis or negatively impact the outcome and to assess the need for alternative approaches to placement of the stimulator electrodes.(150)(151)(152) **(EG 2)**

For spinal dysraphism, evidence demonstrates a net benefit, but of less than moderate certainty, and may consist of a consensus opinion of experts, case studies, and common standard care. **(RG A2)** Spinal dysraphism includes a variety of malformations with incomplete fusion of midline mesenchymal, bony, and neurologic structures; a broad bony defect of the vertebral arch is seen at the level of the malformation.(66)(154)(155) **(EG 2)** Spinal dysraphism may be associated with neurogenic scoliosis. Intraspinal pathology has been reported in 12% to 50% of patients with congenital scoliosis; the MRI findings may include anomalies such as filum lipoma, syrinx, or Chiari malformation.(135)(159) **(EG 2)**

For spinal stenosis of the thoracic spine, evidence demonstrates at least moderate certainty of at least moderate net benefit. **(RG A1)** A literature review on the imaging of spinal stenosis noted that CT scan, CT myelography, and MRI can all visualize the ventral and dorsal contributions to spinal canal compromise.(164) **(EG 2)**

For spondylolysis, evidence demonstrates a net benefit, but of less than moderate certainty, and may consist of a consensus opinion of experts, case studies, and common standard care. **(RG A2)** Prior to the development of detectable stress fracture on plain radiograph

or CT scan, bone marrow edema of the pars interarticularis may be identified using thin-slice MRI in patients with spondylolysis.(6)(168) (169) **(EG 2)** MRI may also detect associated ligament disruption in traumatic spondylolysis and is used to evaluate neurologic findings, such as radiculopathy, in patients with spondylolysis.(6)(170)(171) **(EG 2)**

For stereotactic spine radiotherapy, evidence demonstrates a net benefit, but of less than moderate certainty, and may consist of a consensus opinion of experts, case studies, and common standard care. **(RG A2)** CT scan and MRI are increasingly used for treatment planning prior to stereotactic spine radiotherapy for spinal metastases.(172)(173)(175)(177)(178) **(EG 2)**

For syringomyelia, evidence demonstrates a net benefit, but of less than moderate certainty, and may consist of a consensus opinion of experts, case studies, and common standard care. **(RG A2)** A specialty society guideline indicates that MRI is an appropriate imaging modality for assessment of syringomyelia.(21) **(EG 2)** When Chiari malformation is suspected on cranial CT scan, a review article recommends MRI of the whole spine.(181) **(EG 2)**

For tethered cord, evidence demonstrates a net benefit, but of less than moderate certainty, and may consist of a consensus opinion of experts, case studies, and common standard care. **(RG A2)** MRI is considered to be the study of choice when evaluating tethered cord, as it demonstrates the level of the conus and nature of the filum, identifies the cause of the tethering, and provides anatomic details needed for preoperative planning.(64)(153) **(EG 2)** A retrospective study of 790 patients with anorectal malformation reported that the incidence of tethered cord was 36%, and noted that screening for tethered cord using MRI or ultrasound has become standard practice for newborns with anorectal malformations.(185) **(EG 2)**

For trauma or fracture of the thoracic spine, evidence demonstrates at least moderate certainty of at least moderate net benefit. **(RG A1)** Systematic reviews of studies of patients with spinal cord injuries who did not have abnormalities on normal plain x-rays or CT scans demonstrated abnormal spine MRI findings in 57% of children and 93% of adults.(193)(194) **(EG 1)** In a retrospective cohort study of 206 pediatric patients evaluated for spinal injuries in a trauma center, MRI was obtained more frequently in patients between 0 and 3 years of age (67%) compared with patients between 4 and 9 years of age (20%). The authors hypothesized that this disparity may be due to difficulty in obtaining reliable neurologic examinations in infants and toddlers. MRI is also a better tool for imaging cartilage, of which a significant portion of the spinal column in this age group is composed.(195) **(EG 2)** A study of patients with 152 thoracic or lumbar vertebral compression fractures showed that specific combinations of MRI findings were able to distinguish malignant, osteoporotic, and infective etiologies with accuracy ranging from 76% to 98%, with biopsy as the gold standard.(196) **(EG 2)** MRI is helpful in assessment of acute and subacute posttraumatic spinal cord injury, including damage to disks(197) and spinal cord ligaments, as well as assessment of spinal canal stenosis, edema, and hematoma.(198)(199) **(EG 2)** MRI may also be helpful at predicting higher-risk compression fractures that result in nonunion.(200) **(EG 2)** A specialty society practice guideline stated that MRI may be used to evaluate the acute thoracic and thoracolumbar traumatic spine to assess for soft tissue injury, detect fractures, and ascertain the need for surgery. The authors noted that a small study population and heterogeneity limited conclusions about the use of MRI in this setting; further long-term studies with a higher level of evidence were recommended.(201) **(EG 2)** Another specialty society guideline recommended that MRI be performed in patients with osteoporotic or malignancy-related vertebral compression fractures who are being considered for vertebral augmentation (ie, vertebroplasty, kyphoplasty) to help determine the age and healing status of the fracture.(202) **(EG 2)**

For tuberculosis of the spine, evidence demonstrates at least moderate certainty of at least moderate net benefit. **(RG A1)** Tuberculous infection of the spine (Pott disease) presents most commonly in the thoracic spine and can lead to kyphosis.(227)(228)(229) **(EG 2)** MRI is considered the most accurate imaging modality for diagnosis and classification of tuberculosis, including early spinal cord changes.(81)(228) **(EG 2)**

Rationale

Use of this MCG care guideline helps the clinician determine if a particular treatment, medication, or service might be appropriate for a specific patient, taking into account their unique health complexities.

Use of these evidence-based clinical criteria to support decision making benefits the patient by identifying patient-specific complex clinical factors and conditions, promoting personalized treatment. Utilizing evidence-based clinical criteria promotes patient safety by helping ensure that potential patient benefits outweigh the risks. In addition, the use of evidence-based guidelines can increase consistency in treatment thresholds, leading to less variation in care and promoting equitable treatment among patients.

Related CMS Coverage Guidance

This guideline supplements but does not replace, modify, or supersede existing Medicare regulations or applicable National Coverage Determinations (NCDs) or Local Coverage Determinations (LCDs).

Code of Federal Regulations (CFR): 42 CFR 419.22(230); 42 CFR 422.101(231)

Internet-Only Manual (IOM) Citations: CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 14 - Medical Devices(232); CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 15 - Covered Medical and Other Health Services(233); CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 16 - General Exclusions from Coverage(234)

Medicare Coverage Determinations: Medicare Coverage Database(235)

References

1. Klein JP. A practical approach to spine imaging. *Continuum : Lifelong Learning in Neurology* 2015;21(1 Spinal Cord Disorders):36-51. DOI: 10.1212/01.CON.0000461083.33500.ec. [Context Link 1] View abstract...
2. Rossi A, Martinetti C, Morana G, Severino M, Tortora D. Diagnostic approach to pediatric spine disorders. *Magnetic Resonance Imaging Clinics of North America* 2016;24(3):621-644. DOI: 10.1016/j.mric.2016.04.001. [Context Link 1] View abstract...
3. Singh K, Mechtler LL, Klein JP. Imaging of spinal cord disorders. *Continuum : Lifelong Learning in Neurology* 2016;22(5, Neuroimaging):1595-1612. DOI: 10.1212/CON.0000000000000381. [Context Link 1] View abstract...
4. George E, Guenette JP, Lee TC. Introduction to neuroimaging. *American Journal of Medicine* 2018;131(4):346-356. DOI: 10.1016/j.amjmed.2017.11.014. [Context Link 1] View abstract...
5. Lavi ES, Pal A, Bleicher D, Kang K, Sidani C. MR imaging of the spine: urgent and emergent indications. *Seminars in Ultrasound, CT and MRI* 2018;39(6):551-569. DOI: 10.1053/j.sult.2018.10.006. [Context Link 1] View abstract...
6. Dahmouch H, et al. Back Pain-Child. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2024 Accessed at: <https://www.acr.org>. [accessed 2024 Nov 01] [Context Link 1, 2, 3, 4, 5, 6, 7]
7. Squires JH, Narayanan S, Tadros S. Fundamentals of pediatric radiology. In: Zitelli BJ, McIntire SC, Nowalk AJ, Garrison J, editors. *Zitelli and Davis' Atlas of Pediatric Physical Diagnosis*. 8th ed. Elsevier; 2022:923-992. [Context Link 1, 2, 3, 4]
8. Primary Panel, et al. 2021 update on safety of magnetic resonance imaging: joint statement from Canadian Cardiovascular Society/Canadian Society for Cardiovascular Magnetic Resonance/Canadian Heart Rhythm Society. *Canadian Journal of Cardiology* 2021;37(6):835-847. DOI: 10.1016/j.cjca.2021.02.012. (Reaffirmed 2024 Jul) [Context Link 1, 2, 3] View abstract...
9. ACR Committee on Drugs and Contrast Media. ACR Manual on Contrast Media. [Internet] American College of Radiology. 2022 Accessed at: <https://www.acr.org/>. [accessed 2024 Sep 18] [Context Link 1]
10. Weinreb JC, et al. Use of intravenous gadolinium-based contrast media in patients with kidney disease: consensus statements from the American College of Radiology and the National Kidney Foundation. *Radiology* 2021;298(1):28-35. DOI: 10.1148/radiol.20202903. (Reaffirmed 2024 Aug) [Context Link 1] View abstract...
11. Thomsen HS. Nephrogenic systemic fibrosis: a serious adverse reaction to gadolinium - 1997-2006-2016. Part 2. *Acta Radiologica* 2016;57(6):643-648. DOI: 10.1177/0284185115626481. [Context Link 1] View abstract...
12. Kanda T, Nakai Y, Hagiwara A, Oba H, Toyoda K, Furui S. Distribution and chemical forms of gadolinium in the brain: a review. *British Journal of Radiology* 2017;90(1079):20170115. DOI: 10.1259/bjr.20170115. [Context Link 1] View abstract...
13. Ramalho J, Semelka RC, Ramalho M, Nunes RH, AlObaidy M, Castillo M. Gadolinium-based contrast agent accumulation and toxicity: an update. *AJNR. American Journal of Neuroradiology* 2016;37(7):1192-1198. DOI: 10.3174/ajnr.A4615. [Context Link 1] View abstract...
14. Gulani V, Calamante F, Shellock FG, Kanal E, Reeder SB, International Society for Magnetic Resonance in Medicine. Gadolinium deposition in the brain: summary of evidence and recommendations. *Lancet Neurology* 2017;16(7):564-570. DOI: 10.1016/S1474-4422(17)30158-8. [Context Link 1] View abstract...
15. ACR Practice Parameter for Performing and Interpreting Magnetic Resonance Imaging (MRI). Resolution 8 [Internet] American College of Radiology. 2022 Accessed at: <https://www.acr.org/>. [created 1992; accessed 2024 Sep 18] [Context Link 1]
16. Lowe MD, Plummer CJ, Manisty CH, Linker NJ, British Heart Rhythm Society. Safe use of MRI in people with cardiac implantable electronic devices. *Heart* 2015;101(24):1950-1953. DOI: 10.1136/heartjnl-2015-308495. [Context Link 1] View abstract...
17. Nazarian S, et al. Safety of magnetic resonance imaging in patients with cardiac devices. *New England Journal of Medicine* 2017;377(26):2555-2564. DOI: 10.1056/NEJMoa1604267. [Context Link 1] View abstract...
18. Nordbeck P, Ertl G, Ritter O. Magnetic resonance imaging safety in pacemaker and implantable cardioverter defibrillator patients: how far have we come? *European Heart Journal* 2015;36(24):1505-1511. DOI: 10.1093/eurheartj/ehv086. [Context Link 1] View abstract...
19. Russo RJ, et al. Assessing the risks associated with MRI in patients with a pacemaker or defibrillator. *New England Journal of Medicine* 2017;376(8):755-764. DOI: 10.1056/NEJMoa1603265. [Context Link 1] View abstract...
20. Bhuvu A, et al. Joint British Society consensus recommendations for magnetic resonance imaging for patients with cardiac implantable electronic devices. *Heart* 2024;110(4):Online. DOI: 10.1136/heartjnl-2022-320810. [Context Link 1] View abstract...
21. ACR-ASNR-SABI-SSR Practice Parameter for the Performance of Magnetic Resonance Imaging (MRI) of the Adult Spine. Resolution 6 [Internet] American College of Radiology. 2023 Accessed at: <https://www.acr.org/>. [created 2001; accessed 2024 Sep 19] [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13]
22. Hazenfield JM, Gaskill-Shipley MF. Neoplastic and paraneoplastic involvement of the spinal cord. *Seminars in Ultrasound, CT and MRI* 2016;37(5):482-497. DOI: 10.1053/j.sult.2016.05.009. [Context Link 1] View abstract...
23. Merhemic Z, Stosic-Opincal T, Thurnher MM. Neuroimaging of spinal tumors. *Magnetic Resonance Imaging Clinics of North America* 2016;24(3):563-579. DOI: 10.1016/j.mric.2016.04.007. [Context Link 1] View abstract...
24. Chokshi FH, Law M, Gibbs WN. Conventional and advanced imaging of spine oncologic disease, nonoperative post-treatment effects, and unique spinal conditions. *Neurosurgery* 2018;82(1):1-23. DOI: 10.1093/neuros/nyx491. [Context Link 1] View abstract...
25. Barwick T, Bretszajn L, Wallitt K, Amiras D, Rockall A, Messiou C. Imaging in myeloma with focus on advanced imaging techniques. *British Journal of Radiology* 2019;92(1095):20180768. DOI: 10.1259/bjr.20180768. [Context Link 1] View abstract...
26. Biermann JS, et al. Bone Cancer. NCCN Clinical Practice Guidelines in Oncology [Internet] National Comprehensive Cancer Network (NCCN). v. 2.2024; 2024 Mar 12 Accessed at: <https://www.nccn.org/>. [accessed 2024 Mar 25] [Context Link 1]

27. Chamberlain MC. Neoplastic myelopathies. *Continuum : Lifelong Learning in Neurology* 2015;21(1 Spinal Cord Disorders):132-145. DOI: 10.1212/01.CON.0000461089.02113.1b. [Context Link 1, 2, 3] View abstract...
28. Scott BJ. Neuro-oncologic emergencies. *Seminars in Neurology* 2015;35(6):675-682. DOI: 10.1055/s-0035-1564684. [Context Link 1, 2, 3] View abstract...
29. Lin AL, Avila EK. Neurologic emergencies in the patients with cancer. *Journal of Intensive Care Medicine* 2017;32(2):99-115. DOI: 10.1177/0885066615619582. [Context Link 1, 2, 3, 4] View abstract...
30. Calloni SF, Huisman TA, Poretti A, Soares BP. Back pain and scoliosis in children: When to image, what to consider. *Neuroradiology Journal* 2017;30(5):393-404. DOI: 10.1177/1971400917697503. [Context Link 1] View abstract...
31. Shah SA, Saller J. Evaluation and diagnosis of back pain in children and adolescents. *Journal of the American Academy of Orthopedic Surgeons* 2016;24(1):37-45. DOI: 10.5435/JAAOS-D-14-00130. [Context Link 1] View abstract...
32. Huisman TAGM, et al. Advances in imaging modalities for pediatric brain and spinal cord tumors. *Pediatric Neurosurgery* 2023;58(5):240-258. DOI: 10.1159/000531998. [Context Link 1] View abstract...
33. Strauss SJ, et al. Bone sarcomas: ESMO-EURACAN-GENTURIS-ERNPaedCan Clinical Practice Guideline for diagnosis, treatment and follow-up. *Annals of Oncology* 2021;32(12):1520-1536. DOI: 10.1016/j.annonc.2021.08.1995. (Reaffirmed 2024 Aug) [Context Link 1] View abstract...
34. Rogers SN, et al. Imaging of pediatric spine and spinal cord tumors: a COG Diagnostic Imaging Committee/SPR Oncology Committee/ASPNR White Paper. *Pediatric Blood and Cancer* 2023;70 Suppl 4(Suppl 4):Online. DOI: 10.1002/pbc.30150. [Context Link 1, 2, 3] View abstract...
35. Tsukamoto S, Mavrogenis AF, Langevelde KV, Vucht NV, Kido A, Errani C. Imaging of spinal bone tumors: principles and practice. *Current Medical Imaging* 2022;18(2):142-161. DOI: 10.2174/1573405617666210301110446. [Context Link 1] View abstract...
36. Bestic JM, et al. Primary Bone Tumors. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2019 Accessed at: <https://www.acr.org>. [created 1995; accessed 2024 Jun 06] [Context Link 1, 2]
37. Go RS, et al. Histiocytic Neoplasms. NCCN Clinical Practice Guideline in Oncology [Internet] National Comprehensive Cancer Network (NCCN). v. 2.2024; 2024 Jul Accessed at: <https://www.nccn.org/>. [accessed 2024 Aug 16] [Context Link 1, 2, 3, 4]
38. Khoury JD, et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. *Leukemia* 2022;36(7):1703-1719. DOI: 10.1038/s41375-022-01613-1. [Context Link 1, 2] View abstract...
39. Emile JF, et al. Revised classification of histiocytoses and neoplasms of the macrophage-dendritic cell lineages. *Blood* 2016;127(22):2672-2681. DOI: 10.1182/blood-2016-01-690636. [Context Link 1] View abstract...
40. Haupt R, et al. Langerhans cell histiocytosis (LCH): guidelines for diagnosis, clinical work-up, and treatment for patients till the age of 18 years. *Pediatric Blood and Cancer* 2013;60(2):175-184. DOI: 10.1002/pbc.24367. [Context Link 1, 2] View abstract...
41. Goyal G, et al. The Mayo Clinic Histiocytosis Working Group consensus statement for the diagnosis and evaluation of adult patients with histiocytic neoplasms: Erdheim-Chester disease, Langerhans Cell Histiocytosis, and Rosai-Dorfman Disease. *Mayo Clinic Proceedings* 2019;94(10):2054-2071. DOI: 10.1016/j.mayocp.2019.02.023. [Context Link 1, 2] View abstract...
42. Pinter NK, Pfiffner TJ, Mechtler LL. Neuroimaging of spine tumors. *Handbook of Clinical Neurology* 2016;136:689-706. DOI: 10.1016/B978-0-444-53486-6.00033-8. [Context Link 1] View abstract...
43. Heindel W, Gubitz R, Vieth V, Weckesser M, Schober O, Schafers M. The diagnostic imaging of bone metastases. *Deutsches Arzteblatt International* 2014;111(44):741-7. DOI: 10.3238/arztebl.2014.0741. [Context Link 1] View abstract...
44. Yang HL, Liu T, Wang XM, Xu Y, Deng SM. Diagnosis of bone metastases: a meta-analysis comparing 18FDG PET, CT, MRI and bone scintigraphy. *European Radiology* 2011;21(12):2604-2617. DOI: 10.1007/s00330-011-2221-4. [Context Link 1] View abstract...
45. Ottenhausen M, et al. Intradural spinal tumors in adults-update on management and outcome. *Neurosurgical Review* 2019;42(2):371-388. DOI: 10.1007/s10143-018-0957-x. [Context Link 1] View abstract...
46. Moron FE, Delumpa A, Szklaruk J. Spinal tumors. In: Haaga JR, Boll DT, editors. *CT and MRI of the Whole Body*. 6th ed. Philadelphia, PA: Elsevier; 2017:844-874. [Context Link 1, 2, 3]
47. Ledbetter LN, Leever JD. Imaging of intraspinal tumors. *Radiologic Clinics of North America* 2019;57(2):341-357. DOI: 10.1016/j.rcl.2018.09.007. [Context Link 1] View abstract...
48. Missenard G, Bouthors C, Fadel E, Court C. Surgical strategies for primary malignant tumors of the thoracic and lumbar spine. *Orthopaedics & Traumatology, Surgery & Research : OTSR* 2020;106(1S):S53-S62. DOI: 10.1016/j.otsr.2019.05.028. [Context Link 1] View abstract...
49. Tettenborn B, Hagele-Link S. Spinal cord disorders. *European Neurology* 2015;74(3-4):141-146. DOI: 10.1159/000437417. [Context Link 1, 2] View abstract...
50. Lee MJ, Aronberg R, Manganaro MS, Ibrahim M, Parmar HA. Diagnostic approach to intrinsic abnormality of spinal cord signal intensity. *Radiographics* 2019;39(6):1824-1839. DOI: 10.1148/rg.2019190021. [Context Link 1, 2] View abstract...
51. Nabors LB, et al. Central Nervous System Cancers. NCCN Clinical Practice Guidelines in Oncology [Internet] National Comprehensive Cancer Network (NCCN). v. 2.2024; 2024 Jul Accessed at: <https://www.nccn.org/guidelines/>. [accessed 2024 Aug 16] [Context Link 1, 2]
52. Gajjar A, et al. Pediatric Central Nervous System Cancers. NCCN Clinical Practice Guideline in Oncology [Internet] National Comprehensive Cancer Network (NCCN). v. 1.2024; 2024 Feb 26 Accessed at: http://www.nccn.org/professionals/physician_gls/f_guidelines.asp. [accessed 2024 Mar 25] [Context Link 1]
53. Louis DN, et al. The 2021 WHO classification of tumors of the central nervous system: a summary. *Neuro-Oncology* 2021;23(8):1231-1251. DOI: 10.1093/neuonc/noab106. [Context Link 1] View abstract...
54. Fangusaro J, et al. Response assessment in paediatric low-grade glioma: recommendations from the Response Assessment in Pediatric Neuro-Oncology (RAPNO) working group. *Lancet Oncology* 2020;21(6):e305-e316. DOI: 10.1016/S1470-2045(20)30064-4. [Context Link 1, 2] View abstract...

55. Ivanidze J, et al. Brain Tumors. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2024 Accessed at: <https://www.acr.org>. [created 2024; accessed 2024 Nov 01] [Context Link 1, 2, 3]
56. Le Rhun E, et al. Leptomeningeal metastasis from solid tumours: EANO-ESMO Clinical Practice Guideline for diagnosis, treatment and follow-up. *ESMO Open Cancer Horizons* 2023;8(5):Online. DOI: 10.1016/j.esmoop.2023.101624. [Context Link 1] View abstract...
57. Evans DG. NF2-Related Schwannomatosis. Synonyms: Neurofibromatosis 2, Neurofibromatosis Type II [Internet] GeneReviews. 2023 Apr 20 Accessed at: <https://www.ncbi.nlm.nih.gov/books/NBK1201/>. [created 1998; accessed 2024 Sep 27] [Context Link 1]
58. Evans DGR, et al. Cancer and central nervous system tumor surveillance in pediatric neurofibromatosis 2 and related disorders. *Clinical Cancer Research* 2017;23(12):e54-e61. DOI: 10.1158/1078-0432.CCR-17-0590. [Context Link 1] View abstract...
59. Louise M Binderup M, et al. von Hippel-Lindau disease: Updated guideline for diagnosis and surveillance. *European Journal of Medical Genetics* 2022;65(8):Online. DOI: 10.1016/j.ejmg.2022.104538. [Context Link 1] View abstract...
60. Rednam SP, et al. Von Hippel-Lindau and hereditary pheochromocytoma/paraganglioma syndromes: clinical features, genetics, and surveillance recommendations in childhood. *Clinical Cancer Research* 2017;23(12):e68-e75. DOI: 10.1158/1078-0432.CCR-17-0547. [Context Link 1] View abstract...
61. Luetzen N, Dovi-Akue P, Fung C, Beck J, Urbach H. Spontaneous intracranial hypotension: diagnostic and therapeutic workup. *Neuroradiology* 2021;63(11):1765-1772. DOI: 10.1007/s00234-021-02766-z. [Context Link 1] View abstract...
62. Timpone VM, et al. Imaging of Suspected Intracranial Hypotension. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2024 Accessed at: <https://www.acr.org>. [created 2024; accessed 2024 Sep 26] [Context Link 1]
63. Wang LL, Bierbrauer KS. Congenital and hereditary diseases of the spinal cord. *Seminars in Ultrasound, CT and MRI* 2017;38(2):105-125. DOI: 10.1053/j.sult.2016.07.002. [Context Link 1] View abstract...
64. Sewell MJ, Chiu YE, Drolet BA. Neural tube dysraphism: review of cutaneous markers and imaging. *Pediatric Dermatology* 2015 Mar-Apr;32(2):161-170. DOI: 10.1111/pde.12485. [Context Link 1, 2, 3, 4] View abstract...
65. Shankar P, Zamora C, Castillo M. Congenital malformations of the brain and spine. *Handbook of Clinical Neurology* 2016;136:1121-1137. DOI: 10.1016/B978-0-444-53486-6.00058-2. [Context Link 1] View abstract...
66. Dias M, Partington M, Section on Neurologic Surgery. Congenital brain and spinal cord malformations and their associated cutaneous markers. *Pediatrics* 2015;136(4):e1105-e1119. DOI: 10.1542/peds.2015-2854. [Context Link 1, 2, 3, 4, 5] View abstract...
67. Gunny RS, Saunders DE, Argyropoulou MI. Paediatric neuroradiology. In: Adam A, Dixon AK, Gillard JH, Schaefer-Prokop CM, editors. *Grainger & Allison's Diagnostic Radiology*. 7th ed. Elsevier; 2021:1984-2045. [Context Link 1, 2, 3, 4, 5, 6]
68. Albano D, Messina C, Gitto S, Papakonstantinou O, Sconfienza LM. Differential diagnosis of spine tumors: my favorite mistake. *Seminars in Musculoskeletal Radiology* 2019;23(1):26-35. DOI: 10.1055/s-0038-1675551. [Context Link 1] View abstract...
69. Leake RL, Mills MK, Hanrahan CJ. Spinal marrow imaging: clues to disease. *Radiologic Clinics of North America* 2019;57(2):359-375. DOI: 10.1016/j.rcl.2018.09.008. [Context Link 1] View abstract...
70. Gaudino S, et al. A systematic approach to vertebral hemangioma. *Skeletal Radiology* 2015;44(1):25-36. DOI: 10.1007/s00256-014-2035-y. [Context Link 1] View abstract...
71. Peckham ME, Hutchins TA. Imaging of vascular disorders of the spine. *Radiologic Clinics of North America* 2019;57(2):307-318. DOI: 10.1016/j.rcl.2018.09.005. [Context Link 1] View abstract...
72. Ortiz AO, et al. Suspected Spine Infection. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2021 Accessed at: <https://www.acr.org>. [created 2021; accessed 2024 Oct 01] [Context Link 1]
73. Berbari EF, et al. 2015 Infectious Diseases Society of America (IDSA) clinical practice guidelines for the diagnosis and treatment of native vertebral osteomyelitis in adults. *Clinical Infectious Diseases* 2015;61(6):e26-e46. DOI: 10.1093/cid/civ482. (Reaffirmed 2024 Aug) [Context Link 1] View abstract...
74. Johnson SM, Shah LM. Imaging of acute low back pain. *Radiologic Clinics of North America* 2019;57(2):397-413. DOI: 10.1016/j.rcl.2018.10.001. [Context Link 1, 2] View abstract...
75. Le Saux N. Diagnosis and management of acute osteoarticular infections in children. *Paediatrics Child Health* 2018;23(5):336-343. DOI: 10.1093/pch/pxy049. [Context Link 1] View abstract...
76. Thurnher MM, Olatunji RB. Infections of the spine and spinal cord. *Handbook of Clinical Neurology* 2016;136:717-31. DOI: 10.1016/B978-0-444-53486-6.00035-1. [Context Link 1] View abstract...
77. Herren C, Jung N, Pishnamaz M, Breuninger M, Siewe J, Sobottke R. Spondylodiscitis: diagnosis and treatment options. *Deutsches Arzteblatt International* 2017;114(51-52):875-882. DOI: 10.3238/arztebl.2017.0875. [Context Link 1] View abstract...
78. Talbott JF, et al. Imaging-based approach to extradural infections of the spine. *Seminars in Ultrasound, CT and MRI* 2018;39(6):570-586. DOI: 10.1053/j.sult.2018.09.003. [Context Link 1] View abstract...
79. Thammaroj J, Kitkuandee A, Sawanyawisuth K. Differences of MRI features between tuberculous and bacterial spondylitis in a TB-endemic area. *Southeast Asian Journal of Tropical Medicine and Public Health* 2015;46(1):71-9. [Context Link 1] View abstract...
80. Talbott JF, Narvid J, Chazen JL, Chin CT, Shah V. An imaging-based approach to spinal cord infection. *Seminars in Ultrasound, CT and MRI* 2016;37(5):411-30. DOI: 10.1053/j.sult.2016.05.006. [Context Link 1] View abstract...
81. Kilborn T, van Rensburg PJ, Candy S. Pediatric and adult spinal tuberculosis: imaging and pathophysiology. *Neuroimaging Clinics of North America* 2015;25(2):209-231. DOI: 10.1016/j.nic.2015.01.002. [Context Link 1, 2, 3, 4] View abstract...
82. Raghavan M, Palestro CJ. Imaging of spondylodiscitis: an update. *Seminars in Nuclear Medicine* 2023;53(2):152-166. DOI: 10.1053/j.semnuclmed.2022.11.005. [Context Link 1] View abstract...

83. Babic M, Simpfordorfer CS. Infections of the spine. *Infectious Disease Clinics of North America* 2017;31(2):279-297. DOI: 10.1016/j.idc.2017.01.003. [Context Link 1, 2, 3] View abstract...
84. Lacasse M, et al. 2022 SPILF - clinical practice guidelines for the diagnosis and treatment of disco-vertebral infection in adults. *Infectious Diseases Now* 2023;53(3):Online. DOI: 10.1016/j.idnow.2023.01.007. [Context Link 1] View abstract...
85. Lener S, Hartmann S, Barbagallo GMV, Certo F, Thome C, Tschugg A. Management of spinal infection: a review of the literature. *Acta Neurochirurgica* 2018;160(3):487-496. DOI: 10.1007/s00701-018-3467-2. [Context Link 1] View abstract...
86. Achar S, Yamanaka J. Back pain in children and adolescents. *American Family Physician* 2020;102(1):19-28. [Context Link 1, 2, 3] View abstract...
87. Theodore N. Degenerative cervical spondylosis. *New England Journal of Medicine* 2020;383(2):159-168. DOI: 10.1056/NEJMr2003558. [Context Link 1, 2, 3] View abstract...
88. Colip CG, Lotfi M, Buch K, Holalkere N, Setty BN. Emergent spinal MRI in IVDU patients presenting with back pain: do we need an MRI in every case? *Emergency Radiology* 2018;25(3):247-256. DOI: 10.1007/s10140-017-1572-9. [Context Link 1] View abstract...
89. Laur O, Nandu H, Titelbaum DS, Nunez DB, Khurana B. Nontraumatic spinal cord compression: MRI primer for emergency department radiologists. *Radiographics* 2019;39(6):1862-1880. DOI: 10.1148/rg.2019190024. [Context Link 1, 2, 3, 4] View abstract...
90. Bart G, et al. Is there an association between magnetic resonance imaging and neurological signs in patients with vertebral osteomyelitis?: A retrospective observational study on 121 patients. *Medicine* 2016;95(3):e2373. DOI: 10.1097/MD.0000000000002373. [Context Link 1] View abstract...
91. Witte DH III. Advanced imaging in orthopaedics. In: Azar FM, Beatty JH, editors. *Campbell's Operative Orthopaedics*. 14th ed. Philadelphia, PA: Elsevier; 2021:141-176.e4. [Context Link 1, 2]
92. Huyskens J, et al. Overview of the complications and sequelae in spinal infections. *Neuroimaging Clinics of North America* 2015;25(2):309-21. DOI: 10.1016/j.nic.2015.01.007. [Context Link 1] View abstract...
93. Thwaites G, et al. British Infection Society guidelines for the diagnosis and treatment of tuberculosis of the central nervous system in adults and children. *Journal of Infection* 2009;59(3):167-187. DOI: 10.1016/j.jinf.2009.06.011. (Reaffirmed 2024 Jul) [Context Link 1] View abstract...
94. Rovira A, Auger C, Rovira A. Other noninfectious inflammatory disorders. *Handbook of Clinical Neurology* 2016;135:425-46. DOI: 10.1016/B978-0-444-53485-9.00021-0. [Context Link 1] View abstract...
95. Kranz PG, Amrhein TJ. Imaging approach to myelopathy: acute, subacute, and chronic. *Radiologic Clinics of North America* 2019;57(2):257-279. DOI: 10.1016/j.rcl.2018.09.006. [Context Link 1, 2] View abstract...
96. Wattjes MP, et al. 2021 MAGNIMS-CMSC-NAIMS consensus recommendations on the use of MRI in patients with multiple sclerosis. *Lancet Neurology* 2021;20(8):653-670. DOI: 10.1016/S1474-4422(21)00095-8. (Reaffirmed 2024 Sep) [Context Link 1, 2, 3] View abstract...
97. Barraza G, Deiva K, Husson B, Adamsbaum C. Imaging in pediatric multiple sclerosis : an iconographic review. *Clinical Neuroradiology* 2021;31(1):61-71. DOI: 10.1007/s00062-020-00929-8. [Context Link 1, 2] View abstract...
98. Filippi M, et al. Diagnosis of progressive multiple sclerosis from the imaging perspective: a review. *JAMA Neurology* 2021;78(3):351-364. DOI: 10.1001/jamaneurol.2020.4689. [Context Link 1] View abstract...
99. Fabian MT, Krieger SC, Lublin FD. Multiple sclerosis and other inflammatory demyelinating diseases of the central nervous system. In: Jankovic J, Mazziotta JC, Pomeroy SL, Newman NJ, editors. *Bradley and Daroff's Neurology in Clinical Practice*. 8th ed. Elsevier; 2022:1226-1254.e5. [Context Link 1, 2, 3]
100. McGinley MP, Goldschmidt CH, Rae-Grant AD. Diagnosis and treatment of multiple sclerosis: a review. *Journal of the American Medical Association* 2021;325(8):765-779. DOI: 10.1001/jama.2020.26858. [Context Link 1, 2] View abstract...
101. Bower A, Makhani N. Radiologically isolated syndrome and the multiple sclerosis prodrome in pediatrics: early features of the spectrum of demyelination. *Seminars in Pediatric Neurology* 2023;46:Online. DOI: 10.1016/j.spen.2023.101053. [Context Link 1] View abstract...
102. Brownlee WJ, Hardy TA, Fazekas F, Miller DH. Diagnosis of multiple sclerosis: progress and challenges. *Lancet* 2017;389(10076):1336-1346. DOI: 10.1016/S0140-6736(16)30959-X. [Context Link 1] View abstract...
103. Filippi M, Preziosa P, Rocca MA. Multiple sclerosis. *Handbook of Clinical Neurology* 2016;135:399-423. DOI: 10.1016/B978-0-444-53485-9.00020-9. [Context Link 1] View abstract...
104. Gass A, et al. MRI monitoring of pathological changes in the spinal cord in patients with multiple sclerosis. *Lancet Neurology* 2015;14(4):443-54. DOI: 10.1016/S1474-4422(14)70294-7. [Context Link 1] View abstract...
105. Muccilli A, Seyman E, Oh J. Spinal cord MRI in multiple sclerosis. *Neurologic Clinics* 2018;36(1):35-57. DOI: 10.1016/j.ncl.2017.08.009. [Context Link 1] View abstract...
106. Reich DS, Lucchinetti CF, Calabresi PA. Multiple sclerosis. *New England Journal of Medicine* 2018;378(2):169-180. DOI: 10.1056/NEJMr201401483. [Context Link 1] View abstract...
107. Jewells VL, Latchaw RE. What can mimic multiple sclerosis? *Seminars in Ultrasound, CT and MRI* 2020;41(3):284-295. DOI: 10.1053/j.sult.2020.02.003. [Context Link 1] View abstract...
108. Wendebourg MJ, Nagy S, Derfuss T, Parmar K, Schlaeager R. Magnetic resonance imaging in immune-mediated myelopathies. *Journal of Neurology* 2020;267(5):1233-1244. DOI: 10.1007/s00415-019-09206-2. [Context Link 1] View abstract...
109. Budd KR, Johnson M, Glaubiger SA, Jewells VL. Understanding the radiologist's role in evaluating multiple sclerosis: a review of the tip of the iceberg. *Seminars in Ultrasound, CT and MRI* 2020;41(3):275-283. DOI: 10.1053/j.sult.2020.02.002. [Context Link 1] View abstract...
110. Langille MM, Rutatangwa A, Francisco C. Pediatric multiple sclerosis: a review. *Advances in Pediatrics* 2019;66:209-229. DOI: 10.1016/j.yapd.2019.03.003. [Context Link 1] View abstract...

111. Hartung HP, Graf J, Aktas O, Mares J, Barnett MH. Diagnosis of multiple sclerosis: revisions of the McDonald criteria 2017 - continuity and change. *Current Opinion in Neurology* 2019;32(3):327-337. DOI: 10.1097/WCO.0000000000000699. [Context Link 1] View abstract...
112. Goh C, Desmond PM, Phal PM. MRI in transverse myelitis. *Journal of Magnetic Resonance Imaging : JMRI* 2014;40(6):1267-1279. DOI: 10.1002/jmri.24563. [Context Link 1, 2] View abstract...
113. Young V, Quaghebeur G. Transverse myelitis and neuromyelitis optica spectrum disorders. *Seminars in Ultrasound, CT and MRI* 2016;37(5):384-95. DOI: 10.1053/j.sult.2016.05.004. [Context Link 1] View abstract...
114. Dodson SC, Koontz NA. Spinal manifestations of systemic disease. *Radiologic Clinics of North America* 2019;57(2):281-306. DOI: 10.1016/j.rcl.2018.10.005. [Context Link 1] View abstract...
115. Clarencon F, Law-Ye B, Bienvenot P, Cormier E, Chiras J. The degenerative spine. *Magnetic Resonance Imaging Clinics of North America* 2016;24(3):495-513. DOI: 10.1016/j.mric.2016.04.008. [Context Link 1, 2] View abstract...
116. Shur N, Corrigan A, Agrawal K, Desai A, Gnanasegaran G. Radiological and radionuclide imaging of degenerative disease of the facet joints. *Indian Journal of Nuclear Medicine* 2015;30(3):191-198. DOI: 10.4103/0972-3919.158526. [Context Link 1, 2] View abstract...
117. Hegmann KT, et al. Diagnostic tests for low back disorders. *Journal of Occupational and Environmental Medicine* 2019;61(4):e155-e168. DOI: 10.1097/JOM.0000000000001551. [Context Link 1] View abstract...
118. Popescu A, Lee H. Neck pain and lower back pain. *Medical Clinics of North America* 2020;104(2):279-292. DOI: 10.1016/j.mcna.2019.11.003. [Context Link 1] View abstract...
119. Knezevic NN, Candido KD, Vlaeyen JWS, Van Zundert J, Cohen SP. Low back pain. *Lancet* 2021;398(10294):78-92. DOI: 10.1016/S0140-6736(21)00733-9. [Context Link 1] View abstract...
120. Hutchins TA, et al. Low Back Pain. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2021 Accessed at: <https://www.acr.org>. [created 1996; accessed 2024 Sep 26] [Context Link 1]
121. Hall AM, Aubrey-Bassler K, Thorne B, Maher CG. Do not routinely offer imaging for uncomplicated low back pain. *British Medical Journal* 2021;372:n291. DOI: 10.1136/bmj.n291. [Context Link 1] View abstract...
122. Ropper AH, Zafonte RD. Sciatica. *New England Journal of Medicine* 2015;372(13):1240-1248. DOI: 10.1056/NEJMr1410151. [Context Link 1] View abstract...
123. Patrick N, Emanski E, Knaub MA. Acute and chronic low back pain. *Medical Clinics of North America* 2016;100(1):169-181. DOI: 10.1016/j.mcna.2015.08.015. [Context Link 1] View abstract...
124. Hooten WM, Cohen SP. Evaluation and treatment of low back pain: A clinically focused review for primary care specialists. *Mayo Clinic Proceedings* 2015;90(12):1699-1718. DOI: 10.1016/j.mayocp.2015.10.009. [Context Link 1] View abstract...
125. Jones CMP, et al. Analgesia for non-specific low back pain. *British Medical Journal* 2024;385:Online. DOI: 10.1136/bmj-2024-080064. [Context Link 1, 2] View abstract...
126. Childress MA, Becker BA. Nonoperative management of cervical radiculopathy. *American Family Physician* 2016;93(9):746-754. [Context Link 1] View abstract...
127. Makin J, et al. Effectiveness and safety of manual therapy when compared with oral pain medications in patients with neck pain: a systematic review and meta-analysis. *BMC Sports Science, Medicine and Rehabilitation* 2024;16(1):Online. DOI: 10.1186/s13102-024-00874-w. [Context Link 1] View abstract...
128. Bellini M, Ferrara M, Grazzini I, Cerase A. Neuroimaging of the postoperative spine. *Magnetic Resonance Imaging Clinics of North America* 2016;24(3):601-620. DOI: 10.1016/j.mric.2016.04.006. [Context Link 1, 2] View abstract...
129. Pollice S, Muto M, Scarabino T. Post-therapeutic imaging findings. *European Journal of Radiology* 2015;84(5):799-806. DOI: 10.1016/j.ejrad.2014.06.010. [Context Link 1, 2] View abstract...
130. Willson MC, Ross JS. Postoperative spine complications. *Neuroimaging Clinics of North America* 2014;24(2):305-326. DOI: 10.1016/j.nic.2014.01.002. [Context Link 1, 2] View abstract...
131. Jones JY, et al. Scoliosis - Child. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2018 Accessed at: <https://www.acr.org>. [created 2018; accessed 2024 Jun 06] [Context Link 1, 2]
132. Hresko MT. Clinical practice. Idiopathic scoliosis in adolescents. *New England Journal of Medicine* 2013;368(9):834-841. DOI: 10.1056/NEJMc1209063. [Context Link 1, 2, 3] View abstract...
133. El-Hawary R, Chukwunyerenna C. Update on evaluation and treatment of scoliosis. *Pediatric Clinics of North America* 2014;61(6):1223-1241. DOI: 10.1016/j.pcl.2014.08.007. [Context Link 1] View abstract...
134. Pereira EAC, Oxenham M, Lam KS. Intraspinous anomalies in early-onset idiopathic scoliosis. *Bone and Joint Journal* 2017;99-B(6):829-833. DOI: 10.1302/0301-620X.99B6.BJJ-2016-1159.R1. [Context Link 1] View abstract...
135. Zhang W, Sha S, Xu L, Liu Z, Qiu Y, Zhu Z. The prevalence of intraspinal anomalies in infantile and juvenile patients with "presumed idiopathic" scoliosis: a MRI-based analysis of 504 patients. *BMC Musculoskeletal Disorders* 2016;17:189. DOI: 10.1186/s12891-016-1026-7. [Context Link 1, 2] View abstract...
136. Kim W, Porrino JA, Hood KA, Chadaz TS, Klauser AS, Taljanovic MS. Clinical evaluation, imaging, and management of adolescent idiopathic and adult degenerative scoliosis. *Current Problems in Diagnostic Radiology* 2019;48(4):402-414. DOI: 10.1067/j.cpradiol.2018.08.006. [Context Link 1, 2] View abstract...
137. Ng SY, Bettany-Saltikov J. Imaging in the diagnosis and monitoring of children with idiopathic scoliosis. *Open Orthopaedics Journal* 2017;11:1500-1520. DOI: 10.2174/1874325001711011500. [Context Link 1, 2] View abstract...
138. Barber JL, Offiah AC. Skeletal radiology in children: non-traumatic and non-malignant. In: Adam A, Dixon AK, Gillard JH, Schaefer-Prokop CM, editors. *Grainger & Allison's Diagnostic Radiology*. 7th ed. Elsevier; 2021:1886-1934. [Context Link 1, 2]

139. Agarwal V, et al. Myelopathy. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2020 Accessed at: <https://www.acr.org>. [created 1996; accessed 2024 Sep 27] [Context Link 1]
140. Murray EL, Misulis KE. Hemiplegia and monoplegia. In: Jankovic J, Mazziotta JC, Pomeroy SL, Newman NJ, editors. Bradley and Daroff's Neurology in Clinical Practice. 8th ed. Elsevier; 2022:345-355.e3. [Context Link 1]
141. Oyinkan Marquis B, Capone PM. Myelopathy. Handbook of Clinical Neurology 2016;136:1015-1026. DOI: 10.1016/B978-0-444-53486-6.00052-1. [Context Link 1] View abstract...
142. Lyerly MJ, Bag AK, Geldmacher DS. Spinal cord vascular disease. In: Jankovic J, Mazziotta JC, Pomeroy SL, Newman NJ, editors. Bradley and Daroff's Neurology in Clinical Practice. 8th ed. Elsevier; 2022:1058-1065.e1. [Context Link 1, 2, 3]
143. Pikija S, et al. Delayed hospital presentation and neuroimaging in non-surgical spinal cord infarction. Frontiers in Neurology 2017;8:Online. DOI: 10.3389/fneur.2017.00143. [Context Link 1, 2] View abstract...
144. Vargas MI, et al. Spinal cord ischemia: practical imaging tips, pearls, and pitfalls. AJNR. American Journal of Neuroradiology 2015;36(5):825-830. DOI: 10.3174/ajnr.A4118. [Context Link 1, 2] View abstract...
145. Thomas G, Alakbarzade V, Sammaraiye Y, Cociasu I, Dalton C, Pereira AC. Spontaneous spinal cord infarction: a practical approach. Practical Neurology 2022;22(6):497-502. DOI: 10.1136/pn-2022-003441. [Context Link 1] View abstract...
146. Khoueiry M, Moussa H, Sawaya R. Spinal cord infarction in a young adult: What is the culprit? Journal of Spinal Cord Medicine 2021;44(6):1015-1018. DOI: 10.1080/10790268.2019.1671077. [Context Link 1] View abstract...
147. Rock AK, Truong H, Park YL, Piliitsis JG. Spinal cord stimulation. Neurosurgery Clinics of North America 2019;30(2):169-194. DOI: 10.1016/j.nec.2018.12.003. [Context Link 1] View abstract...
148. Sitzman BT, Provenzano DA. Best practices in spinal cord stimulation. Spine 2017;42 Suppl 14:S67-S71. DOI: 10.1097/BRS.0000000000002220. [Context Link 1] View abstract...
149. Best BJ, Porwal MH, Pahapill PA. Preoperative thoracic spine magnetic resonance imaging for spinal cord stimulation: should such a recommendation be an absolute requirement? Neuromodulation 2022;25(5):758-762. DOI: 10.1111/ner.13518. [Context Link 1] View abstract...
150. Deer TR, et al. The Neurostimulation Appropriateness Consensus Committee (NACC) safety guidelines for the reduction of severe neurological injury. Neuromodulation 2017;20(1):15-30. DOI: 10.1111/ner.12564. [Context Link 1] View abstract...
151. Deer TR, et al. The Neurostimulation Appropriateness Consensus Committee (NACC): recommendations on best practices for cervical neurostimulation. Neuromodulation 2022;25(1):35-52. DOI: 10.1016/j.neurom.2021.10.013. [Context Link 1] View abstract...
152. Deer TR, et al. The Neurostimulation Appropriateness Consensus Committee (NACC): recommendations for surgical technique for spinal cord stimulation. Neuromodulation 2022;25(1):1-34. DOI: 10.1016/j.neurom.2021.10.015. [Context Link 1] View abstract...
153. Thompson DN. Spinal dysraphic anomalies; classification, presentation and management. Paediatrics and Child Health 2014;24(10):431-8. [Context Link 1, 2, 3]
154. Kumar J, Afsal M, Garg A. Imaging spectrum of spinal dysraphism on magnetic resonance: A pictorial review. World Journal of Radiology 2017;9(4):178-190. DOI: 10.4329/wjr.v9.i4.178. [Context Link 1, 2] View abstract...
155. Murphy CJ, Stanley E, Kavanagh E, Lenane PE, McCaul CL. Spinal dysraphisms in the parturient: implications for perioperative anaesthetic care and labour analgesia. International Journal of Obstetric Anesthesia 2015;24(3):252-63. DOI: 10.1016/j.ijoa.2015.04.002. [Context Link 1, 2] View abstract...
156. Nguyen Q, Tomei K. Spinal dysraphisms. In: Martin RJ, Fanaroff AA, editors. Fanaroff and Martin's Neonatal-Perinatal Medicine. 12th ed. Elsevier; 2025:1145-1153. [Context Link 1, 2]
157. Behbahani M, Lam SK, Bowman R. Cutaneous Stigmata of the Spine: A Review of Indications for Imaging and Referral. Pediatric Clinics of North America 2021;68(4):895-913. DOI: 10.1016/j.pcl.2021.04.017. [Context Link 1] View abstract...
158. White C, Milla SS, Maloney JA, Neuberger I. Imaging of congenital spine malformations. Clinics in Perinatology 2022;49(3):623-640. DOI: 10.1016/j.clp.2022.05.003. [Context Link 1] View abstract...
159. Martin BD, McClung A, Denning JR, Laine JC, Johnston CE. Intrathecal anomalies in presumed infantile idiopathic scoliosis: when is MRI necessary? Spine Deformity 2014;2(6):444-447. DOI: 10.1016/j.jspd.2014.03.003. [Context Link 1] View abstract...
160. Tuite GF, Thompson DNP, Austin PF, Bauer SB. Evaluation and management of tethered cord syndrome in occult spinal dysraphism: Recommendations from the international children's continence society. Neurourology and Urodynamics 2018;37(3):890-903. DOI: 10.1002/nau.23382. [Context Link 1, 2, 3] View abstract...
161. O'Neill BR, et al. Use of magnetic resonance imaging to detect occult spinal dysraphism in infants. Journal of Neurosurgery. Pediatrics 2017;19(2):217-226. DOI: 10.3171/2016.8.PEDS16128. [Context Link 1, 2] View abstract...
162. Kinsman SL, Johnston MV. Congenital anomalies of the central nervous system. In: Kliegman RM, St. Geme JW, Blum NJ, Shah SS, Tasker RC, Wilson KM, editors. Nelson Textbook of Pediatrics. 21st ed. Philadelphia, PA: Elsevier; 2020:3063-3082.e1. [Context Link 1]
163. Chen ZQ, Sun CG, Spine Surgery Group of Chinese Orthopedic Association. Clinical guideline for treatment of symptomatic thoracic spinal stenosis. Orthopaedic Surgery 2015;7(3):208-212. DOI: 10.1111/os.12190. [Context Link 1] View abstract...
164. Cowley P. Neuroimaging of spinal canal stenosis. Magnetic Resonance Imaging Clinics of North America 2016;24(3):523-539. DOI: 10.1016/j.mric.2016.04.009. [Context Link 1, 2] View abstract...
165. Sartip KA, et al. Ossification of the posterior longitudinal ligament: imaging findings in the era of cross-sectional imaging. Journal of Computer Assisted Tomography 2015;39(6):835-841. DOI: 10.1097/RCT.0000000000000303. [Context Link 1] View abstract...
166. Mizuno J. Radiologic evaluation of ossification of the posterior longitudinal ligament with dural ossification. Neurosurgery Clinics of North America 2018;29(1):55-61. DOI: 10.1016/j.nec.2017.09.007. [Context Link 1] View abstract...

167. Melancia JL, Francisco AF, Antunes JL. Spinal stenosis. *Handbook of Clinical Neurology* 2014;119:541-549. DOI: 10.1016/B978-0-7020-4086-3.00035-7. [Context Link 1] View abstract...
168. Li N, Amarasinghe S, Boudreaux K, Fakhre W, Sherman W, Kaye A. Spondylolysis. *Orthopedic Reviews* 2022;14(3):37470. DOI: 10.52965/001c.37470. [Context Link 1, 2] View abstract...
169. Mohile NV, Kuczmarski AS, Lee D, Warburton C, Rakoczy K, Butler AJ. Spondylolysis and isthmic spondylolisthesis: a guide to diagnosis and management. *Journal of the American Board of Family Medicine* 2022;35(6):1204-1216. DOI: 10.3122/jabfm.2022.220130R1. [Context Link 1, 2] View abstract...
170. Scholz M, et al. Recommendations for diagnosis and reatment of fractures of the ring of axis. *Zeitschrift fur Orthopadie und Unfallchirurgie* 2018;156(6):662-671. DOI: 10.1055/a-0620-9170. [Context Link 1, 2] View abstract...
171. Merrow AC, Hariharan S. Approach to pediatric spine. In: Merrow AC, Hariharan S, editors. *Imaging in Pediatrics*. Philadelphia, PA 19103-2899: Elsevier; 2018:328-329. [Context Link 1]
172. Redmond KJ, et al. Consensus guidelines for postoperative stereotactic body radiation therapy for spinal metastases: results of an international survey. *Journal of Neurosurgery: Spine* 2017;26(3):299-306. DOI: 10.3171/2016.8.SPINE16121. [Context Link 1, 2] View abstract...
173. ACR-ARS Practice Parameter for the Performance of Stereotactic Body Radiation Therapy. [Internet] American College of Radiology. 2024 Accessed at: <https://www.acr.org/>. [accessed 2024 May 31] [Context Link 1, 2]
174. Sahgal A, et al. The Canadian Association of Radiation Oncology scope of practice guidelines for lung, liver and spine stereotactic body radiotherapy. *Clinical Oncology* 2012;24(9):629-639. DOI: 10.1016/j.clon.2012.04.006. [Context Link 1] View abstract...
175. Thibault I, et al. Response assessment after stereotactic body radiotherapy for spinal metastasis: a report from the SPIne response assessment in Neuro-Oncology (SPINO) group. *Lancet Oncology* 2015;16(16):e595-e603. DOI: 10.1016/S1470-2045(15)00166-7. [Context Link 1, 2] View abstract...
176. Khan M, et al. Neuroimaging and stereotactic body radiation therapy (SBRT) for spine metastasis. *Topics in Magnetic Resonance Imaging: TMRI* 2019;28(2):85-96. DOI: 10.1097/RMR.000000000000199. [Context Link 1] View abstract...
177. Redmond KJ, et al. Consensus contouring guidelines for postoperative stereotactic body radiation therapy for metastatic solid tumor malignancies to the spine. *International Journal of Radiation Oncology, Biology and Physics* 2017;97(1):64-74. DOI: 10.1016/j.ijrobp.2016.09.014. [Context Link 1] View abstract...
178. Tseng CL, et al. Spine stereotactic body radiotherapy: indications, outcomes, and points of caution. *Global Spine Journal* 2017;7(2):179-97. DOI: 10.1177/2192568217694016. [Context Link 1] View abstract...
179. Poretti A, Boltshauser E, Huisman TA. Chiari malformations and syringohydromyelia in children. *Seminars in Ultrasound, CT and MRI* 2016;37(2):129-142. DOI: 10.1053/j.sult.2015.12.001. [Context Link 1] View abstract...
180. Rodriguez A, Kuhn EN, Somasundaram A, Couture DE. Management of idiopathic pediatric syringohydromyelia. *Journal of Neurosurgery. Pediatrics* 2015;16(4):452-457. DOI: 10.3171/2015.3.PEDS14433. [Context Link 1] View abstract...
181. McVige JW, Leonardo J. Imaging of Chiari type I malformation and syringohydromyelia. *Neurologic Clinics* 2014;32(1):95-126. DOI: 10.1016/j.ncl.2013.07.002. [Context Link 1, 2] View abstract...
182. Watts J, Box GA, Galvin A, Van Tonder F, Trost N, Sutherland T. Magnetic resonance imaging of intramedullary spinal cord lesions: a pictorial review. *Journal of Medical Imaging and Radiation Oncology* 2014;58(5):569-581. DOI: 10.1111/1754-9485.12202. [Context Link 1] View abstract...
183. Fehnel KP, Proctor MR. Spinal cord disorders. In: Kliegman RM, et al., editors. *Nelson Textbook of Pediatrics*. 22nd ed. Elsevier; 2025:3785-3790. [Context Link 1]
184. Roguski M, Groves ML. Adult tethered cord syndrome. In: Winn RH, editor. *Youmans and Winn Neurological Surgery*. 8th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:2594-2600.e2. [Context Link 1]
185. Samuk I, Bischoff A, Freud E, Pena A. Tethered cord in children with anorectal malformations with emphasis on rectobladder neck fistula. *Pediatric Surgery International* 2019;35(2):221-226. DOI: 10.1007/s00383-018-4399-x. [Context Link 1, 2] View abstract...
186. van den Hondel D, Sloots C, de Jong TH, Lequin M, Wijnen R. Screening and treatment of tethered spinal cord in anorectal malformation patients. *European Journal of Pediatric Surgery* 2016;26(1):22-8. DOI: 10.1055/s-0035-1563673. [Context Link 1] View abstract...
187. Tamura G, Morota N, Ihara S. Impact of magnetic resonance imaging and urodynamic studies on the management of sacrococcygeal dimples. *Journal of Neurosurgery. Pediatrics* 2017;20(3):289-297. DOI: 10.3171/2017.5.PEDS16719. [Context Link 1] View abstract...
188. Wolf M, Weber MA. Neuroimaging of the traumatic spine. *Magnetic Resonance Imaging Clinics of North America* 2016;24(3):541-561. DOI: 10.1016/j.mric.2016.04.004. [Context Link 1] View abstract...
189. Guarnieri G, Izzo R, Muto M. The role of emergency radiology in spinal trauma. *British Journal of Radiology* 2016;89(1061):20150833. DOI: 10.1259/bjr.20150833. [Context Link 1] View abstract...
190. Sixta S, et al. Screening for thoracolumbar spinal injuries in blunt trauma: an Eastern Association for the Surgery of Trauma practice management guideline. *Journal of Trauma and Acute Care Surgery* 2012;73(5 Suppl 4):S326-S332. DOI: 10.1097/TA.0b013e31827559b8. (Reaffirmed 2024 Aug) [Context Link 1] View abstract...
191. BOAST- Cervical Spine Clearance in the Trauma Patient. [Internet] British Orthopaedic Association. 2021 May Accessed at: <https://www.boa.ac.uk/>. [accessed 2024 Sep 20] [Context Link 1]
192. Masson de Almeida Prado R, Masson de Almeida Prado JL, Ueta RHS, Guimaraes JB, Yamada AF. Subaxial spine trauma: radiological approach and practical implications. *Clinical Radiology* 2021;76(12):941.e1-941.e10. DOI: 10.1016/j.crad.2021.09.006. [Context Link 1] View abstract...
193. Boese CK, Lechler P. Spinal cord injury without radiologic abnormalities in adults: a systematic review. *Journal of Trauma and Acute Care Surgery* 2013;75(2):320-330. DOI: 10.1097/TA.0b013e31829243c9. [Context Link 1, 2] View abstract...

194. Boese CK, Oppermann J, Siewe J, Eysel P, Scheyerer MJ, Lechler P. Spinal cord injury without radiologic abnormality in children: a systematic review and meta-analysis. *Journal of Trauma and Acute Care Surgery* 2015;78(4):874-882. DOI: 10.1097/TA.0000000000000579. [Context Link 1, 2] View abstract...
195. Knox JB, Schneider JE, Cage JM, Wimberly RL, Riccio AI. Spine trauma in very young children: a retrospective study of 206 patients presenting to a level 1 pediatric trauma center. *Journal of Pediatric Orthopedics* 2014;34(7):698-702. DOI: 10.1097/BPO.0000000000000167. [Context Link 1, 2] View abstract...
196. Abdel-Wanis ME, Solyman MT, Hasan NM. Sensitivity, specificity and accuracy of magnetic resonance imaging for differentiating vertebral compression fractures caused by malignancy, osteoporosis, and infections. *Journal of Orthopaedic Surgery (Hong Kong)* 2011;19(2):145-150. [Context Link 1] View abstract...
197. Shah LM, Flanders AE. Update on new imaging techniques for trauma. *Neurosurgery Clinics of North America* 2017;28(1):1-21. DOI: 10.1016/j.nec.2016.08.006. [Context Link 1] View abstract...
198. Kumar Y, Hayashi D. Role of magnetic resonance imaging in acute spinal trauma: a pictorial review. *BMC Musculoskeletal Disorders* 2016;17:310. DOI: 10.1186/s12891-016-1169-6. [Context Link 1] View abstract...
199. Pierce JL, et al. Spinal hematomas: what a radiologist needs to know. *Radiographics* 2018;38(5):1516-1535. DOI: 10.1148/rg.2018180099. [Context Link 1, 2] View abstract...
200. Takahashi S, et al. Predicting delayed union in osteoporotic vertebral fractures with consecutive magnetic resonance imaging in the acute phase: a multicenter cohort study. *Osteoporosis International* 2016;27(12):3567-3575. DOI: 10.1007/s00198-016-3687-3. [Context Link 1] View abstract...
201. Qureshi S, et al. Congress of Neurological Surgeons systematic review and evidence-based guidelines on the evaluation and treatment of patients with thoracolumbar spine trauma: radiological evaluation. *Neurosurgery* 2019;84(1):E28-E31. DOI: 10.1093/neuros/nyy373. [Context Link 1, 2] View abstract...
202. Tsoumakidou G, et al. CIRSE guidelines on percutaneous vertebral augmentation. *Cardiovascular and Interventional Radiology* 2017;40(3):331-42. DOI: 10.1007/s00270-017-1574-8. (Reaffirmed 2024 Aug) [Context Link 1, 2, 3] View abstract...
203. Moriarty HK, O Cearbhaill R, Moriarty PD, Stanley E, Lawler LP, Kavanagh EC. MR imaging of spinal haematoma: a pictorial review. *British Journal of Radiology* 2019;92(1095):20180532. DOI: 10.1259/bjr.20180532. [Context Link 1] View abstract...
204. Spinal Injury: Assessment and Initial Management. NICE Guideline NG41 [Internet] National Institute for Health and Care Excellence. 2016 Feb Accessed at: <https://www.nice.org.uk/guidance/>. [accessed 2024 Oct 01] [Context Link 1, 2] View abstract...
205. Jo AS, Wilseck Z, Manganaro MS, Ibrahim M. Essentials of spine trauma imaging: radiographs, CT, and MRI. *Seminars in Ultrasound, CT and MRI* 2018;39(6):532-550. DOI: 10.1053/j.sult.2018.10.002. [Context Link 1, 2, 3] View abstract...
206. Vincent SA, Anderson PA. The unstable spine: a surgeon's perspective. *Seminars in Ultrasound, CT and MRI* 2018;39(6):618-629. DOI: 10.1053/j.sult.2018.10.001. [Context Link 1, 2] View abstract...
207. Bernstein MP, Young MG, Baxter AB. Imaging of spine trauma. *Radiologic Clinics of North America* 2019;57(4):767-785. DOI: 10.1016/j.rcl.2019.02.007. [Context Link 1] View abstract...
208. Hassankhani A, et al. Acute Spinal Trauma. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2024 Accessed at: <https://www.acr.org>. [created 2024; accessed 2024 Nov 01] [Context Link 1]
209. Kyriakou C, et al. The role of cement augmentation with percutaneous vertebroplasty and balloon kyphoplasty for the treatment of vertebral compression fractures in multiple myeloma: a consensus statement from the International Myeloma Working Group (IMWG). *Blood Cancer Journal* 2019;9(3):Online. DOI: 10.1038/s41408-019-0187-7. [Context Link 1] View abstract...
210. Khan MA, et al. Management of Vertebral Compression Fractures. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2022 Accessed at: <https://www.acr.org>. [created 2010; accessed 2024 Sep 26] [Context Link 1]
211. Katsuura Y, Osborn JM, Cason GW. The epidemiology of thoracolumbar trauma: A meta-analysis. *Journal of Orthopaedics* 2016;13(4):383-8. DOI: 10.1016/j.jor.2016.06.019. [Context Link 1] View abstract...
212. Izzo R, Popolizio T, Balzano RF, Pennelli AM, Simeone A, Muto M. Imaging of cervical spine traumas. *European Journal of Radiology* 2019;117:75-88. DOI: 10.1016/j.ejrad.2019.05.007. [Context Link 1] View abstract...
213. Sjeklocha L, Gatz JD. Traumatic injuries to the spinal cord and peripheral nervous system. *Emergency Medicine Clinics of North America* 2021;39(1):1-28. DOI: 10.1016/j.emc.2020.09.001. [Context Link 1] View abstract...
214. Hagan MJ, et al. Spinal cord injury in adult and pediatric populations. *Interdisciplinary Neurosurgery : Advanced techniques and Case Management* 2022;29:Online. DOI: 10.1016/j.inat.2022.101594. [Context Link 1]
215. Zohrabian VM, Flanders AE. Imaging of trauma of the spine. *Handbook of Clinical Neurology* 2016;136:747-67. DOI: 10.1016/B978-0-444-53486-6.00037-5. [Context Link 1, 2] View abstract...
216. de Almeida Prado RM, et al. Spine trauma: Radiological approach and new concepts. *Skeletal Radiology* 2021;50(6):1065-1079. DOI: 10.1007/s00256-020-03668-6. [Context Link 1] View abstract...
217. Srinivasan V, Jea A. Pediatric thoracolumbar spine trauma. *Neurosurgery Clinics of North America* 2017;28(1):103-114. DOI: 10.1016/j.nec.2016.07.003. [Context Link 1] View abstract...
218. Hirsch JA, et al. Management of vertebral fragility fractures: a clinical care pathway developed by a multispecialty panel using the RAND/UCLA Appropriateness Method. *Spine Journal* 2018;18(11):2152-2161. DOI: 10.1016/j.spinee.2018.07.025. [Context Link 1] View abstract...
219. Kemp A, Cowley L, Maguire S. Spinal injuries in abusive head trauma: patterns and recommendations. *Pediatric Radiology* 2014;44 Suppl 4:S604-S612. DOI: 10.1007/s00247-014-3066-1. [Context Link 1] View abstract...

220. Palasis S, Hayes LL. Acquired pathology of the pediatric spine and spinal cord. *Pediatric Radiology* 2015;45 Suppl 3:S420-S432. DOI: 10.1007/s00247-015-3328-6. [Context Link 1] View abstract...

221. Kadom N, et al. Suspected Spine Trauma - Child. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2018 Accessed at: <https://www.acr.org/>. [created 2018; accessed 2024 Oct 01] [Context Link 1]

222. Traylor KS, Kralik SF, Radhakrishnan R. Pediatric spine emergencies. *Seminars in Ultrasound, CT and MRI* 2018;39(6):605-617. DOI: 10.1053/j.sult.2018.09.002. [Context Link 1] View abstract...

223. Odle TG. Computed tomography of thoracolumbar spine trauma. *Radiologic Technology* 2017;88(3):299-319. [Context Link 1] View abstract...

224. Bouthors C, Benzakour A, Court C. Surgical treatment of thoracic disc herniation: an overview. *International Orthopaedics* 2019;43(4):807-816. DOI: 10.1007/s00264-018-4224-0. [Context Link 1] View abstract...

225. Chaudhary V, Bano S, Garga UC. Central nervous system tuberculosis: an imaging perspective. *Canadian Association of Radiologists Journal* 2017;68(2):161-170. DOI: 10.1016/j.carj.2016.10.007. [Context Link 1] View abstract...

226. Prodi E, Grassi R, Iacobellis F, Cianfoni A. Imaging in spondylodiskitis. *Magnetic Resonance Imaging Clinics of North America* 2016;24(3):581-600. DOI: 10.1016/j.mric.2016.04.005. [Context Link 1] View abstract...

227. Hirji H, Saifuddin A. Paediatric acquired pathological vertebral collapse. *Skeletal Radiology* 2014;43(4):423-436. DOI: 10.1007/s00256-013-1792-3. [Context Link 1] View abstract...

228. Torres C, Riascos R, Figueroa R, Gupta RK. Central nervous system tuberculosis. *Topics in Magnetic Resonance Imaging: TMRI* 2014;23(3):173-189. DOI: 10.1097/RMR.000000000000023. [Context Link 1, 2] View abstract...

229. Schaller MA, Wicke F, Foerch C, Weidauer S. Central nervous system tuberculosis : etiology, clinical manifestations and neuroradiological features. *Clinical Neuroradiology* 2019;29(1):3-18. DOI: 10.1007/s00062-018-0726-9. [Context Link 1, 2] View abstract...

230. Centers for Medicare and Medicaid Services. "Hospital services excluded from payment under the hospital outpatient prospective payment system." 42 CFR 419.22 Washington, DC 2023 Jul [accessed 2024 Oct 08] Accessed at: <https://www.ecfr.gov/>. [Context Link 1]

231. Centers for Medicare and Medicaid Services. "Requirements relating to basic benefits." 42 CFR 422.101 Washington, DC 2024 Sep 30 [accessed 2024 Oct 08] Accessed at: <https://www.ecfr.gov/>. [Context Link 1]

232. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 14 - Medical Devices Rev. 198 [Internet] Centers for Medicare and Medicaid Services. 2014 Nov 06 Accessed at: <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/>. [accessed 2024 Sep 26] [Context Link 1]

233. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 15 - Covered Medical and Other Health Services Rev. 12425 [Internet] Centers for Medicare & Medicaid Services. 2023 Dec 21 Accessed at: <https://www.cms.gov/manuals/>. [accessed 2024 Feb 01] [Context Link 1]

234. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 16 - General Exclusions From Coverage Rev. 198 [Internet] Centers for Medicare and Medicaid Services. 2014 Nov 06 Accessed at: <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/>. [accessed 2024 Jun 06] [Context Link 1]

235. Medicare Coverage Database. [Internet] Centers for Medicare and Medicaid Services. Accessed at: <https://www.cms.gov/medicare-coverage-database/search.aspx? Updated 2024> [accessed 2024 Sep 26] [Context Link 1]

Footnotes

[A] Histiocytic and dendritic cell neoplasms are a heterogeneous group of rare hematologic disorders that arise from neoplastic proliferations of cells in the monocytic, histiocytic, or dendritic lineage that arise from common myeloid progenitors.(37)(38) They are classified on the basis of clinical, radiographic, histologic, and molecular features. Major categories include the plasmacytoid dendritic cell neoplasms; Langerhans cell and other dendritic cell neoplasms, including Langerhans cell histiocytosis; and histiocytic neoplasms, including Erdheim-Chester disease and Rosai-Dorfman disease.(37)(38)(39) Depending on the subtype, histiocytic and dendritic cell neoplasms may be characterized by vertebral lesions.(37)(40)(41) [A in Context Link 1]

Codes

CPT®: 72146, 72147, 72157

CPT copyright 2024 American Medical Association. All rights reserved.

MCG Health
Ambulatory Care 29th Edition
Copyright © 2025 MCG Health, LLC
All Rights Reserved

Last Update: 6/13/2025 2:11:20 AM
Build Number: 29.1.20250613001137.022065