

Lumbar Fusion

ORG: S-820 (ISC)
Link to Codes

- Care Planning - Inpatient Admission and Alternatives
 - Clinical Indications for Procedure
 - Alternatives to Procedure
 - Operative Status Criteria
- Hospitalization
 - Optimal Recovery Course
 - Goal Length of Stay - **Ambulatory**
 - Extended Stay
- Discharge
 - Discharge Planning
 - Discharge Destination
- Evidence Summary
 - Background
 - Criteria
 - Alternatives
 - Length of Stay
 - Rationale
 - Related CMS Coverage Guidance
- References
- Footnotes
- Definitions
- Codes

Care Planning - Inpatient Admission and Alternatives

Clinical Indications for Procedure

- Procedure is indicated for **1 or more** of the following(1):**NNNNNN**
 - ☐ Spinal fracture repair, as indicated by **1 or more** of the following(4)(5)(12):
 - Spinal instability (eg, burst fracture)
 - Neural compression
 - ☐ Spinal stenosis surgery requiring stabilization with fusion, as indicated by **ALL** of the following(1)(6)(13):
 - Unacceptable postoperative instability is judged to be likely due to extent of disease or surgery (eg, multiple vertebral levels involved).
 - Lumbar spinal stenosis treatment needed, as indicated by **1 or more** of the following:
 - Rapidly progressive or very severe symptoms of neurogenic claudication with imaging findings of lumbar spinal stenosis that correlate to clinical findings(14)
 - Leg or buttock neurogenic claudication symptoms and **ALL** of the following:
 - Symptoms that are persistent and disabling
 - Imaging findings of lumbar spinal stenosis that correlate with clinical findings
 - Failure of 3 months of nonoperative therapy
 - ☐ Lumbar spondylolisthesis, as indicated by **1 or more** of the following(1)(7)(15)(16):
 - Rapidly progressive or very severe neurologic deficits (eg, bowel or bladder dysfunction)
 - Symptoms requiring treatment, as indicated by **ALL** of the following:
 - Patient has persistent disabling symptoms, including **1 or more** of the following:
 - Low back pain
 - Neurogenic claudication
 - Radicular pain
 - Treatment is indicated by **ALL** of the following:
 - Listhesis demonstrated on imaging
 - Symptoms correlate with findings on MRI or other imaging.
 - Failure of 3 months of nonoperative therapy
 - Child or adolescent with high-grade (greater than 50% slippage) spondylolisthesis (to prevent progression)(17)(18)(19)

- ☐ Adjacent segment disease, as indicated by **ALL** of the following(1)(8):
 - Radiographic evidence of adjacent segment disease (eg, neural compression) that correlates with symptoms
 - Persistent disabling symptoms (low back pain, radiculopathy)
 - Failure of 3 months of nonoperative therapy
- ☐ Lumbar pseudarthrosis and **ALL** of the following(1)(9):
 - Symptoms (eg, pain) unresponsive to nonoperative therapy
 - Lumbar imaging findings consistent with symptoms
 - Alternative etiologies of symptoms ruled out
- Spinal repair with fusion (eg, for instability due to extensive surgery) in conjunction with other procedures (eg, laminectomy) for neural decompression, fracture, dislocation, infection, abscess, tumor, or coronal imbalance(10)(11)(20)(21)

Alternatives to Procedure

- Alternatives include(1)(7)(12)(13)(22):[\[1\]](#)
 - Nonoperative measures, including(17)(25):
 - Anti-inflammatory medication
 - Analgesics
 - Lumbar bracing
 - Physical therapy
 - Activity modification
 - Laminectomy without fusion. See Lumbar Laminectomy [\[2\]](#) ISC guideline.
 - Discectomy without fusion. See Lumbar Discectomy, Foraminotomy, or Laminotomy [\[3\]](#) ISC guideline.
 - Hemilaminectomy
 - Multilevel unilateral or bilateral laminotomy
 - Fracture fixation without fusion(5)
 - Radiation therapy, chemotherapy, and glucocorticoids (eg, neoplastic epidural compression)(21)

Operative Status Criteria

Goal Length of Stay: Ambulatory

Note: The definition of an ambulatory procedure depends on payer-provider contractual agreement or regulatory language (eg, CMS' Two-Midnight Rule). An ambulatory procedure may include one postoperative overnight stay in a facility; therefore, MCG's ambulatory Goal Length of Stay (GLOS) attainment calculation reports the sum of same-day and next-day postoperative discharges. Depending on various patient and procedural factors, some patients undergoing a procedure with an ambulatory GLOS require inpatient care (eg, medical necessity for hospital-based care across 2 or more postoperative midnights). Some of these factors are described in the Extended Stay section of this guideline.

- Ambulatory
- Inpatient (eg, medical necessity for hospital-based care across 2 or more postoperative midnights)
- Inpatient (Medicare patient, and specific procedure is on CMS Inpatient Only List)

Hospitalization

Optimal Recovery Course

Day	Level of Care	Clinical Status	Activity	Routes	Interventions	Medications
1	• Social Determinants of Health Assessment • OR to recovery to discharge[A] • Discharge planning	• Hemodynamic stability • No new neurologic deficit • Pain absent or managed • No evidence of postoperative or surgical site infection • Voiding ability at baseline • Adequate ambulation • Discharge plans and education	• Ambulatory or acceptable for next level of care[B]	• Oral hydration[C] • Oral medications or regimen acceptable for next level of care • Oral diet or acceptable for next level of care	• DVT prophylaxis[D]	• Prophylactic antibiotics[E] • PCA and parenteral analgesics absent[F]

Recovery Milestones are indicated in **bold**.

Goal Length of Stay: Ambulatory

Note: Goal Length of Stay assumes optimal recovery, decision making, and care. Patients may be discharged to a lower level of care (either later than or sooner than the goal) when it is appropriate for their clinical status and care needs.

Extended Stay

Minimal (a few hours to 1 day), Brief (1 to 3 days), Moderate (4 to 7 days), and Prolonged (more than 7 days).

- Inpatient stay (eg, need for hospital-based care beyond postoperative day 1) may be needed for(29):
 - Failure to meet discharge criteria (recovery milestones within final day in Optimal Recovery Course)
 - Expect brief stay extension.
 - Complications of surgery (eg, prolonged ileus, respiratory insufficiency)(30)
 - Expect brief stay extension.
 - Pathologic fracture or neoplasm
 - Expect brief stay extension.
 - Spinal abscess or osteomyelitis
 - Expect brief to moderate stay extension.
 - Severe preoperative deficit or injury
 - Patient with significant neurologic compromise or multiple injuries will require longer acute care and recovery times.
 - Stay extension varies depending on injury.
 - Dural tear(31)
 - Anticipate possible CSF drainage and surgical repair.
 - Expect brief to moderate stay extension.
 - Intraspinous hemorrhage
 - Intraspinous hemorrhage may need surgical repair.
 - Expect brief to moderate stay extension.
 - Extraspinous hemorrhage
 - Anticipate possible evacuation and surgical repair.
 - Expect brief to moderate stay extension.
 - Nerve root or cauda equina injury
 - Anticipate possible physical therapy and rehabilitation.
 - Expect brief to moderate stay extension.
 - Visual loss (eg, ischemic optic retinopathy)
 - Expect brief to moderate stay extension.
 - Combined (anterior and posterior) procedures
 - Expect brief stay extension.
 - Active comorbid illness (eg, cardiovascular, renal, or pulmonary disease)(32)
 - Anticipate relevant monitoring, treatment, and consultation.
 - Expect brief stay extension.
 - Postoperative altered mental status (eg, delirium)
 - Anticipate evaluation for etiology (eg, electrolyte abnormalities, pain, medications) and specific management as appropriate.
 - Expect brief stay extension.

See Common Complications and Conditions [ISC](#) for further information.

Discharge

Discharge Planning

- Discharge planning includes[G]:
 - Assessment of needs and planning for care, including(34)(35):
 - Develop and modify treatment plan (involving multiple providers) as needed.
 - Evaluate and address preadmission functioning as needed.

- Evaluate and address psychosocial status issues as indicated. See Psychosocial Assessment [SR](#) for further information.
- Evaluate and address social determinants of health (eg, housing, food). See Social Determinants of Health Screening Tool [SR](#) for further information.(33)
- Evaluate and address patient or caregiver preferences as indicated.
- Identify skilled services needed at next level of care, with specific attention to:
 - Medication management, adherence instruction, and side effects assessment(36)
 - Neurologic status assessment(37)
 - Pain management(37)
 - Rehabilitation therapy or equipment coordination(38)
 - Wound or dressing management(39)
- Early identification of anticipated discharge destination; options include(35)(40):
 - Home; considerations include:
 - Home safety assessment. See Home Safety Assessment [SR](#) for further information.
 - ☐ Patient safe to go home; examples include(41)(42)(43):
 - Medical status stable for patient's condition
 - Functional care can safely be provided with available resources.
 - Mental status stable for patient's condition
 - Medication availability confirmed and reconciliation complete
 - Patient/caregiver education completed with written discharge instructions provided
 - Community resources identified and referrals made, as needed
 - Home care arranged, if indicated
 - Necessary medical equipment delivery arranged or available in home, if indicated
 - Necessary medical supplies ordered, or patient/caregiver can obtain, if indicated
 - Access to follow-up care
 - Self-management ability if appropriate. See Activities of Daily Living (ADL) and Instrumental Activities of Daily Living (IADL) Assessment [SR](#) for further information.
 - Caregiver need, ability, and availability
 - Post-acute skilled care or custodial care as indicated. See Discharge Planning Tool [SR](#) for further information.
- Transitions of care plan complete, including(35)(40)(44):
 - Patient and caregiver education complete.
 - See Teach Back Tool [SR](#) for further information.
 - See Lumbar Fusion: Patient Education for Clinicians [SR](#) for further information.
 - ☐ Medication reconciliation completion includes(36)(45):
 - Compare patient's discharge list of medications (prescribed and over-the-counter) against provider's admission or transfer orders.
 - Assess each medication for correlation to disease state or medical condition.
 - Report medication discrepancies to prescribing provider, attending physician, and primary care provider, and ensure accurate medication order is identified.
 - Provide reconciled medication list to all treating providers.
 - Confirm that patient or caregiver can acquire medication.
 - Educate patient and caregiver.
 - Provide complete medication list to patient and caregiver.
 - Importance of presenting personal medication list to all providers at each care transition, including all provider appointments
 - Reason, dosage, and timing of medication (eg, use "teach-back" techniques)(46)
 - Encourage communication between patient, caregiver, and pharmacy for obtaining prescriptions, setting up home medication delivery, and reviewing for drug-drug interactions.
 - See Medication Reconciliation Tool [SR](#) for further information.
 - Plan communicated to patient, caregiver, and all members of care team, including(47):
 - Inpatient care and service providers
 - Primary care provider
 - All post-discharge care and service providers
 - Appointments planned or scheduled, which may include(1):
 - Primary care provider
 - Neurosurgeon(48)
 - Orthopedic surgeon
 - Rehabilitation therapy services
 - Specialists for management of comorbidities as needed
 - Other

- Outpatient testing and procedure plans made, which may include:
 - Laboratory testing
 - Other
- Referrals made for assistance or support, which may include:
 - Financial, for follow-up care, medication, and transportation(49)
 - Tobacco use treatment(50)
 - Vocational rehabilitation(37)
 - Other
- Medical equipment and supplies coordinated (ie, delivered or delivery confirmed), which may include:
 - Ambulation devices (eg, cane, crutches, walker)(51)
 - Antiembolic or compression stockings(51)
 - Immobilizers (eg, braces, splints)(52)
 - Wound care equipment and supplies(53)
 - Other

Discharge Destination

- Post-hospital levels of admission may include:
 - Home.
 - Home healthcare. See Home Care Indications for Admission Section [HC](#) in Lumbar Spine Surgery guideline in Home Care.
 - Recovery facility care. See Recovery Facility Care Indications for Admission Section [RFC](#) in Lumbar Spine Surgery guideline in Recovery Facility Care.

Evidence Summary

Background

Lumbar fusion may be performed either directly through an open incision, via image-guided minimally invasive surgery, or by endoscopic surgery, based on patient characteristics and surgeon experience.(2)(3) **(EG 2)**

Criteria

The evidence for the clinical indications found in this guideline includes 16 published peer reviewed articles and 5 book sections.

For spinal fracture repair, a narrative review states that lumbosacral fusion is indicated for patients with unstable sacral fractures.(4) **(EG 2)** An orthopedic surgery textbook states that lumbar fusion surgery is appropriate for patients with spinal fractures associated with neural compression or instability.(5) **(EG 2)**

For spinal stenosis surgery, an orthopedic surgery textbook notes that additional lumbar fusion may be needed if excessive bony resection occurs which destabilizes the spine.(1) **(EG 2)** A narrative review states that interbody fusion of the lumbar spine can be used in the setting of spinal stenosis to provide indirect decompression and improve symptoms.(6) **(EG 2)**

For lumbar spondylolisthesis, an orthopedic surgery textbook states that lumbar fusion is an appropriate treatment for spondylolisthesis of all types when patients have chronic pain from instability.(1)(7) **(EG 2)** Narrative reviews note that lumbar fusion is appropriate for patients with high-grade spondylolisthesis.(4)(8) **(EG 2)**

For adjacent segment disease, a narrative review notes that lateral lumbar fusion can be a helpful procedure to improve symptoms.(8) **(EG 2)**

For lumbar pseudarthrosis, an orthopedic surgery textbook and a narrative review state that repeat lumbar fusion is a recommended procedure to improve symptoms in patients with localized pain over the fusion area and when lumbar imaging findings are consistent with the symptoms.(1)(9) **(EG 2)**

For spinal repair with fusion in conjunction with other procedures, an orthopedic surgery textbook states that fusion can be a helpful adjunctive procedure when any significant amount of bone is resected due to tumor treatment.(10) **(EG 2)** A narrative review states that surgical management of spinal tuberculosis includes debridement, decompression, and adequate fusion to provide for early mobilization.(11) **(EG 2)**

Alternatives

A randomized controlled study of 233 patients (mean age 67 years) undergoing surgery for lumbar spinal stenosis (at 1 or 2 levels) found that the addition of fusion to laminectomy did not result in improved clinical outcomes at 2-year and 5-year follow-up, and that fusion resulted in longer operative times and more blood loss.(23) **(EG 1)** In the same trial, when patients were stratified by the presence or absence of spondylolisthesis, the results were similar.(23) **(EG 1)** A study of 211 patients (mean age 66 years) with lumbar spinal stenosis randomized to 1-level or 2-level decompression alone or decompression with fusion found that, although spinal

restenosis was less frequent after decompression with fusion at the operative level (4% vs 14% for decompression with fusion vs decompression alone, respectively), this was more than offset by an increase in the rate of new stenosis at an adjacent level (44% vs 17%).(24) **(EG 1)** A randomized noninferiority trial compared lumbar decompression alone with lumbar decompression with instrumental fusion in 262 patients (mean age 66 years) with leg or back pain due to lumbar spinal stenosis and single-level spondylolisthesis of 3 mm or more that has persisted despite nonoperative treatment (most patients with symptoms for more than 1 year), and concluded that there was noninferiority of decompression alone in patient-reported outcomes (eg, Oswestry Disability Index) at 2-year follow-up.(15) **(EG 1)** In the same trial, the decompression alone group had a slightly higher reoperation rate (12.5% vs 9.1%).(15) **(EG 1)**

Length of Stay

Analysis of procedure data for a commercially insured population shows 42% of patients undergoing lumbar fusion were discharged the day of or the day after surgery.(28) **(EG 3)** Analysis of procedure data for a Medicare-insured population shows 30% of patients undergoing lumbar fusion were discharged the day of or the day after surgery.(28) **(EG 3)**

Rationale

Use of this MCG care guideline helps the clinician identify, for a given procedure, which patient-specific factors and clinical conditions are appropriate for that procedure. The evidence-based clinical criteria assist the clinician in the decision to appropriately perform a procedure, evaluating whether the potential benefits of a procedure outweigh the potential risks. For Medicare enrollees, surgical MCG care guidelines also identify which procedures CMS has designated as inpatient only.

Use of these evidence-based clinical criteria to support decision making around the need for a given procedure is of benefit to the patient, as all procedures come with inherent risk that must be balanced by anticipated clinical benefit. Utilizing evidence-based clinical criteria enables a more accurate and patient-specific decision-making process. In addition, the use of evidence-based guidelines can help reduce unwarranted variation in care, such as divergent clinical thresholds to perform a procedure for clinically similar patients that vary across geographic regions, between facilities, and among individual clinicians.

Related CMS Coverage Guidance

This guideline supplements but does not replace, modify, or supersede existing Medicare regulations or applicable National Coverage Determinations (NCDs) or Local Coverage Determinations (LCDs).

Code of Federal Regulations (CFR): 42 CFR 412.3(54); 42 CFR 419.22(55); 42 CFR 422.101(56)

Internet-Only Manual (IOM) Citations: CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 1 - Inpatient Hospital Services Covered Under Part A(57); CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 6 - Hospital Services Covered Under Part B(58); CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 15 - Covered Medical and Other Health Services(59); CMS IOM Publication 100-08, Medicare Program Integrity Manual, Chapter 6, Section 6.5 - Medical Review of Inpatient Hospital Claims for Part A Payment(60)

Medicare Coverage Determinations: Medicare Coverage Database(61)

References

1. Gardocki RJ, Park AL. Degenerative disorders of the thoracic and lumbar spine. In: Azar FM, Beaty JH, editors. Campbell's Operative Orthopaedics. 14th ed. Philadelphia, PA: Elsevier; 2021:1719-1801 e9. [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11]
2. Drossopoulos PN, et al. Evolution of the transforaminal lumbar interbody fusion (TLIF): from open to percutaneous to patient-specific. Journal of Clinical Medicine 2024;13(8):Online. DOI: 10.3390/jcm13082271. [Context Link 1] View abstract...
3. Kumar RP, Elsayed GA, Hafez DM, Agarwal N. Advances in anterolateral approaches to the lumbar spine: a focus on technological developments. Neurosurgery Clinics of North America 2024;35(2):199-205. DOI: 10.1016/j.nec.2023.11.006. [Context Link 1] View abstract...
4. Esmende SM, Shah KN, Daniels AH. Spinopelvic fixation. Journal of the American Academy of Orthopedic Surgeons 2018;26(11):396-401. DOI: 10.5435/JAAOS-D-15-00738. [Context Link 1, 2, 3] View abstract...
5. Williams KD. Fractures, dislocations, and fracture-dislocations of the spine. In: Azar FM, Beaty JH, editors. Campbell's Operative Orthopaedics. 14th ed. Philadelphia, PA: Elsevier; 2021:1832-1923. [Context Link 1, 2, 3]
6. Darlow M, et al. A pathway for the diagnosis and treatment of lumbar spinal stenosis. Orthopedic Clinics of North America 2022;53(4):523-534. DOI: 10.1016/j.ocl.2022.05.006. [Context Link 1, 2] View abstract...
7. Williams KD. Spondylolisthesis. In: Azar FM, Beaty JH, editors. Campbell's Operative Orthopaedics. 14th ed. Philadelphia, PA: Elsevier; 2021:1802-1831.e1. [Context Link 1, 2, 3, 4]
8. Taba HA, Williams SK. Lateral lumbar interbody fusion. Neurosurgery Clinics of North America 2020;31(1):33-42. DOI: 10.1016/j.nec.2019.08.004. [Context Link 1, 2, 3] View abstract...
9. Derman PB, Singh K. Surgical strategies for the treatment of lumbar pseudarthrosis in degenerative spine surgery: a literature review and case study. HSS Journal 2020;16(2):183-187. DOI: 10.1007/s11420-019-09732-9. [Context Link 1, 2] View abstract...
10. Williams KD. Infections and tumors of the spine. In: Azar FM, Beaty JH, editors. Campbell's Operative Orthopaedics. 14th ed. Philadelphia, PA: Elsevier; 2021:1924-1956.e7. [Context Link 1, 2]

11. Ruparel S, et al. Surgical management of spinal tuberculosis-the past, present, and future. *Diagnostics* 2022;12(6):Online. DOI: 10.3390/diagnostics12061307. [Context Link 1, 2] View abstract...
12. Russo GS, et al. Thoracolumbar trauma. In: Browner BD, Jupiter JB, Krettek C, Anderson PA, editors. *Skeletal Trauma: Basic Science, Management, and Reconstruction*. 6th ed. Elsevier; 2020:988-1064. [Context Link 1, 2, 3]
13. Katz JN, Zimmerman ZE, Mass H, Makhni MC. Diagnosis and management of lumbar spinal stenosis: a review. *Journal of the American Medical Association* 2022;327(17):1688-1699. DOI: 10.1001/jama.2022.5921. [Context Link 1, 2] View abstract...
14. Gandhi J, et al. Neuro-urological sequelae of lumbar spinal stenosis. *International Journal of Neuroscience* 2018;128(6):554-562. DOI: 10.1080/00207454.2017.1400973. [Context Link 1] View abstract...
15. Austevoll IM, et al. Decompression with or without fusion in degenerative lumbar spondylolisthesis. *New England Journal of Medicine* 2021;385(6):526-538. DOI: 10.1056/NEJMoa2100990. [Context Link 1, 2, 3] View abstract...
16. Shukla GG, et al. Laminectomy with fusion is associated with greater functional improvement compared with laminectomy alone for the treatment of degenerative lumbar spondylolisthesis: a systematic review and meta-analysis. *Spine* 2023;48(12):874-884. DOI: 10.1097/BRS.0000000000004673. [Context Link 1] View abstract...
17. Crawford CH, et al. Current evidence regarding the treatment of pediatric lumbar spondylolisthesis: a report from the Scoliosis Research Society Evidence Based Medicine Committee. *Spine Deformity* 2017;5(5):284-302. DOI: 10.1016/j.jspd.2017.03.011. [Context Link 1, 2] View abstract...
18. Li Y, Fletcher ND. Indications for lumbar fusion in the skeletally mature adolescent: how to address oblique takeoff and limb length discrepancy. *Journal of Pediatric Orthopedics* 2021;41(Suppl 1):S59-S63. DOI: 10.1097/BPO.0000000000001805. [Context Link 1] View abstract...
19. Kouchehi R, Rocos B, Gandhi R, Lewis SJ, Lebel DE. Surgical management of high-grade paediatric spondylolisthesis: meta-analysis and systematic review. *European Spine Journal* 2023;32(2):436-446. DOI: 10.1007/s00586-022-07408-3. [Context Link 1] View abstract...
20. Waddell WH, Gupta R, Stephens BF. Thoracolumbar spine trauma. *Orthopedic Clinics of North America* 2021;52(4):481-489. DOI: 10.1016/j.ocl.2021.05.014. [Context Link 1] View abstract...
21. Nieblas-Bedolla E, Zuccato J, Kluger H, Zadeh G, Brastianos PK. Central nervous system metastases. *Hematology/Oncology Clinics of North America* 2022;36(1):161-188. DOI: 10.1016/j.hoc.2021.08.004. [Context Link 1, 2] View abstract...
22. Austevoll IM, et al. Comparative effectiveness of microdecompression alone vs decompression plus instrumented fusion in lumbar Degenerative Spondylolisthesis. *JAMA Network Open* 2020;3(9):e2015015. DOI: 10.1001/jamanetworkopen.2020.15015. [Context Link 1] View abstract...
23. Forsth P, et al. A randomized, controlled trial of fusion surgery for lumbar spinal stenosis. *New England Journal of Medicine* 2016;374(15):1413-1423. DOI: 10.1056/NEJMoa1513721. [Context Link 1, 2] View abstract...
24. Karlsson T, et al. Decompression alone or decompression with fusion for lumbar spinal stenosis: a randomized clinical trial with two-year MRI follow-up. *Bone and Joint Journal* 2022;104-B(12):1343-1351. DOI: 10.1302/0301-620X.104B12.BJJ-2022-0340.R1. [Context Link 1] View abstract...
25. Beck AW, Simpson AK. High-grade lumbar spondylolisthesis. *Neurosurgery Clinics of North America* 2019;30(3):291-298. DOI: 10.1016/j.nec.2019.02.002. [Context Link 1] View abstract...
26. Falck-Ytter Y, et al. Prevention of VTE in orthopedic surgery patients: antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2012;141(2 Suppl):e278S-e325S. DOI: 10.1378/chest.11-2404. (Reaffirmed 2024 Jul) [Context Link 1] View abstract...
27. Shaffer WO, Baisden JL, Fernand R, Matz PG, North American Spine Society. An evidence-based clinical guideline for antibiotic prophylaxis in spine surgery. *Spine Journal* 2013;23(10):1387-1392. DOI: 10.1016/j.spinee.2013.06.030. [Context Link 1] View abstract...
28. Proprietary health insurance data sources (2022-2023); and Medicare 5% Standard Analytical File (2021-2022). [Context Link 1, 2, 3]
29. Premier PINC AI™ Healthcare Database (PHD), 01/01/2022-12/31/2023. Premier, Inc. [Context Link 1]
30. Horowitz JA, Jain A, Puvanesarajah V, Qureshi R, Hassanzadeh H. Risk factors, additional length of stay, and cost associated with postoperative ileus following anterior lumbar interbody fusion in the elderly. *World Neurosurgery* 2018;115:e185-e189. DOI: 10.1016/j.wneu.2018.04.006. [Context Link 1] View abstract...
31. Hassanzadeh H, Bell J, Bhatia M, Puvanesarajah V. Incidental durotomy in lumbar spine surgery; risk factors, complications, and perioperative management. *Journal of the American Academy of Orthopedic Surgeons* 2021;29(6):e279-e286. DOI: 10.5435/JAAOS-D-20-00210. [Context Link 1] View abstract...
32. Cloud GW, et al. Increasing patient complexity is associated with longer in-hospital lengths of stay and higher rates of medical complications following primary 1- to 2-level lumbar fusion. *International Journal of Spine Surgery* 2022;16(6):1029-1033. DOI: 10.14444/8364. [Context Link 1] View abstract...
33. Hudson T. The role of social determinates of health in discharge practices. *Nursing Clinics of North America* 2021;56(3):369-378. DOI: 10.1016/j.cnur.2021.04.004. [Context Link 1, 2] View abstract...
34. Chovanec KA, Arsene C, Beck A, Zachrich K, Liedel B, Wolff-Elliott J. Association of discharge disposition with outcomes. *Population Health Management* 2021;24(1):116-121. DOI: 10.1089/pop.2019.0176. [Context Link 1] View abstract...
35. Centers for Medicare and Medicaid Services. "Condition of participation: Discharge planning." 42 CFR Pt. 482.43 Washington, DC 2022 Oct [accessed 2024 Oct 08] Accessed at: <https://www.ecfr.gov/>. [Context Link 1, 2, 3]
36. Adverse drug reactions and medication errors. In: Burchum JR, Rosenthal LD, editors. *Lehne's Pharmacology for Nursing Care*. 12th ed. Elsevier; 2025:64-75. [Context Link 1, 2]
37. Filson R. Musculoskeletal trauma and orthopedic surgery. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:1638-1672. [Context Link 1, 2, 3]

38. The intervention. In: Dutton M, editor. Dutton's Orthopaedic Examination, Evaluation, and Intervention. 6th ed. McGraw-Hill Education; 2023:343-368. [Context Link 1]
39. Skin integrity and wound care. In: Lynn P, editor. Taylor's Clinical Nursing Skills. 6th ed. Wolters Kluwer; 2023:452-540. [Context Link 1]
40. Targeted care planning and care transitions. In: Perez R, editor. CMSA's Integrated Case Management. 2nd ed. Springer Publishing Company LLC; 2023:178-189. [Context Link 1, 2]
41. Elements of Excellence in Transitions of Care (TOC). TOC Checklist [Internet] National Transitions of Care Coalition. Accessed at: <https://www.ntocc.org>. [accessed 2024 Sep 24] [Context Link 1]
42. Medical-surgical nursing. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:33-55. [Context Link 1]
43. Transition of care. In: Gillingham DC, editor. Foundations of Case Management. Blue Bayou Press; 2021:137-144. [Context Link 1]
44. Becker C, et al. Interventions to improve communication at hospital discharge and rates of readmission: a systematic review and meta-analysis. JAMA Network Open 2021;4(8):e2119346. DOI: 10.1001/jamanetworkopen.2021.19346. [Context Link 1] View abstract...
45. The nursing process in drug therapy and patient safety. In: Tucker RG, editor. Karch's Focus on Nursing Pharmacology. 9th ed. Wolters Kluwer; 2023:47-57. [Context Link 1]
46. Safe medication preparation. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. Clinical Nursing Skills and Techniques. 11th ed. Elsevier; 2025:586-607. [Context Link 1]
47. The case manager's roles, functions, and activities. In: Gillingham DC, editor. Foundations of Case Management. Blue Bayou Press; 2021:9-14. [Context Link 1]
48. Elsamadicy AA, et al. Patient- and hospital-related risk factors for non-routine discharge after lumbar decompression and fusion for spondylolisthesis. Clinical Neurology and Neurosurgery 2021;209:Online. DOI: 10.1016/j.clineuro.2021.106902. [Context Link 1] View abstract...
49. Bragg-Underwood T. Family health. In: Nies MA, editor. Community/Public Health Nursing. 8th ed. Elsevier; 2024:404-427. [Context Link 1]
50. Harding MM. Substance use disorders in acute care. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems. 12th ed. St. Louis, MO: Mosby; 2023:162-178. [Context Link 1]
51. Activity. In: Lynn P, editor. Taylor's Clinical Nursing Skills. 6th ed. Wolters Kluwer; 2023:541-632. [Context Link 1, 2]
52. Ryzner D. Musculoskeletal problems. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems. 12th ed. St. Louis, MO: Mosby; 2023:1674-1695. [Context Link 1]
53. Management of patients with dermatologic disorders. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:1815-1864. [Context Link 1]
54. Centers for Medicare and Medicaid Services. "Admissions." 42 CFR 412.3 Washington, DC 2023 Jul [accessed 2024 Oct 08] Accessed at: <https://www.ecfr.gov/current/title-42/chapter-IV/subchapter-B/part-412>. [Context Link 1]
55. Centers for Medicare and Medicaid Services. "Hospital services excluded from payment under the hospital outpatient prospective payment system." 42 CFR 419.22 Washington, DC 2023 Jul [accessed 2024 Oct 08] Accessed at: <https://www.ecfr.gov/>. [Context Link 1]
56. Centers for Medicare and Medicaid Services. "Requirements relating to basic benefits." 42 CFR 422.101 Washington, DC 2024 Sep 30 [accessed 2024 Oct 08] Accessed at: <https://www.ecfr.gov/>. [Context Link 1]
57. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2024 Jan 18] [Context Link 1]
58. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 6 - Hospital Services Covered Under Part B Rev. 12425 [Internet] Centers for Medicare and Medicaid Services. 2023 Dec Accessed at: <http://www.cms.gov/manuals/>. [accessed 2024 Jun 05] [Context Link 1]
59. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 15 - Covered Medical and Other Health Services Rev. 12425 [Internet] Centers for Medicare & Medicaid Services. 2023 Dec 21 Accessed at: <https://www.cms.gov/manuals/>. [accessed 2024 Feb 01] [Context Link 1]
60. Centers for Medicare and Medicaid Services. Medicare Program Integrity Manual. Chapter 6, Section 6.5 - Medical Review of Inpatient Hospital Claims for Part A Payment Rev. 10365 [Internet] Centers for Medicare and Medicaid Services. 2020 Oct 02 Accessed at: <https://www.cms.gov/medicare/regulations-guidance/manuals/internet-only-manuals-ioms>. [accessed 2024 Sep 26] [Context Link 1]
61. Medicare Coverage Database. [Internet] Centers for Medicare and Medicaid Services. Accessed at: <https://www.cms.gov/medicare-coverage-database/search.aspx?> Updated 2024 [accessed 2024 Sep 26] [Context Link 1]

Footnotes

[A] See Ambulatory Surgery Discharge and Complications: Common Complications and Conditions  ISC for further information. [A in Context Link 1]

[B] Patient is ambulatory or near baseline activity for age and development. [B in Context Link 1]

[C] Some patients may have their hydration needs met via alternative means (eg, percutaneous endoscopic gastrostomy tube). [C in Context Link 1]

[D] Low-dose unfractionated heparin, low-molecular-weight heparin, or intermittent pneumatic compression devices may be appropriate.(26) [D in Context Link 1]

[E] Prophylactic antibiotics may be considered to decrease the rate of infections following instrumented spine fusion.(27) [E in Context Link 1]

[F] Use Multimodal analgesia or individual analgesic agent as indicated. [F in Context Link 1]

[G] Discharge instructions should be given in the patient's and caregiver's native language using trained language interpreters whenever possible.(33) [G in Context Link 1]

Definitions

Hemodynamic stability

- Hemodynamic stability, as indicated by **1 or more** of the following:
 - Hemodynamic abnormalities at baseline or acceptable for next level of care
 - Patient hemodynamically stable, as indicated by **ALL** of the following(1)(2)(3)(4)(5):
 - Tachycardia absent
 - Hypotension absent
 - No evidence of inadequate perfusion (eg, no myocardial ischemia)
 - No other hemodynamic abnormalities (eg, no Orthostatic hypotension)

References

1. Puskarich MA, Jones AE. Shock. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:34-41.e1.
2. Lewis J, Patel B. Shock. In: Gershel JC, Rauch DA, editors. Caring for the Hospitalized Child: A Handbook of Inpatient Pediatrics. 3rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2023:107-113.
3. Ingbar DH, Thiele H. Cardiogenic shock and pulmonary edema. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:2250-2257.
4. Brant EB, Seymour CW, Angus DC. Sepsis and septic shock. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:2241-2249.
5. Singer M, et al. The Third International Consensus definitions for sepsis and septic shock (Sepsis-3). Journal of the American Medical Association 2016;315(8):801-810. DOI: 10.1001/jama.2016.0287.

Hypotension absent

- Hypotension absent[A], as indicated by **1 or more** of the following(1)(2)(3)(4)(5):
 - SBP greater than or equal to 90 mm Hg in adult or child age 10 years or older
 - Mean arterial pressure[B] greater than or equal to 70 mm Hg in adult or child age 10 years or older  MAP Calculator
 - Mean arterial pressure[B] at patient's baseline (eg, healthy adult with low SBP), at intentional therapeutic goal (eg, patient with heart failure), or acceptable for next level of care (eg, blood pressure stable and no significant signs or symptoms due to low blood pressure)  MAP Calculator
 - SBP greater than or equal to sum of 70 mm Hg plus twice patient's age in years in child 1 to 9 years of age
 - SBP greater than or equal to 70 mm Hg in infant 1 to 11 months of age

References

1. Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Angus DC. Approach to the patient with shock. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:658-665.
3. Massaro AF. Approach to the patient with shock. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:2235-2241.
4. Horeczko T. Pediatric cardiac disorders. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:2109-2131.e1.
5. Singh S, Holmes JF. Pediatric trauma. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:2052-2066.e3.

Footnotes

- A. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for

deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

B. The mean arterial pressure (MAP) takes into account both SBP and DBP readings.

Multimodal analgesia

- Multimodal analgesia involves the utilization of 2 or more analgesic agents with different mechanisms of action in order to provide additive or synergistic pain control, while minimizing side effects and reliance on opioids.(1)(2)(3)

References

1. D'Souza RS, Johnson RL. Regional and multimodal treatments of perioperative pain. In: Benzon H, editor. Practical Management of Pain. 6th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:355-373.e8.
2. George S, Johns M. Review of nonopioid multimodal analgesia for surgical and trauma patients. American Journal of Health-System Pharmacy 2020;77(24):2052-2063. DOI: 10.1093/ajhp/zxaa301.
3. Finnerup NB. Nonnarcotic methods of pain management. New England Journal of Medicine 2019;380(25):2440-2448. DOI: 10.1056/NEJMr1807061.

Orthostatic hypotension

- Orthostatic hypotension,[A][B] as indicated by **1 or more** of the following(1)(2)(3):
 - Fall in SBP of 20 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position
 - Fall in DBP of 10 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position

References

1. Shibao C, Lipsitz LA, Biaggioni I, American Society of Hypertension Writing Group. Evaluation and treatment of orthostatic hypotension. Journal of the American Society of Hypertension 2013 Jul-Aug;7(4):317-324. DOI: 10.1016/j.jash.2013.04.006.
2. Dalal AS, Van Hare GF. Syncope. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:592-599.
3. Fang JC, O'Gara PT. History and physical examination: an evidence-based approach. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:123-140.

Footnotes

- A. Concomitant measurements of the heart rate are important to measure to help diagnose subtypes of orthostatic hypotension (eg, the lack of a compensatory increase in heart rate is typical of autonomic failure and an exaggerated tachycardia may be reflective of volume depletion). However, the heart rate is not a component of the definition of orthostatic hypotension, which relies upon blood pressure alone.(1)(2)(3)
- B. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

Social Determinants of Health Assessment

- Risk of poor health outcomes may be increased by the presence of **1 or more** of the following social determinants of health(1)(2)(3):
 - Housing insecurity, as indicated by **1 or more** of the following:
 - Individual or caregiver's current living situation is **1 or more** of the following(4):
 - Does not have own housing (eg, staying in a hotel, shelter, or with others)
 - Has own housing (eg, house, apartment), but at risk of losing it in the future (ie, behind on rent or mortgage)
 - Has own housing (eg, house, apartment), but has lived in 3 or more places in past year
 - Current housing has **1 or more** of the following:
 - Electrical appliances (eg, stove, refrigerator) not working or unavailable
 - Insufficient heating or cooling
 - Insufficient ventilation
 - Lead paint or pipes
 - Mold
 - Pests (eg, bugs) or rodents
 - Smoke detectors not working or unavailable
 - Food insecurity, as indicated by **1 or more** of the following(5):
 - In the past year, individual or caregiver ran out of food and did not have money to buy more food.

- In the past year, individual or caregiver worried that they would run out of food before they received money to buy more food.
- Insufficient transportation, as indicated by **1 or more** of the following(6):
 - In the past year, individual or caregiver missed medical appointments or could not get medications due to lack of transportation.
 - In the past year, individual or caregiver missed nonmedical activities, work, or could not get things needed for daily living due to lack of transportation.
- Insufficient utilities, as indicated by **1 or more** of the following(7):
 - Utilities (eg, electricity, water, gas, or oil) are currently shut off or unavailable.
 - In the past year, electric, water, gas, or oil company threatened to shut off services.
- Personal safety risk, as indicated by **2 or more** of the following(5):
 - Individual is sometimes or frequently physically hurt by another person (including family member).
 - Individual is sometimes or frequently insulted or talked down to by another person (including family member).
 - Individual is sometimes or frequently threatened with physical harm by another person (including family member).
 - Individual is sometimes or frequently screamed or cursed at by another person (including family member).
- Insufficient dependent care, as indicated by **1 or more** of the following:
 - In the past year, individual or caregiver was unable to work due to lack of dependent care.
 - In the past year, individual or caregiver was unable to work more (additional) hours due to lack of dependent care.
 - In the past year, individual or caregiver missed medical appointments or could not get medications due to lack of dependent care.
 - In the past year, individual or caregiver missed nonmedical activities (eg, school, church, social activity) due to lack of dependent care.
- Depression risk, as indicated by **ALL** of the following(8):
 - In the past 2 weeks, individual had little interest or pleasure in normal activities on at least several days.
 - In the past 2 weeks, individual felt down, depressed, or hopeless on at least several days.

References

1. Scanlon A, Reinisch C. Social determinants of health. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:18-20.
2. Billioux A, Verlander K, Anthony S, Alley D. Standardized Screening for Health-Related Social Needs in Clinical Settings: the Accountable Health Communities Screening Tool. [Internet] National Academy of Sciences. 2017 May Accessed at: <https://nam.edu/>. [accessed 2024 Oct 01]
3. Addressing social determinants of health. In: Perez R, editor. *CMSA's Integrated Case Management*. 2nd ed. Springer Publishing Company LLC; 2023:62-75.
4. Sandel M, et al. Unstable housing and caregiver and child health in renter families. *Pediatrics* 2018;14(2):e20172199. DOI: 10.1542/peds.2017-2199.
5. Children's HealthWatch Survey. Screening Instrument [Internet] Children's HealthWatch. 2020 Sep Accessed at: <https://childrenshealthwatch.org/>. [accessed 2024 Sep 23]
6. PRAPARE®: Protocol for Responding to and Assessing Patient Assets, Risks, and Experiences Screening Tool. [Internet] Association of Asian Pacific Community Health Organizations (AAPCHO) and National Association of Community Health Centers (NACHC). 2016 Sep Accessed at: <https://prapare.org/the-prapare-screening-tool/>. [accessed 2024Oct 01]
7. Cook JT, et al. A brief indicator of household energy security: associations with food security, child health, and child development in US infants and toddlers. *Pediatrics* 2008;122(4):e867-75. DOI: 10.1542/peds.2008-0286.
8. Kroenke K, Spitzer RL, Williams JB. The Patient Health Questionnaire-2: validity of a two-item depression screener. *Medical Care* 2003;41(11):1284-92. DOI: 10.1097/01.MLR.0000093487.78664.3C.

Tachycardia absent

- Tachycardia^{[A][B]} absent, as indicated by **1 or more** of the following(1)(2):
 - Heart rate less than or equal to 100 beats per minute in adult or child 6 years or older
 - Heart rate less than or equal to 115 beats per minute in child 3 to 5 years of age
 - Heart rate less than or equal to 125 beats per minute in child 1 or 2 years of age
 - Heart rate less than or equal to 130 beats per minute in infant 6 to 11 months of age
 - Heart rate less than or equal to 150 beats per minute in infant 3 to 5 months of age
 - Heart rate less than or equal to 160 beats per minute in infant 1 or 2 months of age

References

1. Schrager DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. *Goldman-Cecil Medicine*. 27th ed. Elsevier; 2024:32-35.
2. Cico SJ, Caglar D. Care of the pediatric patient. In: Walls RM, editor. *Rosen's Emergency Medicine: Concepts and Clinical Practice*. 10th ed. Elsevier; 2023:1996-2004.e1.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. Interpretation of heart rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline heart rate, medications, and clinical impact. For example, an elderly patient on a beta-blocker medication with a baseline resting heart rate of 60 beats per minute may be clinically tachycardic at a heart rate of 94 beats per minute. Likewise, a patient who is upset, in pain, or nervous in the emergency department with a heart rate of 106 beats per minute may meet the technical definition of tachycardia, but this tachycardia (absent associated findings such as chest pain or hypotension) may not be clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachycardia. Whether a heart rate above or below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.

Codes

ICD-10 Diagnosis: A18.01, C41.2, C79.40, C79.49, D16.6, D32.1, D33.4, D42.1, D43.4, D48.0, D49.2, G06.1, M43.05, M43.06, M43.07, M43.09, M43.15, M43.16, M43.17, M43.19, M46.45, M46.46, M46.47, M47.15, M47.16, M47.25, M47.26, M47.27, M47.815, M47.816, M47.817, M47.895, M47.896, M47.897, M48.05, M48.061, M48.062, M48.07, M51.05, M51.06, M51.15, M51.16, M51.17, M51.25, M51.26, M51.27, M51.35, M51.360, M51.361, M51.362, M51.369, M51.370, M51.371, M51.372, M51.379, M51.85, M51.86, M51.87, M51.A0, M51.A1, M51.A2, M51.A3, M51.A4, M51.A5, M54.15, M54.16, M54.17, M96.1, M99.13, M99.23, M99.33, M99.43, M99.53, M99.63, M99.73, Q76.2, S33.0XXA, S33.100A, S33.101A, S33.110A, S33.111A, S33.120A, S33.121A, S33.130A, S33.131A, S33.140A, S33.141A, S34.101A, S34.102A, S34.103A, S34.104A, S34.105A, S34.109A, S34.111A, S34.112A, S34.113A, S34.114A, S34.115A, S34.119A, S34.121A, S34.122A, S34.123A, S34.124A, S34.125A, S34.129A [Hide]

ICD-10 Procedure: 0RGA070, 0RGA071, 0RGA07J, 0RGA0A0, 0RGA0AJ, 0RGA0J0, 0RGA0J1, 0RGA0JJ, 0RGA0K0, 0RGA0K1, 0RGA0KJ, 0RGA370, 0RGA371, 0RGA37J, 0RGA3A0, 0RGA3AJ, 0RGA3J0, 0RGA3J1, 0RGA3JJ, 0RGA3K0, 0RGA3K1, 0RGA3KJ, 0SG0070, 0SG0071, 0SG007J, 0SG00A0, 0SG00AJ, 0SG00J0, 0SG00J1, 0SG00JJ, 0SG00K0, 0SG00K1, 0SG00KJ, 0SG0370, 0SG0371, 0SG037J, 0SG03A0, 0SG03AJ, 0SG03J0, 0SG03J1, 0SG03JJ, 0SG03K0, 0SG03K1, 0SG03KJ, 0SG1070, 0SG1071, 0SG107J, 0SG10A0, 0SG10AJ, 0SG10J0, 0SG10J1, 0SG10JJ, 0SG10K0, 0SG10K1, 0SG10KJ, 0SG1370, 0SG1371, 0SG137J, 0SG13A0, 0SG13AJ, 0SG13J0, 0SG13J1, 0SG13JJ, 0SG13K0, 0SG13K1, 0SG13KJ, 0SG3070, 0SG3071, 0SG307J, 0SG30A0, 0SG30AJ, 0SG30J0, 0SG30J1, 0SG30JJ, 0SG30K0, 0SG30K1, 0SG30KJ, 0SG3370, 0SG3371, 0SG337J, 0SG33A0, 0SG33AJ, 0SG33J0, 0SG33J1, 0SG33JJ, 0SG33K0, 0SG33K1, 0SG33KJ, XRGA0EA, XRGA0R7, XRGA3R7, XRGA4R7, XRGB0EA, XRGB0R7, XRGB3R7, XRGB4R7, XRGC0EA, XRGC0R7, XRGC3R7, XRGC4R7, XRGD0EA, XRGD0R7, XRGD3R7 [Hide]

CPT®: 22533, 22534, 22558, 22585, 22586, 22612, 22614, 22630, 22632, 22633, 22634, 63048

CPT copyright 2024 American Medical Association. All rights reserved.

MCG Health
Inpatient & Surgical Care 29th Edition
Copyright © 2025 MCG Health, LLC
All Rights Reserved

Last Update: 6/13/2025 6:38:59 AM
Build Number: 29.1.20250613001137.022065