

# CS 5565 - Intro to Statistical Learning

## **Lecture: Linear Regression**

Adu Baffour, PhD

# Lecture Objectives


- Perform linear regression with a single predictor variable.
- Estimate the standard error of regression coefficients.
- Evaluate the goodness of fit of a regression.
- Perform linear regression with multiple predictor variables.
- Evaluate the relative importance of variables in a multiple linear regression.
- Include interaction effects in a multiple linear regression.
- Perform linear regression with qualitative predictor variables.
- Model non-linear relationships using polynomial regression.
- Identify non-linearity in a data set.

## Linear regression

- Linear regression is a simple approach to supervised learning. It assumes that the dependence of  $Y$  on  $X_1, X_2, \dots, X_p$  is linear.


## Linear regression

- Linear regression is a simple approach to supervised learning. It assumes that the dependence of  $Y$  on  $X_1, X_2, \dots, X_p$  is linear.
- True regression functions are never linear!



## Linear regression

- Linear regression is a simple approach to supervised learning. It assumes that the dependence of  $Y$  on  $X_1, X_2, \dots, X_p$  is linear.
- True regression functions are never linear!



- although it may seem overly simplistic, linear regression is extremely useful both conceptually and practically.




# Linear regression for the advertising data

Consider the advertising data shown on the next slide.

Questions we might ask:

- Is there a relationship between advertising budget and sales?
- How strong is the relationship between advertising budget and sales?
- Which media contribute to sales?
- How accurately can we predict future sales?
- Is the relationship linear?
- Is there synergy among the advertising media?

# Advertising data



## Simple linear regression using a single predictor $X$ .

- We assume a model

$$Y = \beta_0 + \beta_1 X + \epsilon,$$

where  $\beta_0$  and  $\beta_1$  are two unknown constants that represent the *intercept* and *slope*, also known as *coefficients* or *parameters*, and  $\epsilon$  is the error term.

- Given some estimates  $\hat{\beta}_0$  and  $\hat{\beta}_1$  for the model coefficients, we predict future sales using

$$\hat{y} = \hat{\beta}_0 + \hat{\beta}_1 x,$$

where  $\hat{y}$  indicates a prediction of  $Y$  on the basis of  $X = x$ . The *hat* symbol denotes an estimated value.



## Estimation of the parameters by least squares

- Let  $\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i$  be the prediction for  $Y$  based on the  $i$ th value of  $X$ . Then  $e_i = y_i - \hat{y}_i$  represents the  $i$ th *residual*

## Estimation of the parameters by least squares

- Let  $\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i$  be the prediction for  $Y$  based on the  $i$ th value of  $X$ . Then  $e_i = y_i - \hat{y}_i$  represents the  $i$ th *residual*
- We define the *residual sum of squares* (RSS) as

$$\text{RSS} = e_1^2 + e_2^2 + \cdots + e_n^2,$$

or equivalently as

$$\text{RSS} = (y_1 - \hat{\beta}_0 - \hat{\beta}_1 x_1)^2 + (y_2 - \hat{\beta}_0 - \hat{\beta}_1 x_2)^2 + \cdots + (y_n - \hat{\beta}_0 - \hat{\beta}_1 x_n)^2.$$

## Estimation of the parameters by least squares

- Let  $\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i$  be the prediction for  $Y$  based on the  $i$ th value of  $X$ . Then  $e_i = y_i - \hat{y}_i$  represents the  $i$ th *residual*
- We define the *residual sum of squares* (RSS) as


$$\text{RSS} = e_1^2 + e_2^2 + \cdots + e_n^2,$$

or equivalently as

$$\text{RSS} = (y_1 - \hat{\beta}_0 - \hat{\beta}_1 x_1)^2 + (y_2 - \hat{\beta}_0 - \hat{\beta}_1 x_2)^2 + \cdots + (y_n - \hat{\beta}_0 - \hat{\beta}_1 x_n)^2.$$

- The least squares approach chooses  $\hat{\beta}_0$  and  $\hat{\beta}_1$  to minimize the RSS.

## Example: advertising data



The least squares fit for the regression of **sales** onto **TV**.  
In this case a linear fit captures the essence of the relationship, although it is somewhat deficient in the left of the plot.

## Assessing the Accuracy of the Coefficient Estimates

- The standard error of an estimator reflects how it varies under repeated sampling. We have

$$\text{SE}(\hat{\beta}_1)^2 = \frac{\sigma^2}{\sum_{i=1}^n (x_i - \bar{x})^2}, \quad \text{SE}(\hat{\beta}_0)^2 = \sigma^2 \left[ \frac{1}{n} + \frac{\bar{x}^2}{\sum_{i=1}^n (x_i - \bar{x})^2} \right],$$

where  $\sigma^2 = \text{Var}(\epsilon)$

## Assessing the Accuracy of the Coefficient Estimates

- The standard error of an estimator reflects how it varies under repeated sampling. We have

$$\text{SE}(\hat{\beta}_1)^2 = \frac{\sigma^2}{\sum_{i=1}^n (x_i - \bar{x})^2}, \quad \text{SE}(\hat{\beta}_0)^2 = \sigma^2 \left[ \frac{1}{n} + \frac{\bar{x}^2}{\sum_{i=1}^n (x_i - \bar{x})^2} \right],$$

where  $\sigma^2 = \text{Var}(\epsilon)$

- These standard errors can be used to compute *confidence intervals*. A 95% confidence interval is defined as a range of values such that with 95% probability, the range will contain the true unknown value of the parameter. It has the form

$$\hat{\beta}_1 \pm 2 \cdot \text{SE}(\hat{\beta}_1).$$

## Confidence intervals — continued

That is, there is approximately a 95% chance that the interval

$$\left[ \hat{\beta}_1 - 2 \cdot \text{SE}(\hat{\beta}_1), \hat{\beta}_1 + 2 \cdot \text{SE}(\hat{\beta}_1) \right]$$

will contain the true value of  $\beta_1$  (under a scenario where we got repeated samples like the present sample)

## Confidence intervals — continued

That is, there is approximately a 95% chance that the interval

$$\left[ \hat{\beta}_1 - 2 \cdot \text{SE}(\hat{\beta}_1), \hat{\beta}_1 + 2 \cdot \text{SE}(\hat{\beta}_1) \right]$$

will contain the true value of  $\beta_1$  (under a scenario where we got repeated samples like the present sample)

For the advertising data, the 95% confidence interval for  $\beta_1$  is  $[0.042, 0.053]$



# Hypothesis testing

- Standard errors can also be used to perform *hypothesis tests* on the coefficients. The most common hypothesis test involves testing the *null hypothesis* of

$H_0$  :     There is no relationship between  $X$  and  $Y$   
              versus the *alternative hypothesis*

$H_A$  :     There is some relationship between  $X$  and  $Y$ .

## Hypothesis testing

- Standard errors can also be used to perform *hypothesis tests* on the coefficients. The most common hypothesis test involves testing the *null hypothesis* of

$H_0$  :     There is no relationship between  $X$  and  $Y$   
              versus the *alternative hypothesis*

$H_A$  :     There is some relationship between  $X$  and  $Y$ .

- Mathematically, this corresponds to testing

$$H_0 : \beta_1 = 0$$

versus

$$H_A : \beta_1 \neq 0,$$

since if  $\beta_1 = 0$  then the model reduces to  $Y = \beta_0 + \epsilon$ , and  $X$  is not associated with  $Y$ .

## Hypothesis testing — continued

- To test the null hypothesis, we compute a *t-statistic*, given by

$$t = \frac{\hat{\beta}_1 - 0}{\text{SE}(\hat{\beta}_1)},$$

- This will have a  $t$ -distribution with  $n - 2$  degrees of freedom, assuming  $\beta_1 = 0$ .
- Using statistical software, it is easy to compute the probability of observing any value equal to  $|t|$  or larger. We call this probability the *p-value*.

## Results for the advertising data

	Coefficient	Std. Error	t-statistic	p-value
Intercept	7.0325	0.4578	15.36	< 0.0001
TV	0.0475	0.0027	17.67	< 0.0001

# Multiple Linear Regression

- Here our model is

$$Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \cdots + \beta_p X_p + \epsilon,$$

- We interpret  $\beta_j$  as the *average* effect on  $Y$  of a one unit increase in  $X_j$ , *holding all other predictors fixed*. In the advertising example, the model becomes

$$\text{sales} = \beta_0 + \beta_1 \times \text{TV} + \beta_2 \times \text{radio} + \beta_3 \times \text{newspaper} + \epsilon.$$

## Interpreting regression coefficients

- The ideal scenario is when the predictors are uncorrelated — a *balanced design*:
  - Each coefficient can be estimated and tested separately.
  - Interpretations such as “*a unit change in  $X_j$  is associated with a  $\beta_j$  change in  $Y$ , while all the other variables stay fixed*”, are possible.
- Correlations amongst predictors cause problems:
  - The variance of all coefficients tends to increase, sometimes dramatically
  - Interpretations become hazardous — when  $X_j$  changes, everything else changes.
- *Claims of causality* should be avoided for observational data.

## Results for advertising data

	Coefficient	Std. Error	t-statistic	p-value
Intercept	2.939	0.3119	9.42	< 0.0001
TV	0.046	0.0014	32.81	< 0.0001
radio	0.189	0.0086	21.89	< 0.0001
newspaper	-0.001	0.0059	-0.18	0.8599

### Correlations:

	TV	radio	newspaper	sales
TV	1.0000	0.0548	0.0567	0.7822
radio		1.0000	0.3541	0.5762
newspaper			1.0000	0.2283
sales				1.0000

## Some important questions

1. *Is at least one of the predictors  $X_1, X_2, \dots, X_p$  useful in predicting the response?*



## Some important questions

1. *Is at least one of the predictors  $X_1, X_2, \dots, X_p$  useful in predicting the response?*
2. *Do all the predictors help to explain  $Y$ , or is only a subset of the predictors useful?*

## Some important questions

1. *Is at least one of the predictors  $X_1, X_2, \dots, X_p$  useful in predicting the response?*
2. *Do all the predictors help to explain  $Y$ , or is only a subset of the predictors useful?*
3. *How well does the model fit the data?*

## Some important questions

1. *Is at least one of the predictors  $X_1, X_2, \dots, X_p$  useful in predicting the response?*
2. *Do all the predictors help to explain  $Y$ , or is only a subset of the predictors useful?*
3. *How well does the model fit the data?*
4. *Given a set of predictor values, what response value should we predict, and how accurate is our prediction?*


## Other Considerations in the Regression Model

### *Qualitative Predictors*

- Some predictors are not *quantitative* but are *qualitative*, taking a discrete set of values.
- These are also called *categorical* predictors or *factor variables*.
- See for example the scatterplot matrix of the credit card data in the next slide.

In addition to the 7 quantitative variables shown, there are four qualitative variables: **gender**, **student** (student status), **status** (marital status), and **ethnicity** (Caucasian, African American (AA) or Asian).

## Credit Card Data



## Qualitative Predictors — continued

Example: investigate differences in credit card balance between males and females, ignoring the other variables. We create a new variable

$$x_i = \begin{cases} 1 & \text{if } i\text{th person is female} \\ 0 & \text{if } i\text{th person is male} \end{cases}$$

Resulting model:

$$y_i = \beta_0 + \beta_1 x_i + \epsilon_i = \begin{cases} \beta_0 + \beta_1 + \epsilon_i & \text{if } i\text{th person is female} \\ \beta_0 + \epsilon_i & \text{if } i\text{th person is male.} \end{cases}$$

Intrepretation?

## Modelling interactions — Advertising data

Model takes the form

$$\begin{aligned}\text{sales} &= \beta_0 + \beta_1 \times \text{TV} + \beta_2 \times \text{radio} + \beta_3 \times (\text{radio} \times \text{TV}) + \epsilon \\ &= \beta_0 + (\beta_1 + \beta_3 \times \text{radio}) \times \text{TV} + \beta_2 \times \text{radio} + \epsilon.\end{aligned}$$

Results:

	Coefficient	Std. Error	t-statistic	p-value
Intercept	6.7502	0.248	27.23	< 0.0001
TV	0.0191	0.002	12.70	< 0.0001
radio	0.0289	0.009	3.24	0.0014
TV×radio	0.0011	0.000	20.73	< 0.0001

# Interpretation

- The results in this table suggests that interactions are important.
- The p-value for the interaction term **TV**×**radio** is extremely low, indicating that there is strong evidence for  $H_A: \beta_3 \neq 0$ .



## Interpretation — continued

- The coefficient estimates in the table suggest that an increase in TV advertising of \$1,000 is associated with increased sales of

$$(\hat{\beta}_1 + \hat{\beta}_3 \times \text{radio}) \times 1000 = 19 + 1.1 \times \text{radio units.}$$

- An increase in radio advertising of \$1,000 will be associated with an increase in sales of

$$(\hat{\beta}_2 + \hat{\beta}_3 \times \text{TV}) \times 1000 = 29 + 1.1 \times \text{TV units.}$$

# Hierarchy

- Sometimes it is the case that an interaction term has a very small p-value, but the associated main effects (in this case, **TV** and **radio**) do not.
- The *hierarchy principle*:

*If we include an interaction in a model, we should also include the main effects, even if the p-values associated with their coefficients are not significant.*

## Hierarchy — continued

- The rationale for this principle is that interactions are hard to interpret in a model without main effects — their meaning is changed.
- Specifically, the interaction terms also contain main effects, if the model has no main effect terms.

## Interactions between qualitative and quantitative variables

Consider the **Credit** data set, and suppose that we wish to predict **balance** using **income** (quantitative) and **student** (qualitative).

Without an interaction term, the model takes the form


$$\begin{aligned}\text{balance}_i &\approx \beta_0 + \beta_1 \times \text{income}_i + \begin{cases} \beta_2 & \text{if } i\text{th person is a student} \\ 0 & \text{if } i\text{th person is not a student} \end{cases} \\ &= \beta_1 \times \text{income}_i + \begin{cases} \beta_0 + \beta_2 & \text{if } i\text{th person is a student} \\ \beta_0 & \text{if } i\text{th person is not a student.} \end{cases}\end{aligned}$$

With interactions, it takes the form

$$\begin{aligned}\text{balance}_i &\approx \beta_0 + \beta_1 \times \text{income}_i + \begin{cases} \beta_2 + \beta_3 \times \text{income}_i & \text{if student} \\ 0 & \text{if not student} \end{cases} \\ &= \begin{cases} (\beta_0 + \beta_2) + (\beta_1 + \beta_3) \times \text{income}_i & \text{if student} \\ \beta_0 + \beta_1 \times \text{income}_i & \text{if not student} \end{cases}\end{aligned}$$

# Non-linear effects of predictors

polynomial regression on **Auto** data



The figure suggests that

$$\text{mpg} = \beta_0 + \beta_1 \times \text{horsepower} + \beta_2 \times \text{horsepower}^2 + \epsilon$$

may provide a better fit.

	Coefficient	Std. Error	t-statistic	p-value
Intercept	56.9001	1.8004	31.6	< 0.0001
horsepower	-0.4662	0.0311	-15.0	< 0.0001
horsepower <sup>2</sup>	0.0012	0.0001	10.1	< 0.0001

# Summary

- Simple linear regression assumes a linear relationship between the predictor ( $X$ ) and the response ( $Y$ ).
- The best estimates for the coefficients are obtained by finding the regression line that fits the training dataset points as closely as possible. This line can be obtained by minimizing the least squares criteria.
- The most common hypothesis test involves testing the null hypothesis versus the alternative hypothesis.
- The p-value indicates how likely it is to observe a meaningful association between  $X$  and  $Y$  by some bizarre random error or chance, as opposed to there being a true relationship between  $X$  and  $Y$ .
- A simple way to extend the linear model is through polynomial regression.



Thank you.

Any questions?