



REVIEW ARTICLE

## Beneficial elements for agricultural crops and their functional relevance in defence against stresses

Simranjeet Kaur<sup>a</sup>, Navneet Kaur<sup>a</sup>, Kadambot H. M. Siddique<sup>b</sup> and Harsh Nayyar<sup>a</sup>

<sup>a</sup>Department of Botany, Panjab University, Chandigarh, India; <sup>b</sup>The UWA Institute of Agriculture, The University of Western Australia, Crawley, Australia

### ABSTRACT

The beneficial elements are not deemed essential for all crops but may be vital for particular plant taxa. The distinction between beneficial and essential is often difficult in the case of some trace elements. Elements such as aluminium (Al), cobalt (Co), sodium (Na), selenium (Se) and silicon (Si) are considered beneficial for plants. These elements are not critical for all plants but may improve plant growth and yield. Pertinently, beneficial elements reportedly enhance resistance to abiotic stresses (drought, salinity, high temperature, cold, UV stress, and nutrient toxicity or deficiency) and biotic stresses (pathogens and herbivores) at their low levels. However, the essential-to-lethal range for these elements is somewhat narrow. The effect of beneficial elements at low levels deserves more attention with regard to using them to fertilize crops to boost crop production under stress and to enhance plant nutritional value as a feed or food. A more holistic approach to plant nutrition would not only be restricted to nutrients essential to survival but would also include mineral elements at levels beneficial for best growth. Here, we describe the uptake mechanisms of various beneficial elements, their favourable aspects, and the role of these elements in conferring tolerance against abiotic and biotic stresses.

### ARTICLE HISTORY

Received 17 March 2015  
Accepted 24 September 2015

### KEYWORDS

Aluminium; cobalt;  
selenium; silicon; sodium;  
biotic; abiotic; stress  
mitigation

### Introduction

There is rising awareness of the need to pay greater attention to the role of trace elements in plant nutrition as we seek to elucidate the adverse effects of deficiencies and toxicities, and avoid suboptimal concentrations that limit the attainment of optimum economic yields of crops, animal productivity and welfare. The most important role of these elements does not lie in its general acceptance of essentiality, but rather in its most striking and unique function in conferring tolerance in plants to various abiotic and/or biotic stresses. There are diverse opinions regarding the 'essentiality' of mineral nutrients in plants. Early classical reports on the essentiality of minerals came from Arnon and Stout (1939), who identified three criteria to consider an element as essential: 1) a plant cannot complete its life cycle in the absence of the mineral element, 2) the function of the element is not replaceable by another mineral element, and 3) the element is directly involved in plant metabolism. Other elements that promote growth in many plant species but are not needed to complete the plant life cycle, or fail to meet the Arnon and Stout criteria on other grounds, are termed beneficial elements. Of the 92 known elements, 17 are considered essential and are categorised into macro-elements such as C, H, O, Ca, K, Mg, N, S and P (requirement  $>1000 \text{ mg kg}^{-1}$  dry weight – and micro-/trace elements such as B, Cl, Cu, Fe, Mn,

Mo, Ni and Zn (requirement  $<100 \text{ mg kg}^{-1}$  dry weight). Certain other elements, such as Al, Si, Co, Na and Se, do not meet the 'essentiality' criteria for mineral nutrients but have a ubiquitous presence in both soil and water and can be widely taken up and used by plants. Such elements, which may enhance biomass and yield but may not be required for species to survive, are termed functional/beneficial elements (Marschner 2012). Not all metabolic functions need a unique element to function; there are a number of chemically different but physically similar elements that can substitute the functions of essential elements in various metabolic processes. For example, Na, Rb and K, Se and S, Co and Ni, and Si and C often replace each other in certain non-specific metabolic processes. Therefore, it can be opined that the element replacing the essential element may be more efficient than any other element.

Various approaches have been executed to analyze the growth-promoting effects of beneficial elements. For instance, plants have been grown in the presence or absence of beneficial elements with positive and negative effects at the plant tissue level determined. The beneficial elements were effective at their low concentrations and acted as co-factors for some specific enzymes. It is critical to study beneficial elements in the context of ecological processes to establish how valuable these elements are for alleviating abiotic stresses and how plants accumulating these elements interact with pathogens/herbivores (Pilon-Smits et al. 2009). Better nutrient cycling contributes towards an improved ecological environment, resulting in an ecologically sound and sustainable flow of these elements. Soil–plant system strategies that have been assumed to perk up human micronutrient nutrition mainly include: a) beneficial elements flow from soils to the edible parts of crops, and b) a better knowledge of the relationship between the content and bioavailability of beneficial elements in soils and those in edible crop products. The omission of beneficial nutrients in commercial production may mean that plants are not grown to their optimum genetic potential but are merely produced at a subsistence level. Here, we summarize our current knowledge about the abundance of the beneficial elements in soil, their uptake mechanisms, and roles in plant growth, especially under various types of abiotic and biotic stresses.

### **Uptake mechanisms of beneficial elements**


To illustrate the beneficial role of these elements, it is fundamental to understand their uptake and transport mechanisms as well as their movement between different tissues and within cell compartments. This section describes the uptake mechanisms of various beneficial elements, which are summarized in Figure 1.

#### **Aluminium**

Aluminium is present in acidic soils ( $\text{pH} < 5.5$ ) and released in the form of  $\text{Al}^{3+}$ ,  $\text{Al}(\text{OH})_2^+$ , and  $\text{Al}(\text{OH})^{2+}$ . However,  $\text{Al}^{3+}$  uptake, whether symplastic (storing  $\text{Al}^{3+}$  in the less toxic forms complexed with organic acids) or apoplastic ( $\text{Al}^{3+}$  binding to cell walls and obstructing its transport to symplasm), is under debate. There is evidence that a high shoot aluminium level in soluble form is transported through the xylem to accumulate in leaf vacuoles or the apoplast (Hodson & Evans 1995).  $\text{Al}^{3+}$  sensing involves a specific receptor – ALMT (aluminium-activated malate transporter) – which encodes the  $\text{Al}^{3+}$ -activated anion channel that facilitates malate efflux from root apices (Figure 1a).

#### **Cobalt**

Soil pH is the most important factor for cobalt uptake in roots, which increases with decreasing pH. Korshunova et al. (1999) found that IRT1, an *Arabidopsis thaliana* iron transporter, assisted in the uptake of cobalt into plant cells. Consequently,  $\text{Co}^{2+}$  uptake occurs in roots via IRT1 transporters (Figure 1b), which seems to have eight membrane-spanning domains and a putative histidine-rich, metal-binding domain. Cobalt transport occurs in the form of the cobalt ion ( $\text{Co}^{2+}$ ) through cortical cells via passive diffusion and active transport. Upward transport in the xylem tissue chiefly takes place by transpirational flow (Jarosick et al. 1988; Figure 1b).



**Figure 1.** Uptake mechanisms of beneficial elements (a)  $\text{Al}^{3+}$ -activated organic anion influx takes place via ALMT (aluminium-activated malate transporter) transporters in root apices, and then interacts with the protein to form  $\text{Al}^{3+}$  citrate/oxalate-complexes. (b) For cobalt,  $\text{Co}^{2+}$  ion uptake takes place in roots via IRT1 (iron transporter 1) transporters. Transport of cobalt occurs via passive diffusion and active transport. (c) Selenate ( $\text{SeO}_4^{2-}$ ) uptake takes place across the root plasma membrane by high-affinity sulphate transporters whose expression is regulated positively by *O*-acetylserine and negatively by sulphate and glutathione. Selenate is then transported to chloroplast where it is finally reduced to GS-Se-(glutathione conjugated selenide) via various enzymatic and non-enzymatic reactions (Terry et al. 2000). (d) Si uptake occurs in the form of silicic acid [ $\text{Si}(\text{OH})_4$ ] through roots via silicon transporters Lsi1 (low silicon 1), which is then transported through xylem sap to the leaf cells via Lsi6 transporter as reported in rice (Ma et al. 2006). 1 and 2 shows the Si uptake in maize/barley and rice, respectively, via specific transporters attributed to their root structure. OsLsi1, 2: *Oryza sativa* Low Silicon transporter1, 2; ZmLsi1, 2: *Zea mays* Low Silicon transporter1, 2; HvLsi1, 2: *Hordeum vulgare* Low Silicon transporter1, 2 (e)  $\text{Na}^+$  uptake, efflux, long-distance transport, and cellular partitioning in roots and shoots. Abbreviations: AKT1, inward rectifying  $\text{K}^+$  channel; CHX, cation:  $\text{H}^+$  exchanger; HKT, high affinity  $\text{K}^+$  transporter; NHX, vacuolar  $\text{Na}^+:\text{H}^+$  antiporter; NSCC, non-selective cation channel; ORC, outward rectifying  $\text{K}^+$  channel; SOS1, plasma membrane  $\text{Na}^+:\text{H}^+$  antiporter (Maathuis 2014).

## Selenium

Se is present in different forms (elemental selenium, selenite, selenate, thioselenate, and selenide) in the soil. However, plants can absorb and sequester Se in the form of selenite and selenate. Selenate readily competes with sulphate uptake and is taken up via sulphur transporters in the root plasma membrane. Genes encoding sulphate transporters – SHST1, SHST2, and SHST3 (*Stylosanthes hamata* transporters 1, 2, and 3) – have been isolated from the tropical legume *Stylosanthes hamata* (Smith et al. 1995) and one gene HVST1 (*Hordeum vulgare* sulphate transporter 1) from

barley (*Hordeum vulgare* L.; Smith et al. 1997). The expression of these transporter genes is positively and negatively regulated by special regulators such as glutathione (GSH), O-acetylserine, and S (sulfur) (Davidian et al. 2000; Figure 1c).

### Silicon

Plants take up silicon in the form of monosilicic acid,  $\text{Si}(\text{OH})_4$ , through roots via silicon transporters Lsi1 and Lsi2 (low silicons 1 and 2; Figure 1d), discovered from rice plants (Ma et al. 2006). A second Si transporter, Lsi2, reportedly involves Lsi1 mutant approach, i.e. different from Lsi1, and acts as an efflux transporter in rice. Lsi2 expression and localization in tissues and cells is similar to Lsi1; however, Lsi2 is confined to the proximal side of the exodermis and endodermis cells (Ma et al. 2007). After the  $\text{Si}^{4+}$  uptake through roots via Lsi1 and Lsi2, more than 90% is transported to the shoots by transpirational flow in the xylem. From the stem xylem sap, silicic acid is transported to other leaf cells via Lsi6 transporter reported in rice (Ma et al. 2006). Similar to rice transporters (OsLsi1,2), HvLsi(1,2), ZmLsi(1,2), and CmLsi (1,2) silicon transporters have been discovered in barley (*Hordeum vulgare* L.), maize (*Zea mays* L.), respectively, that are localized at epidermal, hypodermal, and cortical cells (Chiba et al. 2009; Mitani et al. 2009; Figure 1d).

### Sodium

Sodium ( $\text{Na}^+$ ) at its low levels is beneficial in the presence of low potassium ( $\text{K}^+$ ) levels. At high  $\text{Na}^+$  levels, low-affinity fluxes catalyze  $\text{Na}^+$  movement via  $\text{K}^+$  selective ion channels such as AKT1 (inward rectifying  $\text{K}^+$  channel), HKT (high-affinity  $\text{K}^+$  channels), non-selective ion channels (NSCC), and outward rectifying  $\text{K}^+$  channels (ORC) (Maathuis 2014; Figure 1e). However, in salt-sensitive plants, there is a  $\text{Na}^+\text{-H}^+$  antiporter, SOS1 (salt overly sensitive), resulting in the efflux of  $\text{Na}^+$  ions. Elevated  $\text{Ca}^{2+}$  levels are sensed by SOS3 (a  $\text{Ca}^{2+}$  binding protein), which stimulates SOS2 (a serine threonine protein kinase). The SOS3–SOS2 protein complex then phosphorylates SOS1 that induces the plasma membrane  $\text{Na}^+\text{-H}^+$  antiporter (NHX1), leading to efflux of  $\text{Na}^+$  ions. NHX1 antiport activity can be regulated by SOS2 independent of SOS3, probably by SOS3-like  $\text{Ca}^{2+}$  binding proteins (SCaBP). High  $\text{Na}^+$  levels stimulate ABA production, which negatively regulates SOS2/SOS1 and NHX1 (Silva & Gerós 2009; Figure 1e).

## Functional relevance against abiotic stresses

### Aluminium

Aluminium ( $\text{Al}^{3+}$ ) is the third most abundant element in the earth's crust. In soil, 7% of total aluminium is present in solubilized form under acidic conditions, increasing their availability to plants. No convincing evidence implies that aluminium is a vital nutrient for plants, while there are a few reports of its beneficial effects (Huang & Barchelard 1993; Figure 2). Soluble Al is released from acidic soils in the form of  $\text{Al}^{3+}$ ,  $\text{Al}(\text{OH})^{2+}$ , and  $\text{Al}(\text{OH})_2^+$ . Among all soluble forms of Al in soil, inorganic aluminium in the form of  $(\text{Al}(\text{H}_2\text{O})_6)^{3+}$  is considered the most phytotoxic. In contrast, Al-F, Al- $\text{SO}_4$ , and Al-P forms are less toxic/non-toxic to plants. Aluminium-tolerant species store Al-organic acid complexes in the symplast (Ma et al. 2001b). Phosphorus treatments along with aluminium increase root and shoot growth and enhance phosphorous absorption in tea plants (Konishi 1992). Al induced the increase in the activities of antioxidant enzymes, such as superoxide dismutase (SOD), catalase (CAT), and ascorbate peroxidase (APX), resulting in increased membrane integrity and delayed lignification and ageing, thus stimulating the growth of tea (*Camelia sinensis* L.) plants (Ghanati et al. 2005; Table. 1). In non-accumulator plants such as turnip (*Brassica rapa* L. sub. sp. *Campestris*) and (*Glycine max* Merr.) increasing aluminium concentrations increased root length (Rufty et al. 1995). At low concentrations,  $\text{Al}^{3+}$  ameliorates  $\text{H}^+$  ion toxicity in wheat (*Triticum aestivum* L.) (Kinraide et al. 1992). Considering the beneficial effects of Al on plant growth indicated by limited studies, its involvement in countering the effects of abiotic stress situations needs to be explored.




Figure 2. Overview of the functions of beneficial elements at various concentrations.

### Cobalt

Cobalt ( $\text{Co}^{2+}$ ) is a micronutrient in plants, and a constituent of vitamin  $\text{B}_{12}$ . Cobalt has similar chemical properties to nickel ( $\text{Ni}^{2+}$ ), and enters plasma membrane through the same carriers. On the basis of plant dry weight, cobalt concentration ranges from 0.1 to 10  $\text{mg kg}^{-1}$  with the exception of plants related to families like *Lamiaceae*, *Scrophulariaceae*, *Asteraceae*, and *Fabaceae*, which are co-hyperaccumulators containing >1000  $\text{mg kg}^{-1}$  cobalt per plant. Of these, *Haumaniastrum robertii* (Robyns) J. Duvign. & Plancke (a copper-tolerant plant) has a peculiarly highest cobalt concentration among phanerogams (about 4304  $\text{mg kg}^{-1}$  dry weight) (Brooks 1977). Numerous studies on cobalt estimation in diverse plant species found the presence of cobalt in different plant parts, i.e. in the style and stigma of *Lilium longifolium* Thunb., leaves of green beans (*Phaseolus sativus* L.) and radish (*Raphanus sativus* L.), and in leafy plants such as lettuce (*Lactuca sativa* L.), cabbage (*Brassica oleracea* var. capitata L.), and spinach (*Spinacea oleracea* L.) (above 0.6  $\text{mg kg}^{-1}$ ) (Kloke 1980). Cereals contain 2.2  $\text{mg Co kg}^{-1}$  and forage plants 0.6–3.5  $\text{mg Co kg}^{-1}$ , rice (*Oryza sativa* L.) contains 0.02–0.150  $\text{mg Co kg}^{-1}$  plant dry weight (Palit et al. 1994).

The essentiality of  $\text{Co}^{2+}$  has not been recorded in the literature, but is required as a constituent of vitamin  $\text{B}_{12}$  into methyl and adenosyl vitamin  $\text{B}_{12}$ , which function as coenzymes. Cobalt is not found at active sites of respiratory chain enzymes; however, there are two sites of action of cobalt discovered in mitochondrial respiration. It prompts different responses towards dissimilar substances like  $\alpha$ -ketoglutarate and succinate. In higher plants,  $\text{Co}^{2+}$  plays a major physiological role, i.e. nitrogen fixation by leguminous crops. Cobalt is an essential component of cobalamine, which is needed for activities of several enzymes in nitrogen fixation by rhizobia bacteria and cyanobacteria that live in root nodules of leguminous plants. In pea (*Pisum sativum* L.), doses of 8  $\text{mg}$  of cobalt to the soil increased growth, plant nutrient levels, nodule numbers and weight, and seed pod yield and quality (Gad 2006). In sweet potato (*Ipomea batatas* L.), 10  $\text{mg}$  cobalt had a collegial effect on root growth, yield quality as starch, sugars, L-ascorbic acid, and contents of  $\text{N}^+$ ,  $\text{P}^{+3}$ ,  $\text{K}^+$ ,  $\text{Mn}^{2+}$ ,  $\text{Zn}^{2+}$  and  $\text{Cu}^{2+}$  compared to the control and other supplemented concentrations (Gad & Kandil 2008). Additionally, Gad (2012) found that supplementation of 8  $\text{mg}$  cobalt to groundnut (*Arachis hypogaea* L.) plants significantly increased nitrogenase activity and subsequently enhanced growth and yield, leading to improved quality of pods and oil yield.

A study on peas (*Pisum sativum* L.) revealed that three cultivars varied in their horticultural characteristics when primed with a combination of ascorbic acid (2  $\text{mg L}^{-1}$ ) +  $\text{CoCl}_2$  (150  $\text{mg L}^{-1}$ );

**Table 1.** Role of beneficial elements in activating antioxidant enzymes to counteract stress conditions.

Beneficial element	Plant species	Antioxidant enzymes	Stress type	Concentration	Reference
Aluminium	Tea ( <i>Camellia sinensis</i> L.)	Activity of superoxide dismutase (SOD), catalase (CAT), and ascorbate peroxidase (APX) enzymes increases	Improves growth	50 $\mu\text{M}$ as aluminium chloride	Ghanati et al. (2005)
Cobalt	Tomato ( <i>Lycopersicon esculentum</i> Mill.)	Activity of superoxide dismutase (SOD) and catalase (CAT) enzymes increases	Improves growth	20 mM as cobaltous sulphate	Zhu et al. (2000)
Selenium	Clover ( <i>Trifolium repens</i> L.)	Alleviates ascorbate peroxidase (APOX) and glutathione reductase (GR) activity	Drought stress	5 $\mu\text{M}$ Se as sodium selenate	Wang (2011)
	Rapeseed ( <i>Brassica napus</i> L.)	Increase the activity of dehydroascorbate reductase (DHAR), glutathione S-transferase (GST), glutathione peroxidase (GPX), and glyoxalase I (Gly I)	Drought stress	25 $\mu\text{M}$ Se as sodium selenate	Hasanuzzaman and Fujita (2011)
	Rapeseed Seedlings ( <i>Brassica napus</i> L.)	Enhances antioxidative defence and MG detoxification systems	Salt stress	25 $\mu\text{M}$ Se as sodium selenate	Hasanuzzaman et al. (2011)
	Ryegrass roots ( <i>Lolium</i> L.)	Enhancing spontaneous dismutation of superoxide radicals and the subsequent activation of POD enzyme	Salt + Heavy metal stress (Al)	0–10 $\mu\text{M}$ Se as sodium selenite	Cartes et al. (2010)
	Spring barley ( <i>Hordeum vulgare</i> L. cv. Rihane-03)	Increases CAT and GSH-Px activities and lowers lipid peroxidation	Drought stress	30 g $\text{ha}^{-1}$ Se as sodium selenate	Habibi (2013)
	Sorghum seeds ( <i>Sorghum bicolor</i> L.)	Improves ascorbic acid peroxidase and guaiacol peroxidases activity	Chilling stress	0, 3, 6 & 12 $\text{mgL}^{-1}$ Se as sodium selenate	Abbas (2012)
	Sorghum leaves ( <i>Sorghum bicolor</i> (L.) Moench)	Enhancing antioxidant defence	High temperature stress	75 $\text{mgL}^{-1}$ Se as sodium selenate	Djanaguiraman et al. (2010)
	Wheat ( <i>Triticum aestivum</i> L.)	Enhances antioxidant defence	Chilling stress	5 $\text{mgL}^{-1}$ Se	Akladios (2012)
	Wheat ( <i>Triticum aestivum</i> L.)	Enhances anthocyanin, phenolic compounds, flavonoids, peroxidases and catalase activity	Chilling stress	1.0 $\text{mg kg}^{-1}$ Se	Chu et al. (2010)
Silicon	Barley ( <i>Hordeum vulgare</i> L.) cultivars, i.e. Kepin No. 7 (salt sensitive) and Jian 4 (salt tolerant),	Higher activity of superoxide dismutase (SOD), peroxidase (POD), catalase (CAT) and glutathione reductase (GR).	Salt stress	1 $\text{mM L}^{-1}$ Si as potassium silicate ( $\text{K}_2\text{SiO}_3$ )	Liang et al. (2003)
	Cotton ( <i>Gossypium hirsutum</i> L.)	Si lowers Ni uptake and enhance antioxidant enzymes activities	Nickel toxicity	1.0 mM Si as sodium silicate ( $\text{Na}_2\text{SiO}_3$ )	Khaliq et al. (2015)
	Okra ( <i>Abelmoschus esculentus</i> L.)	Enhance the activity of superoxide dismutase, peroxidase and catalase	Salt stress	150 $\text{mgL}^{-1}$ Si as silicic acid ( $\text{Si(OH)}_4$ )	Abbas et al. (2015)
	Rice ( <i>Oryza sativa</i> L.)	Enzymatic antioxidant (catalase, peroxidase and polyphenol oxidase) increased	Salt stress	1mM Si as sodium silicate ( $\text{Na}_2\text{SiO}_3 \cdot 5\text{H}_2\text{O}$ )	Kim et al. (2014)
	Tomato ( <i>Lycopersicon esculentum</i> Mill. c.v.)	Raising SOD and CAT activities	Salt stress	2.5 mM Si as sodium silicate ( $\text{Na}_2\text{SiO}_3$ )	Al-aghabary et al. (2005)
	Wheat ( <i>Triticum aestivum</i> L.)	Improved leaf membrane stability index, epicuticular wax, relative water content and proline level	Drought stress	Silicic acid ( $\text{Si(OH)}_4$ ; 29.16% Si), sodium silicate ( $\text{Na}_2\text{SiO}_3$ ; 22.95% Si) and silica gel ( $\text{SiO(OH)}_2$ ; 45.16% Si)	Ahmed et al. (2015)



with desirable significant increases in N and protein content when compared to non-treated plants (Gheeth et al. 2013). A high concentration of cobalt might be attributed to catalase and peroxidase activities that increase catabolism rather than anabolism in tomatoes (Zhu et al. 2000; Table. 1). Cobalt helps to delay senescence in apples and keeps the fruit fresh (Bulantseva et al. 2001).  $\text{Co}^{2+}$  delays ageing in marigold (*Tagetes patula* L.) and chrysanthemum (*Chrysanthemum* spp.), maiden-hair fern (*Adiantum* spp.), common vetch (*Vicia* spp.), and rose (*Rosa* spp.) thereby extending vase and shelf life (Mitra 2015; Figure 2). Cobalt at high doses suppresses  $\text{Cd}^{2+}$  uptake by roots; moreover, cobalt acts synergistically with  $\text{Zn}^{2+}$  and  $\text{Cr}^{2+}$  (Palit et al. 1994). The interaction of cobalt with other metals mainly depends on the concentration of other metal ions. More studies are required to test the role of  $\text{Co}^{2+}$  in plants growing in stressed environments.

### Selenium

Selenium (Se) is a group VIA metalloid in the periodic table whose properties are intermediate between those of sulphur and tellurium. In cultivated soils, selenate and selenite are the most prevalent forms. Selenite is more toxic than selenate, due to its faster incorporation. Depending on the geographical area, Se content ranges from 0.1 to 2.0  $\text{mg kg}^{-1}$ . The minimal nutritional level of Se for animals is about 0.05–0.10  $\text{mg kg}^{-1}$  dry forage feed, while 2–5  $\text{mg kg}^{-1}$  is toxic (Dhillon & Dhillon 2003).

Se is a beneficial micronutrient for animals and humans, yet its essentiality in higher plants is ambiguous and unresolved. Se is essential for growth of some algae (*Chlamydomonas reinhardtii* P. A.Dang) and bacteria but not for higher plants. It is also a structural component of some specific selenoproteins and seleno-tRNAs. Selenocystiene acts as an opal stop codon in selenoproteins when present in a specific secondary mRNA structure (SeCys insertion sequence) (Lee et al. 1989). However, in higher plants, SeCys insertion sequences have not been identified, rather glutathione peroxidase (GPX) (a plant homologous of selenoproteins) replaced Cys with SeCys at their active sites (Lobanov et al. 2007).

Se may be an essential micronutrient for plants endemic to seleniferous soils. *Astragalus* and *Stanleya* are major hyperaccumulators of Se. However, the beneficial effects of Se on Se-hyper-accumulators is much less when plants were grown under lower phosphate levels, which suggests that Se may antagonize phosphate toxicity in Se hyperaccumulator plants (Terry et al. 2000).

In Se non-accumulator plants, Se essentiality is questionable (Marschner 2012). Despite this, Se has not been validated as an essential element; numerous studies have demonstrated that Se may exert beneficial effects at their low concentration, including increased growth in ryegrass, lettuce, potato, and buckweed (Hartikainen 2005). Se has diverse biological roles at different concentrations: trace amounts are needed for normal growth and development; moderate concentrations maintain homeostatic functions, while elevated concentrations have toxic effects on plants. The sensitivity of each plant species to Se concentrations in the soil varies, agriculturally important plant species when exposed to varied Se levels may respond accordingly.

**Protective effects of selenium in counteracting various abiotic stresses.** Selenium helps to ameliorate various abiotic stress injuries induced in plants by cold, drought, high temperature, water, salinity, heavy metals, UV-B stress, senescence, and desiccation stress. Selenium protects plants against abiotic stresses by (1) helping to maintain ion balance and structural integrity of cells, hence regulating the uptake and redistribution of elements essential in the antioxidative system (Table1), and (2) interfering with the electron transport complex (ETC) of the photosynthetic system (Kaur et al. 2014; Figure 2).

Se affects plants by (1) stimulating plant growth and protecting plants against abiotic stresses and heavy metal stresses at low dosage, and (2) acting as a pro-oxidant, which is toxic to plants at high doses. Hence, the concentrations of Se are either beneficial or toxic depending on the crop type. In ryegrass (*Lolium perenne* L.), 1  $\text{mg kg}^{-1}$  Se ( $\text{H}_2\text{SeO}_4$ ) added to soil was marginally toxic, while a similar level (0.8  $\text{mg L}^{-1}$  Se as  $\text{Na}_2\text{SeO}_3$ ) was more toxic in paddy-rice in a hydroponic

system (Feng et al. 2013). Se application as a foliar spray or base fertilizer increased Se content in the edible portion of crops, thus helping to combat injuries generated by four different abiotic stresses (Pezzarossa et al. 2012).

### Silicon

Silicon does not qualify as an essential element but provides substantial benefits to many crops particularly those growing in greenhouse. Silicon mimics carbon on the basis of chemical properties, and can make four bonds with molecule/molecules. Silicon is available to plants as monosilicic ( $\text{Si}(\text{OH})_4$ ) acid with 0.1–0.6 mM in soil and water (Epstein & Bloom 2005), and is supplied to agricultural crops in the form of potassium silicate ( $\text{K}_2\text{SiO}_4$ ) or sodium silicate ( $\text{Na}_2\text{SiO}_4$ ). In plants, silicon is deposited in cell walls in the form of amorphous silica ( $\text{SiO}_2 \cdot n\text{H}_2\text{O}$ ) and enhances cell wall rigidity and strength interacting with pectins and polyphenols. Monocotyledonous plants (10–15%) contain more  $\text{Si}^{4+}$  levels compared with dicotyledonous plants (0.5% or less), especially members of the commelinids clade orders *Arecales* and *Poales* (Hodson et al. 2005). As  $\text{Si}^{4+}$  deficiency affects plant growth, development, and reproduction, it may be classified as a 'quasi essential' element.

Marschner (2012) demonstrated that  $\text{Si}^{4+}$  is deposited in epidermal cells of leaves, hence improving leaf exposure to light by keeping leaves more erect; in roots, it increases cell elongation thus enhancing cell wall elasticity.  $\text{Si}^{4+}$  helps to alleviate various abiotic stresses such as metal toxicity, drought, high temperature, salt, radiation damage, freezing, and chilling. The reduction in dry weight of 30-day-old rice seedlings when exposed to different doses of X-rays was less in those treated with  $\text{Si}^{4+}$  than those without  $\text{Si}^{4+}$  (Ma 2004). Likewise, when  $\text{Si}^{4+}$  was supplied, subsequent radiation exposure restored plant growth.

$\text{Si}^{4+}$  mitigates water scarcity by lowering the transpiration rate, since  $\text{Si}^{4+}$  is deposited under the cuticle of leaves.  $\text{Si}^{4+}$ -supplemented rice seedlings had 30% lower transpiration rates than controls, which has a thin cuticle. Substantial  $\text{Si}^{4+}$  was deposited on epidermal cells in cell walls and caused partial hindrance to transpiration bypass flow. The effect of  $\text{Si}^{4+}$  is more evident under water-stressed conditions (low humidity) than non-water-stressed conditions (high humidity) (Ma et al. 2001a).  $\text{Si}^{4+}$  thermally stabilizes lipids in cell membranes and helps to prevent heat stress.  $\text{Si}^{4+}$ -supplemented leaves had less electrolyte leakage at high temperature (42.5°C) than those without  $\text{Si}^{4+}$ . The effect of salt stress in rice, wheat, and barley was mitigated by silicon application (Ma 2004). NaCl (100mM) applications for 3 weeks suppressed root and shoot growth by 60%, but  $\text{Si}^{4+}$  addition nullified the salt-induced injury in rice plants (Yeo et al. 1999). In barley,  $\text{Si}^{4+}$  may assist changes in structural integrity and function of plasma membranes by suppressing lipid peroxidation caused by salt stress and stimulating root  $\text{H}^+$ -ATPases in membranes and antioxidant machinery (Liang et al. 2002; Rizwan et al. 2015; Figure 2; Table. 1).

### Sodium

Sodium ( $\text{Na}^+$ ) nutrition remains an alluring and evasive topic. While most research has been conducted on sodium toxicity; there are various deliberations and controversies, which suggest that  $\text{Na}^+$  is a beneficial element at low concentrations. As  $\text{K}^+$  is the principal electrolyte for plants,  $\text{Na}^+$  and  $\text{K}^+$  are similar both chemically and structurally. In addition, pervasive use of  $\text{Na}^+$  by numerous salt-tolerant species suggests that  $\text{Na}^+$  can replace  $\text{K}^+$  via non-selective cation transporters. The role of  $\text{Na}^+$  in plant nutrition is based upon a plant's ability to accumulate or eliminate this ion in tissues of glycophytic plant species. However, in halophytic plants, specifically, sodium transporters exclude  $\text{Na}^+$  ions out of roots, into the vacuole or shoot phloem to counteract high sodium levels in the soil. Non-halophytes can use  $\text{Na}^+$  to some extent under  $\text{K}^+$  inadequacy (Subbarao et al. 2000; Figure 2). In  $\text{C}_4/\text{CAM}$  plants,  $\text{Na}^+$  acts as an essential element for regeneration of PEP (phosphoenolpyruvate) from pyruvate to fix carbon for photosynthesis. Sodium acts as a beneficial element in some natrophilic species by replacing potassium ions as co-factors for certain enzymes and also acts as an osmoregulator in stomatal movement and cell expansion, which is particularly beneficial under  $\text{K}^+$ -limiting levels.  $\text{Na}^+$  application to plants of family poaceae,



brassicaceae, apiaceae, asteraceae, malvaceae, fabaceae, and solanaceae enhanced growth in potassium-deficient soils (Flowers & Lauchli 1983; Figure 2). Moreover, the estimated extent to which  $\text{Na}^+$  replaced  $\text{K}^+$  in edible portions of wheat (*Triticum aestivum* L.) and rice (*Oryza sativa* L.) was by 1% and in red beet (*Beta vulgaris* L.) by 90% (Subbarao et al. 2000).


According to the Arnon and Stout (1939) criteria, for  $\text{C}_4$  plant species such as saltbrush (*Atriplex vesicaria* Heward. and *A. tricolor* Heward.), Bassia (*Kochia childsii*), millet (*Panicum miliaceum* L.), and saltgrass (*Distichlis spicata* L.),  $\text{Na}^+$  is essential. Indeed, in  $\text{Na}^+$ -lacking soils,  $\text{C}_4$  plants (*Amaranthaceae*, *Chenopodiaceae*, and *Cyperaceae*) show visual deficiency symptoms like chlorosis and necrosis, grow poorly, and have impaired flower formation; nonetheless, supplementation of 100  $\mu\text{M}$   $\text{Na}^+$  enhanced growth and alleviated visual deficiency symptoms (Johnston et al. 1988). Maathuis (2014) opined that sodium is beneficial but not essential for halophytes and is required at micronutrient levels. Halophytes, particularly succulent halophytes from the *Chenopodiaceae* family, respond to moderate-to-high salinity and reach maximum biomass. Conversely, in other halophytic plant species, growth is stimulated at low sodium levels, which imply that the overall nutritional requirement of the plant is fulfilled.

Some aquatic halophytes use  $\text{Na}^+$  to facilitate nitrate uptake via  $\text{Na}/\text{NO}_3$  co-transporters. Sodium may also interfere in the nitrate and ammonium uptake from soil, thereby impairing nitrogen metabolism (Ullrich 2002). Nitrate is a significant vacuolar solute in many plant species, especially those in the *Chenopodiaceae* and *Graminae* families, under non-saline conditions. Sodium assists in maintaining osmotic balance in plants by synthesizing large amounts of nitrogen-containing compatible solutes like glycinebetaine and proline (to some extent) that leads to the accumulation of amides and polyamines (Pilon-Smits et al. 2009). Although there is considerable influential evidence on the role of  $\text{Na}^+$  in plant metabolism and growth stimulation, the essentiality of  $\text{Na}^+$  in glycophytic plants is under debate.

### Functional relevance against biotic stresses

Plants are the easy targets of regular attempts of attack by biotic stress factors such as fungi, viruses, and herbivores. In response, diverse defence strategies are activated by plants against various invaders, which require the interaction of a susceptible host and a virulent pathogen. In relation to availability of diverse levels of beneficial elements, a range of complex interactions can be expected, taking into account that pathogens, herbivores, and plants also differ in deficiency and toxicity thresholds for such elements. Beneficial metal ions are capable of protecting plants from biotic stresses in various ways: a) 'metal defence' in metal ion-hyperaccumulators by acting either as antifeedants or as plant-systemic pesticides, b) 'trade-off' of organic defences, c) 'metal therapy' by switching a defective signalling system, or d) 'metal-induced fortification' of plants against pathogen attack (Poschenrieder et al. 2006).

Exogenous application of beneficial elements can generate defence reactions in response to pathogen attack in non-accumulator plants. Protection is only possible if the metal is not as much toxic to the plant than to the parasite, i.e. the plant is more resistant to the metal than the pathogen/herbivore. Moreover, a fundamental difference in the relevance of stress signalling between biotic and ion toxicity stress depends on the effectiveness of inducible defence responses. In biotic stress resistance, pathogen elicitor-induced ion fluxes and reactive oxygen species (ROS) are essential for triggering activation of genes responsible for the synthesis of compounds such as phytoalexins that proficiently stop the pathogen, which is the origin of the signal. In the first line of defence, beneficial elements (in low concentration) also act as scavengers for excess of ROS, which lower cell injury induced by the plant pathogen. Secondly, ROS species can result in cell wall lignification that serves as a barrier for pathogen invasion. ROS may also trigger defence signals and defence-related secondary metabolites (Walters et al. 2005). Signal transduction pathways of biotic stress present multiple spots of interaction with metal ion stress signalling (Figure 3). Biotic stress leads to increased



**Figure 3.** The role of beneficial metal ions in plant–pathogen interactions at a metabolic level (Poschenrieder et al. 2006). The shikimate pathway and sulphur metabolism produce compounds that assist in both biotic stress defence and metal binding. An increased level of reduced glutathione is required for stress resistance over a broad range. A shikimate pathway-derivative growth regulator (auxin) makes a connection between stress metabolism and adaptive growth.

cytoplasmic  $\text{Ca}^{2+}$ , ROS production, nitric oxide (NO), salicylate, thioredoxin and mitogen-activated protein kinase (MAPK) interactions between pathogen-elicited reactions and those activated by metal ions (Glazebrook 2005). In biotic stress resistance, pathogen elicitors result in ion fluxes and oxidative burst, which are crucial for prompt activation of genes responsible for the synthesis of compounds like phytoalexins that competently stop the pathogen, which is the origin of the signal (Walters et al. 2005). However, metal ion-induced ROS activates antioxidant production, which assists to limit ROS-derived injury, but does not abolish the stress factor.

Biotic stresses and metal ions can interact at a metabolic level. High levels of reduced glutathione and production of phytochelatins is the key route for interactions between the metal ion and biotic stresses in sulphur metabolism (Figure 3). The shikimate pathway occurring in the host cell is another significant way for interactions between biotic stress and beneficial elements (Figure 3). In this pathway, beneficial ions trigger the production of some defence-related molecules such as essential amino acids; phenolics such as salicylic acid directly concerned in stress signalling; catechin (high-affinity ligand for metal ion) under neutral pH conditions; and flavonoid-type phenolics are released into the rhizosphere, which act as plant signals to  $\text{N}_2$ -fixing symbiotic microorganisms (Barceló & Poschenrieder 2002). In shikimate pathway, benzoic acid 2-hydroxylase (BA2H) is synthesized in leaf vacuoles of cereals, which catalyzes the hydroxylation of benzoic acid to salicylic acid, a mediator of plant responses to pathogens and wounding, which require iron and aluminium for binding. Benzoxazinoids like DIMBOA (2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one) have a high affinity for iron. Moreover, aluminium bound to DIMBOA is not toxic. Upon Al application to the host plant (*Zea mays* L.), enhanced levels of DIMBOA were observed in root tips of an Al-resistant maize variety, while a drastic Al-induced decrease of DIMBOA levels was found in tips of a sensitive variety (Poschenrieder et al. 2005). The indolyl pathway (Figure 3) functional in the host cells presents an extraordinary metabolic range that reflects interactions between plants and insects, predominantly in the case of reactions catalyzed by cytochrome  $\text{P}_{450}$ -type monooxygenase (Chapple 1998). It produces glucosinolates, cyclic hydroxamates, which may be responsible for regulation of interactions among biotic stress and metal ions.

Pathogen stimulates the elicitors such as salicylic acid, benzothiadiazole, benzoic acid, glucosinolates, auxins, cyclic hydroxamates and so forth, which affect production of phenolic compounds and activation of various defence-related enzymes in host cell (Figure 3). In plants, beneficial elements at their low concentrations trigger the antioxidants, metal chelators, antifeedants, antibiotics, phytoalexins, and various growth regulators through a signalling cascade, which impart resistance against biotic stresses (Poschenrieder et al. 2006). Additionally, in response to metal ions, a variety of S-containing secondary metabolites such as thionins and defensins, glucosinolates, and phytoalexins are synthesized, having a key role in defence against pathogen and herbivore attack. Also, the low-molecular-weight thiol glutathione (GSH), evoked by beneficial elements, plays a significant role in response to biotic stresses as part of the glutathione cycle in detoxification of reactive oxygen species (Figure 3).

Limited work has been done regarding the role in beneficial elements in biotic stresses; some case studies are presented further:

### Aluminium

Aluminium helps in disease control because it can be toxic to pathogenic microorganisms. It inhibits spore germination and vegetative growth in the blast rot pathogen (*Thielaviopsis basicola* Ferraris.) (Meyer et al. 1994), and consequently, mycelial growth and sporangial germination of late blight in potato (*Phytophthora infestans*), thus saving it from the disease (Andrison 1995; Table. 2).

**Table 2.** Beneficial elements which help to suppress insect/pathogen attack in various crop plants.

Beneficial element	Plant species	Insect/Pathogen	Reference
Aluminium	Potato ( <i>Solanum tuberosum</i> L.)	<i>Phytophthora infestans</i> (Late blight of potato) <i>Thielaviopsis basicola</i> . Ferraris (Blast rot)	Andrison (1995) Meyer et al. (1994)
Selenium	Desert princes plume ( <i>Stanleya pinnata</i> Pursh.) Indian mustard ( <i>Brassica juncea</i> L.)	<i>Plutella xylostella</i> (Diamond back moth)  <i>Pieris rapae</i> (Caterpillar), <i>Mesodon ferrissi</i> (Snail), <i>Fusarium</i> sp. (Root stem pathogen) <i>Alternaria brassicicola</i> (Leaf pathogen) <i>Myzus persicae</i> (Green peach aphid)	Freeman et al. (2006)  Hanson et al. (2003, 2004)
Silicon	Cucumber ( <i>Cucumis sativus</i> L.)  Grape ( <i>Vitis vinifera</i> L.)  Italian ryegrass ( <i>Lolium multiflorum</i> Lam.) Maize ( <i>Zea mays</i> L.) Pea ( <i>Pisum sativum</i> L.) Rice ( <i>Oryza sativa</i> L.)  Sorghum ( <i>Sorghum bicolor</i> Moench.) Sugarcane ( <i>Saccharum officinarum</i> L.)  Wheat ( <i>Triticum aestivum</i> L.) Wild rice ( <i>Zizania aquatica</i> L.)	<i>Didymella bryoniae</i> <i>Botrytis cineria</i> (Stem rotting & Stem lesions) <i>Pythium ultimum</i> <i>Pythium aphanidermatum</i> (Root disease) <i>Uncinula necator</i> <i>Oidium tuckeri</i> (Powdery mildew) <i>Oscinella frit</i> (Fruit fly) <i>Oscinella frit</i> (Fruit fly)  <i>Sesamia calamistis</i> (Pink stalk borer) <i>Mycosphaerella pinodes</i> (Leaf spot) <i>Magnaportha grisea</i> (Leaf and neck blast) <i>Xanthomonas oryzae</i> pv. <i>oryzae</i> (Xoo) (Bacterial blight) <i>B. graminis</i> f.sp. <i>tritici</i> . <i>M. grisea</i> (Blast disease) <i>Rhizoctonia solani</i> (Sheath blight pathogen) <i>Striga asiatica</i> Kuntze (Witch weed)  <i>Leptosphaeria sacchari</i> (Sugarcane ring spot)  <i>Eldana saccharira</i> (Sugarcane borer) <i>Tribotium castaneum</i> (Flour beetle) <i>Bipolaris oryzae</i> (Fungal brown spot)	O'Neill (1991)  Cherif et al. (1994)  Bowen et al. (1992)  Sang-Young et al. (1996) Moore (1984)  Setamou et al. (1993) Dann and Muir (2002) Seebold et al. (2001) Chang et al. (2002)  Liang et al. (2005) Zhang et al. (2006) Hodson and Sangster (1989)  Raid et al. (1992)  Meyer and Keeping (2001) Setamou et al. (1993) Malvick and Percich (1993)

### Selenium

Selenium (Se)-hyperaccumulating plants such as *Stanleya pinnata* Pursh. (*Brassicaceae*) accumulate more than 1000 mg Se kg<sup>-1</sup> DW from soils that contain 10 mg Se kg<sup>-1</sup> DW or less (Feist & Parker 2001). Se protects *S. pinnata* from caterpillar herbivory (a newly discovered variety of the invasive diamondback moth (*Plutella xylostella*) through deterrence and toxicity). More interestingly, a Se-tolerant wasp (*Diadegma insulare*) can parasitize the tolerant moth (*Plutella xylostella*) (Freeman et al. 2006; Table. 2). In *Brassica juncea* L., Se accumulation defends plants from the *Brassica*-specific herbivore *Pieris rapae* (caterpillar), *Mesodon ferrissi* (snail), *Myzus persicae* (green peach aphids), and fungal infection (Hanson et al. 2003, 2004; Table. 2).

### Silicon

Silicon (Si<sup>4+</sup>) acts as a signal and renders the natural defence from herbivores by instigating enzyme activity such as peroxidases, chitinases, and polyphenoloxidases, or by enhancing the release of phenolic compounds, phytoalexins, antimicrobial compounds, and systemic stress signals (salicylic acid, jasmonic acid, and ethylene). Si<sup>4+</sup> is polymerized in cell walls providing sturdiness, which obstructs fungal germ tubes from penetrating the epidermis and therefore prevents infection (Hayasaka et al. 2008). Si<sup>4+</sup> bioactivity has been correlated with the secondary messengers of systemic acquired resistance (SAR), i.e. analogous to a plant's immune system, reported in cucumber (Kauss et al. 2003). Si<sup>4+</sup> has an equivalent saturable effect and can modulate the activity of the post-lectin intracellular signalling system comprising mitogen-activated protein (MAP) kinases. Fauteux et al. (2005) reported that when Si<sup>4+</sup> invariably accumulates in plants, it is only the soluble forms of Si<sup>4+</sup> within plants that can provide the SAR defence response, while the polymerized form of Si<sup>4+</sup> is almost inert. Furthermore, Si<sup>4+</sup> is an obstacle for insect pests like stem borer, brown plant hopper, rice green leaf hopper, and white-backed plant hopper, and non-insect pests such as leaf spider and mites (Savant et al. 1997; Table. 2).

### Conclusions

We conclude that adequate intracellular concentrations of beneficial metal ions (in traces) are not only required for optimal growth and development of plants and their enemies, but also for pathogen virulence and plant defences. Hyperaccumulators of beneficial elements are the principal basis for the metal defence hypothesis against plant pathogens. Cross-talk in stress signalling and metal ions is related to plant responses against both abiotic and biotic stress factors and is an upcoming research area in metal-hyperaccumulator and non-hyperaccumulator plants. Elements such as aluminium (Al), cobalt (Co), sodium (Na), selenium (Se) and silicon (Si) are considered beneficial for plants. Na and Si are vital for certain plant groups, while Al and Se are recommended as crucial for some hyperaccumulator species. Co is essential for the microbial partners of some plants rather than the plants themselves. In this era of research, the effects of beneficial elements at low levels deserve more awareness in order to fertilize crops with these nutrients to boost crop production under stressed environments as well as enhance plant nutritional value as feed or food. The mechanisms following the beneficial effects (at low levels) to the plants have not been fully explored, unlike the toxic effects (at high levels). How these elements impart protection from pathogens and abiotic stress factors need to be probed further, especially at a molecular level. Whether these elements have synergistic or antagonistic effects on plants growing under unstressed and stressed situations requires to be investigated. Foliar application of these elements needs to be tested in plants growing under stressed environment. It is noteworthy that elements other than those discussed here may also be beneficial for plants, but more validation is needed to support these results.

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

The first and second authors thank the Government of India for providing fellowships under the University Grants Commission and minority fellowship grants to undertake this work.

## References

- Abbas SM. 2012. Effects of low temperature and selenium application on growth and the physiological changes in sorghum seedlings. *J Stress Physiol Biochem*. 8:268–286.
- Abbas T, Balal RM, Shahid MA, Pervez MA, Ayyub CM, Aqueel MA, Javaid MM. 2015. Silicon-induced alleviation of NaCl toxicity in okra (*Abelmoschus esculentus*) is associated with enhanced photosynthesis, osmoprotectants and antioxidant metabolism. *Acta Physiol Plant*. 37:1–15.
- Ahmed M, Qadeer U, Ahmed ZI, Hassan FU. 2015. Improvement of wheat (*Triticum aestivum*) drought tolerance by seed priming with silicon. *Arch Agron Soil Sci*. 1–17. doi:10.1080/03650340.2015.1048235
- Akladios SA. 2012. Influence of different soaking times with selenium on growth, metabolic activities of wheat seedlings under low temperature stress. *Afr J Biotechnol*. 11:14792–14804.
- Al-aghaby K, Zhu Z, Shi Q. 2005. Influence of silicon supply on chlorophyll content, chlorophyll fluorescence, and antioxidative enzyme activities in tomato plants under salt stress. *J Plant Nutr*. 27:2101–2115.
- Andrion D. 1995. Inhibition by aluminum of mycelia growth and of sporangial production and germination in *Phytophthora infestans*. *Eur J Plant Pathol*. 101:527–533.
- Arnon DI, Stout PR. 1939. Experimental methods for the study of the role of copper, manganese, and zinc in the nutrition of higher plants. *Am J Bot*. 26:144–149.
- Barceló J, Poschenrieder C. 2002. Fast root growth responses, root exudates, and internal detoxification as clues to the mechanisms of aluminum toxicity and resistance: a review. *Environ Exp Bot*. 48:75–92.
- Bowen P, Menzies J, Ehret D, Samuel L, Glass ADM. 1992. Soluble silicon sprays inhibit powdery mildew development on grape leaves. *J Am Soc Hortic Sci*. 117:906–912.
- Brooks RS. 1977. Copper, magnesium and cobalt uptake by *Haumanistrum* species. *Plant Soil*. 48:541–544.
- Bulantseva EA, Glinka EM, Protsenko MA, Sal'kova EG. 2001. A protein inhibitor of polygalacturonase in apple fruits treated with aminoethoxyvinylglycine and cobalt chloride. *Prikl Biokhim Mikrobiol*. 37:100–104.
- Cartes P, Jara AA, Pinilla L, Rosas A, Mora ML. 2010. Selenium improves the antioxidant ability against aluminium-induced oxidative stress in ryegrass roots. *Ann Appl Biol*. 156:297–307.
- Chang SJ, Tzeng DDS, Li CC. 2002. Effect of silicon nutrient on bacterial blight resistance of rice (*Oryza sativa* L.). In: Matoh, T, editors., Second Silicon in Agriculture Conference; Press-Net, Kyoto, Japan. p.31–33.
- Chapple C. 1998. Molecular-genetic analysis of plant cytochrome P450-dependent monooxygenases. *Annu Rev Plant Biol*. 49:311–343.
- Cherif M, Menzies JG, Ehret DL, Bogdanoff C, Belanger RR. 1994. Yield of cucumber infected with *Pythium aphanidermatum* when grown with soluble silicon. *Hort Sci*. 29:896–897.
- Chiba Y, Mitani N, Yamaji N, Ma JF. 2009. HvLsi1 is a silicon influx transporter in barley. *Plant J*. 57:810–818.
- Chu X, Yao Z, Zhang ZN. 2010. Responses of wheat seedlings to exogenous selenium supply under cold stress. *Biol Trace Elem Res*. 136:355–363.
- Dann EK, Muir S. 2002. Peas grown in media with elevated plant-available silicon levels have higher activities of chitinases and  $\beta$ -1,3-glucanase, are less susceptible to a fungal leaf spot pathogen and accumulate more foliar silicon. *Aust Plant Pathol*. 31:9–13.
- Davidian JC, Hatzfield Y, Cathala N, Tagmount A, Vidmar JJ. 2000. Sulfate uptake and transport in plants. In: Brunold, C, Rennenberg, H, De, KLJ, Stuhlen, I, Davidian, JC, editors. Sulfur nutrition and sulfur assimilation in higher plants: molecular, biochemical and physiological aspects. Bern: Paul Haupt Verlag; p. 19–40.
- Dhillon KS, Dhillon SK. 2003. Distribution and management of seleniferous soils. *Adv Agron*. 79:119–185.
- Djanaguiraman M, Prasad PVV, Seppänen M. 2010. Selenium protects sorghum leaves from oxidative damage under high temperature stress by enhancing antioxidant defense system. *Plant Physiol Biochem*. 48:999–1007.
- Epstein E, Bloom AJ. 2005. Mineral nutrition of plants: principles and perspectives. Sunderland: Sinauer Associates.
- Fauteux F, Re'mus-Borel W, Menzies JG, Be' langer RR. 2005. Silicon and plant disease resistance against pathogenic fungi. *FEMS Microbiol Lett*. 249:1–6.
- Feist LJ, Parker DR. 2001. Ecotypic variation in selenium accumulation among populations of *Stanleya pinnata*. *New Phytol*. 149:61–69.
- Feng R, Wei C, Tu S. 2013. The roles of selenium in protecting plants against abiotic stresses. *Environ Exp Bot*. 87:58–68.

- Flowers TJ, Lauchli A. 1983. Sodium versus potassium: substitution and compartmentation. *Encyc Plant Physiol.* 15:651–681.
- Freeman JL, Quinn CF, Marcus MA, Fakra S, Pilon-Smits EA. 2006. Selenium-tolerant diamondback moth disarms hyperaccumulator plant defense. *Curr Biol.* 16:2181–2192.
- Gad N. 2006. Increasing the efficiency of nitrogen fertilization through cobalt application to pea plant. *Res J Agr Biol Sci.* 2:433–442.
- Gad N. 2012. Role and importance of cobalt nutrition on groundnut (*Arachis hypogaea*) production. *World Appl Sci J.* 20:359–367.
- Gad N, Kandil H. 2008. Response of sweet potato (*Ipomoea batatas* L.) plants to different levels of cobalt. *Aust J Basic Appl Sci.* 2:945–955.
- Ghanati F, Morita A, Yokota H. 2005. Effects of aluminum on the growth of tea plant and activation of antioxidant system. *Plant Soil.* 276:133–141.
- Gheeth RHM, Moustafa YMM, Abdel-Hakeem WM. 2013. Enhancing growth and increasing yield of peas (*Pisum sativum* L.) by application of ascorbic acid and cobalt chloride. *J Nov Appl Sci.* 2:106–115.
- Glazebrook J. 2005. Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annu Rev Phytopathol.* 43:205–227.
- Habibi G. 2013. Effect of drought stress and selenium spraying on photosynthesis and antioxidant activity of spring barley. *Acta Agric Slov.* 101:31–39.
- Hanson B, Garifullina GF, Lindblom SD, Wangeline A, Ackley A, Kramer K, Pilon-Smits EA. 2003. Selenium accumulation protects *Brassica juncea* from invertebrate herbivory and fungal infection. *New Phytol.* 159:461–469.
- Hanson B, Lindblom SD, Loeffler ML, Pilon-Smits EA. 2004. Selenium protects plants from phloem-feeding aphids due to both deterrence and toxicity. *New Phytol.* 162:655–662.
- Hartikainen H. 2005. Biogeochemistry of selenium and its impact on food chain quality and human health. *J Trace Elem Med Biol.* 18:309–318.
- Hasanuzzaman M, Fujita M. 2011. Selenium pretreatment upregulates the antioxidant defense and methylglyoxal detoxification system and confers enhanced tolerance to drought stress in rapeseed seedlings. *Biol Trace Elem Res.* 143:1758–1776.
- Hasanuzzaman M, Hossain MA, Fujita M. 2011. Selenium-induced up-regulation of the antioxidant defense and methylglyoxal detoxification system reduces salinity-induced damage in rapeseed seedlings. *Biol Trace Elem Res.* 143:1704–1721.
- Hayasaka T, Fujii H, Ishiguro K. 2008. The role of silicon in preventing appressorial penetration by the rice blast fungus. *Phytopathol.* 98:1038–1044.
- Hodson MJ, Evans DE. 1995. Aluminium/silicon interactions in higher plants. *J Exp Bot.* 46:161–171.
- Hodson MJ, Sangster AG. 1989. X-ray microanalysis of the seminal root of *Sorghum bicolor* with particular reference to silicon. *Ann Bot.* 64:659–667.
- Hodson MJ, White PJ, Mead A, Broadley MR. 2005. Phylogenetic variation in the silicon composition of plants. *Ann Bot.* 96:1027–1046.
- Huang JW, Barchelard EP. 1993. Effects of aluminum on growth and cation uptake in seedlings of *Eucalyptus mannifera* and *Pinus radiata*. *Plant Soil.* 149:121–127.
- Jarosick J, Vara PZ, Koneeny J, Obdrzalek M. 1988. Dynamics of cobalt 60 uptake by roots of pea plants. *Sci Total Environ.* 71:225–229.
- Johnston M, Grof CP, Brownell PF. 1988. The effect of sodium nutrition on the pool sizes of intermediates of the C4 photosynthetic pathway. *Aust J Plant Physiol.* 15:749–760.
- Kaur N, Sharma S, Kaur S, Nayyar H. 2014. Selenium in agriculture: a nutrient or contaminant for crops? *Arch Agron Soil Sci.* 60:1593–1624.
- Kauss H, Seehaus K, Franke R, Gilbert S, Dietrich RA, Kroger N. 2003. Silica deposition by a strongly cationic proline-rich protein from systemically resistant cucumber plants. *Plant J.* 33:87–95.
- Khalik A, Ali S, Hameed A, Farooq MA, Farid M, Shakoor MB, Rizwan M. 2015. Silicon alleviates nickel toxicity in cotton seedlings through enhancing growth, photosynthesis, and suppressing Ni uptake and oxidative stress. *Arch Agron Soil Sci.* 1–15. doi:10.1080/03650340.2015.1073263
- Kim YH, Khan AL, Waqas M, Shim JK, Kim DH, Lee KY, Lee IJ. 2014. Silicon application to rice root zone influenced the phytohormonal and antioxidant responses under salinity stress. *J Plant Growth Regul.* 33:137–149.
- Kinraide TB, Ryan PR, Kochian LV. 1992. Interactive effects of  $Al^{3+}$ ,  $H^{+}$ , and other cations on the root elongation considered in terms of cell-surface electrical potential. *Plant Physiol.* 99:1461–1468.
- Kloke A. 1980. Effects of heavy metals on soil fertility, flora and fauna as well as its importance for the nutrition and fodder-chain. *Proceedings of the Utilization of Refuse and Sewage sludge in Agriculture; Zurich*
- Konishi S. 1992. Promotive effects of aluminium on tea plant growth. *JARQ.* 26:26–33.
- Korshunova YO, Eide D, Clarke WG, Guiriot ML, Pakrasi HB. 1999. The IRT1 protein from *Arabidopsis thaliana* is a metal transporter with a broad substrate range. *Plant Mol Biol.* 40:37–44.
- Lee BJ, Worland PJ, Davis JN, Stadtman TC, Hatfield DL. 1989. Identification of a selenocysteylseryl transfer RNA in mammalian cells that recognizes the nonsense codon, UGA. *J Biol Chem.* 264:9724–9727.



- Liang Y, Chen Q, Liu Q, Zhang W, Ding R. 2003. Exogenous silicon (Si) increases antioxidant enzyme activity and reduces lipid peroxidation in roots of salt-stressed barley (*Hordeum vulgare* L.). *J Plant Physiol.* 160:1157–1164.
- Liang Y, Chen Q, Zhang W, Ding R. 2002. Exogenous silicon increases antioxidant enzyme activities and reduces lipid peroxidation in roots of salt-stressed barley (*Hordeum vulgare* L.). Abstract of Second Silicon in Agriculture Conference; 22–26 August; Yamagata. Tsuruoka (Japan): Japanese Society of Soil Science and Plant Nutrition; p. 140–151.
- Liang YC, Sun W, Si J, Römheld V. 2005. Effects of foliar- and root-applied silicon on the enhancement of induced resistance to powdery mildew in *Cucumis sativus*. *Plant Pathol.* 54:678–685.
- Lobanov AV, Fomenko DE, Zhang Y, Sengupta A, Hatfield DL, Gladyshev VN. 2007. Evolutionary dynamics of eukaryotic selenoproteomes: large selenoproteomes may associate with aquatic life and small with terrestrial life. *Genome Biol.* 8:R198.
- Ma JF. 2004. Role of silicon in enhancing the resistance of plants to biotic and abiotic stresses. *Soil Sci Plant Nutr.* 50:11–18.
- Ma JF, Miyake Y, Takahashi E. 2001a. Silicon in Agriculture. Amsterdam: Elsevier Science.
- Ma JF, Ryan PR, Delhaize E. 2001b. Aluminium tolerance in plants and the complexing role of organic acids. *Trends Plant Sci.* 6:273–278.
- Ma JF, Tamai K, Yamaji N, Mitani N, Konishi S, Katsuhara M, Ishiguro M, Murata Y, Yano M. 2006. A silicon transporter in rice. *Nature.* 440:688–691.
- Ma JF, Yamaji N, Mitani N, Tamai K, Konishi S, Fujiwara T, Katsuhara M, Yano M. 2007. An efflux transporter of silicon in rice. *Nature.* 448:209–212.
- Maathuis FJM. 2014. Sodium in plants: perception, signalling, and regulation of sodium fluxes. *J Exp Bot.* 65:849–858.
- Malvick DK, Percich JA. 1993. Hydroponic culture of wild rice (*Zizania palustris* L.) and its application to studies of silicon nutrition and fungal brown spot disease. *Can J Plant Sci.* 73:969–975.
- Marschner H. 2012. Marschner's mineral nutrition of higher plants. London: Academic press.
- Meyer JH, Keeping MG. 2001. Past, present and future research of the role of silicon for sugarcane in southern Africa. *Studies Plant Sci.* 8:257–275.
- Meyer JR, Shew HD, Harrison UJ. 1994. Inhibition of germination and growth of *Thielaviopsis basicola* by aluminum. *Phytopathol.* 84:598–602.
- Mitani N, Chiba Y, Yamaji N, Ma JF. 2009. Identification and characterization of maize and barley Lsi2-like silicon efflux transporters reveals a distinct silicon uptake system from that in rice. *Plant Cell Online.* 21:2133–2142.
- Mitra GN. 2015. Cobalt (Co), Selenium (Se), Vanadium (V), Cadmium (Cd), Lead (Pb) and Titanium (Ti). In: Garrity G, et al., editors. Regulation of Nutrient Uptake by Plants. New Delhi: Springer. pp.189–195.
- Moore D. 1984. The role of silica in protecting Italian ryegrass (*Lolium multiflorum*) from attack by dipterous stem-boring larvae (*Oscinella* fruit and other related species). *Ann Appl Biol.* 104:161–166.
- O'Neill TM. 1991. Investigation of glasshouse structure, growing medium and silicon nutrition as factors affecting disease incidence in cucumber crops. *Med Fac Landbouww Rijksuniv Gent.* 56:359–367.
- Palit S, Sharma A, Talukder G. 1994. Effect of cobalt on plants. *Bot Rev.* 60:149–181.
- Pezzarossa B, Remorini D, Gentile ML, Massai R. 2012. Effects of foliar and fruit addition of sodium selenate on selenium accumulation and fruit quality. *J Sci Food Agric.* 92:781–786.
- Pilon-Smits EA, Quinn CF, Tapken W, Malagoli M, Schiavon M. 2009. Physiological functions of beneficial elements. *Curr Opin Plant Biol.* 12:267–274.
- Poschenrieder C, Tolrà R, Barceló J. 2006. Can metals defend plants against biotic stress? *Trends Plant Sci.* 11:288–295.
- Poschenrieder C, Tolrà RP, Barceló J. 2005. A role for cyclic hydroxamates in aluminium resistance in maize? *J Inorg Biochem.* 99:1830–1836.
- Raid RN, Anderson DL, Ulloa MF. 1992. Influence of cultivar and amendment of soil with calcium silicate slag on foliar disease development and yield of sugarcane. *Crop Prot.* 11:84–88.
- Rizwan M, Ali S, Ibrahim M, Farid M, Adrees M, Bharwana SA, Abbas F. 2015. Mechanisms of silicon-mediated alleviation of drought and salt stress in plants: a review. *Environ Sci Pollut R.* 1–16. doi:10.1007/s11356-015-5305-x
- Ruffy TW, Mackown CT, Lazof DB, Carter TE. 1995. Effects of aluminium on nitrate uptake and assimilation. *Plant Cell Environ.* 18:1325–1331.
- Sang-Young N, Kyong MK, Sang CL, Jong CP. 1996. Effects of lime and silica fertilizer application on grape cracking. *J Agric Sci Soil Fert.* 38:410–415.
- Savant NK, Snyder GH, Datnoff LE. 1997. Silicon management and sustainable rice production. *Adj' Agron.* 58:151–199.
- Seebold KW, Kucharek TA, Datnoff LE, Correa-Victoria FJ, Marchetti MA. 2001. The influence of silicon on components of resistance to blast in susceptible, partially resistant, and resistant cultivars of rice. *Phytopathol.* 91:63–69.
- Setamou M, Schulthess F, Bosque-Perez NA, Thomas-Odjo A. 1993. Effect of plant nitrogen and silicon on the biotomcs of *Sesamia calamistics*. *Bull Ent Res.* 83:405–411.
- Silva P, Gerós H. 2009. Regulation by salt of vacuolar H<sup>+</sup>-ATPase and H<sup>+</sup> pyrophosphatase activities and Na<sup>+</sup>/H<sup>+</sup> exchange. *Plant Signal Behav.* 4:718–726.

- Smith FW, Hawkesford MJ, Ealing PM, Clarkson DT, Berg PJV. 1997. Regulation of expression of a cDNA from barley roots encoding a high affinity sulphate transporter. *Plant J.* 12:875–884.
- Smith FW, Hawkesford MJ, Prosser IM, Clarkson DT. 1995. Isolation of a Cdna from *Saccharomyces cerevisiae* that encodes a high affinity sulphate transporter at the plasma membrane. *Mol Gen Genet.* 247:709–715.
- Subbarao GV, Wheeler RM, Stutte GW. 2000. Feasibility of substituting sodium for potassium in crop plants for advanced life support systems. *Life Supp Biosph Sci.* 7:225–232.
- Terry N, Zayed AM, De Souza MP, Tarun AS. 2000. Selenium in higher plants. *Ann Rev Plant Physiol Plant Mol Biol.* 51:401–432.
- Ullrich WR. 2002. Salinity and nitrogen nutrition. In: Läuchli, A, Lüttge, U, editors. *Salinity: Environment-Plants-Molecules*. Boston: Kluwer Academic Publishers; p. 229–248.
- Walters D, Walsh D, Newton A, Lyon G. 2005. Induced resistance for plant disease control: maximizing the efficacy of resistance elicitors. *Phytopathol.* 95:1368–1373.
- Wang CQ. 2011. Water-stress mitigation by selenium in *Trifolium repens* L. *J Plant Nutri Soil Sci.* 174:276–282.
- Yeo AR, Flowers SA, Rao G, Welfare K, Senanayake N, Flowers TJ. 1999. Silicon reduces sodium uptake in rice (*Oryza sativa* L.) in saline conditions and this is accounted for by a reduction in the transpirational bypass flow. *Plant Cell Environ.* 22:559–565.
- Zhang GL, Dai QG, Zhang HC. 2006. Silicon application enhances resistance to sheath blight (*Rhizoctonia solani*) in rice. *J Plant Physiol Mol Biol.* 32:600–606.
- Zhu H, Shipp E, Sanchez RJ, Liba A, Stine JE, Hart PJ, Gralla EB, Nersissian AM, Valentine JS. 2000. Cobalt (2) binding to tomato copper chaperone for superoxide dismutase: implications for the metal ion transfer mechanism. *Biochemistry.* 39:5413–5542.