

Name, institution, and contact details of the applicant (one person only) and a list of the scientists or institutions involved in the research project (max. 2 A4 pages) CVs and lists of publications need not be submitted.

PD Dr. rer. medic. Susanne Weis, Heinrich Heine University Düsseldorf;
Gruppenleiterin „Variabilität des Gehirns“, Gehirn und Verhalten (INM-7), Institut für Neurowissenschaften und Medizin, Forschungszentrum Jülich;
E-Mail: S.Weis@fz-juelich.de

Simon

INM-7: The proposed work is embedded in a multidisciplinary working team combining knowledge in the field of neuropsychology, structural and functional MRI analysis, computational neuroscience and machine learning.

Dr. Kaustubh Patil, head of the working group “Applied Machine Learning”

Rick Betzel is an associate professor at the Psychological and Brain Sciences Department of Psychological and Brain Sciences at Indiana University Bloomington, USA. He is an expert in principles of time-varying functional network reconfiguration and its relationship to ongoing cognitive processes and one of the developers of the edge time series approach which we will employ within the project suggested here.

Dynamic Cognition: Movies as a Window into Sex Differences in the Brain

Project Description

Background and Research Question

Functional brain imaging techniques, in particular fMRI have been widely used to investigate sex differences in the brain. Such differences in structural and functional brain organization play a crucial role in healthy human brain development, aging, as well as the manifestation of psychiatric and neurological disorders [1, 3]. Consequently, an understanding of sex differences in the human brain and their underlying mechanisms is critical for understanding both normative behavior and psychopathology.

In the present proposal, we aim to employ novel brain imaging methodology utilizing naturalistic viewing (NV), i.e. watching of Hollywood movies clips in the scanner, to broaden our knowledge of sex differences in the brain. While it is common knowledge that women and men often react differently to complex scenarios in films - think of how a couple might debate a movie's emotional impact after leaving the cinema - our study delves deeper than these everyday observations. Instead of relying on stereotypes such as 'women like emotional scenes, men like action', we will analyze in detail how brain activity and functional connectivity (FC) differs when women and men watch various movie scenes. This could reveal, for example, that women process subtle social cues in dialogues differently from men, while men might react more strongly to visual hints in action scenes that foreshadow danger. Through a thorough analysis of females' and males' brain responses to a variety of different movie scenes, we aim to develop a well-founded understanding of cognitive sex differences that advances our knowledge of sex differences in brain function beyond the current state of research.

So far, our understanding of sex differences in the brain is still far from complete. While there is some agreement on the existence of sex differences in specific cognitive domains like language and spatial processing, some researchers argue that female and male brains are altogether more similar than different [4] and that the overlap between the sexes is larger than their differences. Importantly, neuroimaging research on sex difference has so far only been able to capture certain aspects of these differences.

Classically, sex differences in the brain have been studied using task-based (TB) fMRI with tightly controlled tasks, offering insights limited to specific cognitive domains (e.g. [9, 11, 12]). However, due to the highly controlled and artificial nature of the tasks, ecological validity of task-related fMRI is usually very low and does not reflect cognitive sex differences as observed in daily life. More recently, resting state (RS) approaches have been employed, in which fMRI data is acquired while subjects relax in the scanner without any specific task demand or visual or auditory stimulation. Earlier RS studies examined group differences in connectivity patterns between women and men. More recently, however, machine learning (ML) methods have been applied to move beyond group averages: sex classification approaches use RS data to predict the sex of individual subjects and then infer which brain networks contribute most to distinguishing females from males. Using such an ML approach, we identified regionally specific brain networks that support successful classification, which, importantly, generalized from a training sample to an independent sample and revealed that predictive power—and thus cognitive sex differences—are most pronounced in higher-level cognitive regions involved in language, social cognition, and emotion processing [13, 15, 14].

Importantly though, RS fMRI only reflects the brain in a specific state, which has commonly interpreted as intrinsic brain organization and thus constitutes a trait rather than a state. Moreover, while a state of free mind wandering happens often in real life, what remains largely unexplored are sex differences in the "brain in action", i.e. when encountering complex, multimodal stimulation resembling real-life experiences. The present proposal aims to contribute to closing this gap in knowledge by applying the newly emerging NV approach to examine sex-specific brain responses in ecologically valid contexts. NV focuses on cognitive processes in dynamic, temporally extended, naturalistic contexts, which are much more akin to situations which the brain must deal with in real life. In NV tasks, participants in the scanner are presented with naturalistic material like movies. For the participants, there is no other task demand rather than watching the clips. NV offers an engaging task for the subject and thus avoids boredom in the scanner. Importantly, as opposed to RS, all participants are exposed to the same stimulus, for which content and timing is known and can be used in the analyses. NV approaches offer complex, dynamic and ongoing stimulation similar to experiences in everyday situations, where low-level (audiovisual) and high-level (cognitive and emotional) content




Fig. 1: ROI-based sex classification shows high accuracies for (a) within sample CV and (b) across sample classification.

vary fluidly, creating a multimodal and immersive experience [8], offering the opportunity to capture dynamic neural processing in ecologically valid contexts [10]. Furthermore, NV has been shown to improve reliability and individual identifiability over RS [5] with a recent study from our lab [5] showing that the use of NV enhances the detection of FC patterns that are unique at the individual level.

Our lab pioneered NV analysis methods. We recently developed the topography-based predictive framework (TOPF) [7], which identifies individual-specific evoked activity topographies in a data-driven manner and examines their behavioural relevance using a ML predictive framework. TOPF effectively and stably captures individual differences in evoked brain activity and successfully predicts phenotypes across cognition, emotion and personality, and will be employed in the present context to uncover differences between females and males. In preliminary work using TOPF [7] on NV data, we achieved up to 80% accuracy in sex classification on single movie clips. Highly predictive regions were linked to emotion, language, and higher cognition. These promising findings await further validation over the course of the present project. Altogether, NV offers rich, time-evolving stimulation, allowing for the study of dynamic brain states across diverse scenarios. This method better reflects everyday cognitive and emotional challenges and may uncover novel sex differences in functional brain organization.

Surprisingly, to our knowledge, no study has yet used NV to systematically examine sex differences. In a previous publication (Eickhoff, et al. 2020), we have compared the potential of movies for the study of individual brain differences to what running on a treadmill is to the heart during a cardiac stress-test, i.e. to potentially provide a standardized way to study the whole organ while it works to compare function across different levels of intensity and demands. In the present context, this means that NV can be expected to reveal sex differences in the brain that so far have been unobservable, since NV data encompass an extensive range of dynamic real-life situations, such as social interactions, recognition of faces or perception of emotions [8]. While isolated cognitive processes as examined in TB fMRI might not reveal significant sex differences, the complex interplay of these processes - as engaged during movie watching - could highlight more pronounced differences between women and men. This underscores the value of using naturalistic stimuli to uncover subtle but meaningful variations in neural processing that might be missed in more controlled, simplified experimental paradigms. Using careful annotations of the movies, it is also possible to identify which features of a given scene are particularly relevant to evoking sex differences in complex experiences and to identify brain networks that differ most between females and males during these experiences.

We aim to employ novel methodology utilizing NV paradigms to expand our understanding of sex differences in the brain. This approach has the potential to elucidate sex-specific neural mechanisms in response to the complex and dynamic content of movies and to reveal nuanced sex-specific distinctions in brain function that may not be apparent in more constrained experimental paradigms or everyday observations. Our methodology leverages the ecological validity of movie stimuli to probe complex, multimodal cognitive processes in a controlled yet naturalistic setting. Through advanced neuroimaging techniques and sophisticated data analysis methods, we anticipate uncovering subtle yet significant sex-based variations in neural activity and FC that can inform our understanding of broader cognitive and behavioural differences between females and males. To this aim, we will first consider temporally aggregated FC evoked by movies over the duration of several minutes. Subsequently, we will further disentangle which specific events evoke sex differences and examine differences in the brain networks caused by such events. Furthermore, we will examine how the brain's response to the evolving narrative of the movies differs between the sexes. The project's comprehensive approach of examining temporally aggregated as well as time resolved FC, network dynamics and brain activity patterns provides a multi-faceted view of sex differences in the brain and, which can have implications for understanding sex differences in cognition, behavior, and susceptibility to certain neurological and psychiatric conditions.

Results from the proposed project have the potential to add a crucial new dimension to current knowledge on cognitive sex differences. Psychologically, these insights deepen our understanding of normative sex differences; clinically, they may help clarify why certain psychiatric and neurological disorders manifest differently in women and men. Such knowledge can potentially enhance diagnostic accuracy, inform personalized treatments, and support the development of sex-specific healthcare and education strategies.




Fig. 2: TOPF captures individual differences in brain activity and successfully predicts phenotypes across cognition, emotion and personality.

Throughout the proposal, for ease of reading, the term “sex” will refer to the self-reported (biological) sex and appearance of the participants. At the same time, we fully acknowledge that “gender identity”, i.e. the subjective identification of an individual as female, male, or one of the other gender identities which might be also fluid, also play important roles, which however, are beyond the scope of the present project.

Data: The Ju-MOVIES dataset

The work proposed here will be based on a data set that has already been acquired and is ideally suited for the examination of sex differences through a NV approach. Over the course of the project, further data will be collected. The paradigm comprises seven movie clips of about 8 - 10 minutes length, which are all excerpts from Hollywood movies (“Dirty Dancing”, “Scream”, “Dead Poets Society”, “Forrest Gump”, “Dead Man Walking”, “Life is Beautiful”, “The Good, the Bad, the Ugly”). The specific movie stimuli were chosen to cover a variety of social interactions and complex situations as well as multifaceted emotions evolving over the course of the movie to match the complex nature of real-life experiences. Considering that viewers lack the context of the entire movie, each stimulus was chosen to be long enough for the viewers to understand the situation, empathize with characters and follow the evolving narrative. Additionally, participants watch a compilation of 12 short movie clips (“Short Sequences”), which were again taken from Hollywood movies and completed two RS scans of about 9 minutes length with their eyes open.

So far, we collected data of 135 healthy participants (68 males, age 18 - 35 years, mean age = 24.33 years). All fMRI data are acquired on a Siemens 3T Prisma scanning (Siemens, Erlangen, Germany) with a 64-channel head coil at the Imaging Core Facility of Research Centre Jülich, using a T2w multiband echo planar imaging sequence with the following parameters: repetition time (TR) = 980ms, echo time (TE) = 30ms, flip angle = 70°, field of view (FOV) = 207 x 207mm, voxel size=2.2 x 2.2 x 2.0mm³, number of slices: 64, multiband acceleration factor=4, phase encoding direction=AP, FoV=207mm). A mirror fixed on the head coil allows participants to see a screen used to display the movies. In-ear headphones are used for ear protection and to deliver the movie sound. Additionally, a structural T1w image is acquired using an MP-RAGE sequence (TR=2000ms, TE=2.45ms, TI=900ms, flip angle=8°, FoV: 256mm) yielding 1mm³ voxels.

Saliva samples are collected from each participant and levels of cortisol, oestradiol, progesterone, and testosterone are analysed by a specialized laboratory. Control for fluctuating hormone levels is essential for a meaningfully interpretation of potential sex differences in functional brain. Furthermore, data OC intake is noted for female participants.

To characterize the movie stimuli in more detail, emotions perceived by viewers of the movies were assessed by 44 German participants (23 males, age 20-30 years, mean age 25.05 years) who watched all movie stimuli while simultaneously rating six basic emotions (happiness, fear, surprise, sadness, disgust and anger [2]). Rating were collected by the ReMoTa toolbox (Real-time Movie Tagging v0.0; [6]). Participants watched the movies on a laptop and rated their feelings by use of the keyboard. All emotions were rated simultaneously on a scale from 0 to 100 at a sampling rate of 10hz. The emotion ratings confirmed the multifaceted and diverse emotions elicited by the different movie stimuli. In a further annotation, two independent raters quantified the following features: faces, bodies, male / female presenting characters, ethnicity of characters, presence of children, adults, crowds, hands, buildings, vehicles, food, landscapes, animals, plants, movement, social interactions, place (inside or outside / urban vs. non-urban), time of day (day or night), weather, presence of music and camera movements.

References

- [1] Larry Cahill. “Why sex matters for neuroscience”. In: *Nature Reviews. Neuroscience* 7.6 (June 2006), pp. 477–484. ISSN: 1471-003X. DOI: 10.1038/nrn1909.
- [2] P. Ekman and W. V. Friesen. “Constants across cultures in the face and emotion”. In: *Journal of Personality and Social Psychology* 17.2 (Feb. 1971), pp. 124–129. ISSN: 0022-3514. DOI: 10.1037/h0030377.
- [3] Aarthi R. Gobinath, Elena Choleris, and Liisa A. M. Galea. “Sex, hormones, and genotype interact to influence psychiatric disease, treatment, and behavioral research”. In: *Journal of Neuroscience Research* 95.1 (Jan. 2, 2017), pp. 50–64. ISSN: 1097-4547. DOI: 10.1002/jnr.23872.
- [4] Daphna Joel et al. “Sex beyond the genitalia: The human brain mosaic”. In: *Proceedings of the National Academy of Sciences of the United States of America* 112.50 (Dec. 15, 2015), pp. 15468–15473. ISSN: 1091-6490. DOI: 10.1073/pnas.1509654112.
- [5] Jean-Philippe Kröll et al. “Naturalistic viewing increases individual identifiability based on connectivity within functional brain networks”. In: *NeuroImage* 273 (June 2023), p. 120083. ISSN: 1095-9572. DOI: 10.1016/j.neuroimage.2023.120083.

- [6] Giada Lettieri et al. “Emotionotopy in the human right temporo-parietal cortex”. In: *Nature Communications* 10.1 (Dec. 5, 2019), p. 5568. ISSN: 2041-1723. DOI: 10.1038/s41467-019-13599-z.
- [7] Xuan Li et al. “A topography-based predictive framework for naturalistic viewing fMRI”. In: *NeuroImage* 277 (Aug. 15, 2023), p. 120245. ISSN: 1095-9572. DOI: 10.1016/j.neuroimage.2023.120245.
- [8] Saurabh Sonkusare, Michael Breakspear, and Christine Guo. “Naturalistic Stimuli in Neuroscience: Critically Acclaimed”. In: *Trends in Cognitive Sciences* 23.8 (Aug. 2019), pp. 699–714. ISSN: 1879-307X. DOI: 10.1016/j.tics.2019.05.004.
- [9] M. Thimm et al. “Menstrual cycle effects on selective attention and its underlying cortical networks”. In: *Neuroscience* 258 (Jan. 31, 2014), pp. 307–317. ISSN: 1873-7544. DOI: 10.1016/j.neuroscience.2013.11.010.
- [10] Tamara Vanderwal, Jeffrey Eilbott, and F. Xavier Castellanos. “Movies in the magnet: Naturalistic paradigms in developmental functional neuroimaging”. In: *Developmental Cognitive Neuroscience* 36 (Apr. 2019), p. 100600. ISSN: 1878-9307. DOI: 10.1016/j.dcn.2018.10.004.
- [11] Susanne Weis et al. “Dynamic changes in functional cerebral connectivity of spatial cognition during the menstrual cycle”. In: *Human Brain Mapping* 32.10 (Oct. 2011), pp. 1544–1556. ISSN: 1097-0193. DOI: 10.1002/hbm.21126.
- [12] Susanne Weis et al. “Estradiol modulates functional brain organization during the menstrual cycle: an analysis of interhemispheric inhibition”. In: *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 28.50 (Dec. 10, 2008), pp. 13401–13410. ISSN: 1529-2401. DOI: 10.1523/JNEUROSCI.4392-08.2008.
- [13] Susanne Weis et al. “Sex Classification by Resting State Brain Connectivity”. In: *Cerebral Cortex (New York, N.Y.: 1991)* 30.2 (Mar. 21, 2020), pp. 824–835. ISSN: 1460-2199. DOI: 10.1093/cercor/bhz129.
- [14] Lisa Wiersch and Susanne Weis. “Sex differences in the brain: More than just male or female”. In: *Cognitive Neuroscience* 12.3 (2021), pp. 187–188. ISSN: 1758-8936. DOI: 10.1080/17588928.2020.1867084.
- [15] Lisa Wiersch et al. “Accurate sex prediction of cisgender and transgender individuals without brain size bias”. In: *Scientific Reports* 13.1 (Aug. 24, 2023), p. 13868. ISSN: 2045-2322. DOI: 10.1038/s41598-023-37508-z.