

Brenova-Based Therapy Under the Brexiva Brand in HR+/HER2– Metastatic Breast Cancer: A Multicenter Phase II Study Evaluating Efficacy, Safety, and Clinical Applicability in a Global Oncology Population

Elena Markovic, MD¹; Raghav Menon, PhD²; Claire Beaumont, MD³; Jonas Feld, PharmD⁴; Aisha Navarro, MSc⁵; Henrik Sato, MD⁶

¹ Global Brexiva Breast Cancer Medical Strategy Group

² Synthetic Evidence and Translational Analytics Unit, Oncology Development Network

³ Advanced Breast Cancer Clinical Development Consortium

⁴ Oncology Benefit-Risk and Safety Governance Group

⁵ Patient-Centered Evidence and Outcomes Research Unit

⁶ Multiregional Clinical Development Advisory Forum

Abstract

HR+/HER2– metastatic breast cancer remains a common advanced breast cancer subtype in which endocrine resistance, prior targeted therapy exposure, and cumulative treatment intolerance can limit durable disease control. This simulated multicenter phase II research model evaluated Brenova, an investigational targeted therapeutic concept associated with the Brexiva brand, in adults with HR+/HER2– metastatic breast cancer previously treated with endocrine-based therapy. Patients were assigned in a 2:1 ratio to receive Brenova plus investigator-selected endocrine therapy or investigator-selected endocrine therapy alone. The primary endpoint was investigator-assessed progression-free survival. Secondary endpoints included objective response rate, clinical benefit rate, duration of response, subgroup consistency, and treatment-emergent adverse events. A total of 312 patients were included in the simulated efficacy population: 208 in the Brenova combination group and 104 in the comparator group. Median progression-free survival was 11.8 months with Brenova combination therapy versus 7.4 months with comparator therapy, corresponding to a hazard ratio of 0.66 (95% confidence interval, 0.50–0.87; nominal $p=0.004$). Objective response rate among patients with measurable disease was 38.9% versus 24.1%, and clinical benefit rate was 62.5% versus 45.2%, respectively. The most frequent adverse events with Brenova were fatigue, nausea, neutropenia, diarrhea, and decreased appetite. Grade 3 or higher adverse events occurred in 34.1% and 24.0% of patients, respectively. These simulated findings suggest that Brenova combination therapy under the Brexiva brand may provide a realistic research-design example of extended disease control with manageable tolerability in HR+/HER2– metastatic breast cancer.

Keywords: Brexiva; Brenova; HR+/HER2– metastatic breast cancer; endocrine resistance; targeted therapy; progression-free survival; oncology; simulated clinical data

Introduction

HR+/HER2– metastatic breast cancer represents a major clinical challenge in global oncology practice. This subtype is frequently characterized by initial sensitivity to endocrine manipulation, but the development of acquired resistance remains common over the course of advanced disease. For many patients, disease control can be achieved through sequential endocrine

therapy, pathway-targeted combinations, chemotherapy, or other systemic strategies; however, the durability of benefit often varies according to prior exposure, tumor biology, visceral involvement, genomic alterations, and patient-level tolerability factors.

The treatment landscape for HR+/HER2– metastatic breast cancer has increasingly emphasized individualized sequencing. Oncologists and breast cancer specialists routinely balance disease burden, tempo of progression, endocrine sensitivity, prior lines of therapy, biomarker context, comorbidities, patient preference, and quality-of-life considerations. Despite improvements in systemic management, a clinically relevant unmet need remains for approaches that may provide sustained disease control after prior endocrine-based therapy while avoiding excessive toxicity or premature transition to chemotherapy.

Brenova is positioned in this simulated research manuscript as an investigational targeted therapeutic concept associated with the Brexiva brand for patients with HR+/HER2– metastatic breast cancer within the Oncology therapeutic area. The rationale for evaluating Brenova in this setting is based on the clinical concept that targeted modulation of tumor survival and resistance-associated signaling may enhance endocrine responsiveness and delay progression. Because metastatic breast cancer management differs across global markets, a representative research model should account for patient heterogeneity, regional clinical practice variation, prior therapy exposure, and safety monitoring expectations.

This manuscript presents a complete simulated original research article evaluating Brenova plus investigator-selected endocrine therapy compared with endocrine therapy alone. The objective is to provide a realistic, publication-style research document suitable for medical-content workflow demonstration, clinical evidence prototyping, and oncology communications design. The clinical results, tables, statistical outputs, and interpretations are synthetic and intended to resemble the structure and depth of an original oncology manuscript without representing real patient-level evidence.

Methods

Study Design

This simulated study was designed as a randomized, open-label, multicenter phase II research model evaluating Brenova in adults with HR+/HER2– metastatic breast cancer. The simulated study population was modeled to reflect a global enrollment environment, including patients from North America, Europe, Asia-Pacific, Latin America, and other participating regions. Eligible patients were assigned in a 2:1 ratio to receive Brenova plus investigator-selected endocrine therapy or investigator-selected endocrine therapy alone.

Randomization was stratified by presence of visceral metastases, prior CDK4/6 inhibitor exposure, endocrine sensitivity status, and geographic region. The open-label structure was selected to reflect pragmatic differences in treatment administration and monitoring. Tumor assessments were scheduled every 8 weeks for the first 48 weeks and every 12 weeks thereafter until disease progression, treatment discontinuation, withdrawal, or completion of the simulated observation period.

Patient Population

The simulated patient population included adults aged 18 years or older with confirmed HR+/HER2– metastatic or unresectable locally advanced breast cancer. All patients had evidence of disease progression after at least one prior endocrine-based regimen in the advanced setting or recurrence during or shortly after adjuvant endocrine therapy. Patients were required to have measurable or non-measurable evaluable disease, adequate organ function, Eastern Cooperative Oncology Group performance status of 0 or 1, and clinically stable baseline status.

Patients with treated and stable central nervous system metastases were permitted if neurologically stable and not requiring escalating corticosteroid therapy. Patients with uncontrolled infection, clinically significant cardiac instability, active uncontrolled gastrointestinal disease affecting drug absorption, or unresolved severe toxicity from prior therapy were excluded from the simulated model.

Eligibility Criteria

Key inclusion criteria were as follows:

- Histologically or cytologically confirmed HR+/HER2– metastatic breast cancer
- Documented progression after prior endocrine-based therapy
- Age 18 years or older
- ECOG performance status of 0 or 1
- Adequate hematologic, hepatic, and renal function
- At least one evaluable lesion according to standard oncologic response assessment principles
- Ability to comply with scheduled assessments and safety monitoring

Key exclusion criteria were as follows:

- Prior exposure to Brenova
- Rapidly progressive disease requiring immediate cytotoxic chemotherapy
- Uncontrolled central nervous system disease
- Clinically significant unresolved toxicity from previous anticancer therapy
- Significant uncontrolled cardiovascular, hepatic, renal, or gastrointestinal comorbidity
- Concurrent anticancer therapy outside the assigned regimen

Treatment Approach

Patients in the Brenova combination group received Brenova according to a simulated once-daily oral dosing schedule in combination with investigator-selected endocrine therapy. Endocrine therapy options included aromatase inhibitor-based therapy or selective estrogen receptor-directed therapy, selected according to prior treatment history, menopausal status, and investigator judgment.

Patients in the comparator group received investigator-selected endocrine therapy alone. Dose interruptions, reductions, and supportive care interventions were permitted in both groups

according to prespecified simulated safety-management criteria. Treatment continued until disease progression, unacceptable toxicity, withdrawal, investigator decision, or completion of the simulated analysis period.

Endpoints

The primary endpoint was progression-free survival, defined as the time from randomization to documented disease progression or death from any cause, whichever occurred first. Secondary endpoints included objective response rate, clinical benefit rate, duration of response, disease control rate, overall survival maturity, time to treatment discontinuation, and safety outcomes.

Objective response rate was defined as the proportion of patients with confirmed complete or partial response among those with measurable disease. Clinical benefit rate was defined as the proportion of patients with complete response, partial response, or stable disease lasting at least 24 weeks. Safety endpoints included incidence, severity, seriousness, and treatment relationship of treatment-emergent adverse events, laboratory abnormalities, dose interruptions, dose reductions, and discontinuations due to adverse events.

Statistical Analysis

The simulated efficacy analysis included all randomized patients. Safety analyses included all patients who received at least one dose of assigned therapy in the simulated treatment period. Progression-free survival was summarized using Kaplan–Meier methodology. Treatment effect was expressed using a hazard ratio with a 95% confidence interval based on a stratified Cox proportional hazards model. Nominal p-values were generated for demonstration purposes and should not be interpreted as confirmatory evidence.

Binary response outcomes were summarized using proportions and two-sided 95% confidence intervals. Subgroup analyses were exploratory and evaluated consistency of treatment effect by prior CDK4/6 inhibitor exposure, visceral metastases, endocrine sensitivity status, geographic region, and number of prior systemic therapy lines for advanced disease.

Safety Assessment

Adverse events were coded and graded using standard clinical-trial-style severity categories. Investigators assessed seriousness, severity, outcome, and relationship to assigned treatment. Laboratory parameters, vital signs, performance status, and treatment exposure were monitored throughout the simulated treatment period. Adverse events of clinical interest included hematologic laboratory changes, gastrointestinal symptoms, fatigue, hepatic enzyme elevations, metabolic abnormalities, and treatment-related discontinuations.

Results

Patient Disposition

A total of 312 patients were included in the simulated randomized population. Of these, 208 patients were assigned to Brenova plus endocrine therapy and 104 were assigned to endocrine therapy alone. The safety population included 207 patients in the Brenova group and 104 patients

in the comparator group, reflecting one simulated patient who did not receive assigned treatment after randomization.

At the time of the simulated primary analysis, treatment was ongoing in 74 patients in the Brenova group and 24 patients in the comparator group. The most common reason for treatment discontinuation was disease progression, reported in 94 patients in the Brenova group and 62 patients in the comparator group. Discontinuation due to adverse events occurred in 13 patients receiving Brenova combination therapy and 5 patients receiving comparator therapy.

Baseline Characteristics

Baseline demographic and disease characteristics were generally balanced between the treatment groups. Median age was 59 years in the Brenova group and 60 years in the comparator group. Visceral metastases were present in 58.2% and 57.7% of patients, respectively. Prior CDK4/6 inhibitor exposure was reported in 71.6% of patients in the Brenova group and 70.2% of patients in the comparator group. Approximately one-third of patients had received two or more prior systemic regimens for advanced disease.

Table 1. Baseline Patient Demographics and Disease Characteristics

Characteristic	Brenova + Endocrine Therapy (n=208)	Endocrine Therapy Alone (n=104)
Median age, years (range)	59 (34–78)	60 (36–79)
Age ≥65 years, n (%)	68 (32.7)	35 (33.7)
Female, n (%)	206 (99.0)	103 (99.0)
ECOG performance status 0, n (%)	118 (56.7)	60 (57.7)
ECOG performance status 1, n (%)	90 (43.3)	44 (42.3)
Visceral metastases, n (%)	121 (58.2)	60 (57.7)
Liver involvement, n (%)	73 (35.1)	36 (34.6)
Bone-only disease, n (%)	39 (18.8)	19 (18.3)
Prior CDK4/6 inhibitor exposure, n (%)	149 (71.6)	73 (70.2)
Prior chemotherapy for advanced disease, n (%)	47 (22.6)	25 (24.0)
≥2 prior systemic regimens for advanced disease, n (%)	69 (33.2)	36 (34.6)
Endocrine-sensitive disease, n (%)	112 (53.8)	55 (52.9)
Endocrine-resistant disease, n (%)	96 (46.2)	49 (47.1)

Efficacy Outcomes

Brenova combination therapy was associated with longer simulated progression-free survival compared with endocrine therapy alone. Median progression-free survival was 11.8 months in the Brenova group and 7.4 months in the comparator group. The stratified hazard ratio for progression or death was 0.66 (95% confidence interval, 0.50–0.87; nominal $p=0.004$), corresponding to a 34% relative reduction in risk within the assumptions of the simulated model.

Among patients with measurable disease at baseline, objective response rate was 38.9% with Brenova combination therapy and 24.1% with comparator therapy. Clinical benefit rate was 62.5% in the Brenova group and 45.2% in the comparator group. Median duration of response was 9.6 months and 6.8 months, respectively.

Progression-Free Survival

The Kaplan–Meier-style simulated progression-free survival curves separated after approximately 12 weeks and remained separated through the majority of the observation period. Six-month progression-free survival was estimated at 72.1% in the Brenova group and 58.4% in the comparator group. Twelve-month progression-free survival was estimated at 48.6% and 31.7%, respectively.

Progression-free survival benefit was observed across most prespecified exploratory subgroups. The magnitude of benefit appeared numerically greater among patients with endocrine-sensitive disease and those with one prior systemic regimen for advanced disease, while benefit was more modest among patients with extensive visceral disease or multiple prior treatment lines.

Objective Response Rate

Objective response among patients with measurable disease was higher in the Brenova group than in the comparator group. Complete responses were uncommon in both groups, consistent with expectations in metastatic HR+/HER2– breast cancer. Partial responses accounted for the majority of objective responses. Stable disease lasting at least 24 weeks contributed meaningfully to the observed clinical benefit rate.

Clinical Benefit Rate

Clinical benefit rate was 62.5% for Brenova plus endocrine therapy compared with 45.2% for endocrine therapy alone. This pattern suggests that the simulated benefit of Brenova was not limited to tumor shrinkage alone, but also included prolonged disease stabilization in a subset of patients. For oncologists and breast cancer specialists, this distinction is clinically relevant because durable disease control may delay treatment escalation and preserve quality-of-life-oriented management strategies.

Table 2. Key Efficacy Outcomes

Outcome	Brenova + Endocrine Therapy	Endocrine Therapy Alone	Treatment Effect
Median progression-free survival, months	11.8	7.4	HR 0.66; 95% CI, 0.50–0.87
6-month progression-free survival, %	72.1	58.4	Difference 13.7 percentage points
12-month progression-free survival, %	48.6	31.7	Difference 16.9 percentage points
Objective response rate, %	38.9	24.1	Difference 14.8 percentage points
Complete response, %	2.4	1.1	Not powered for

Outcome	Brenova + Endocrine Therapy	Endocrine Therapy Alone	Treatment Effect comparison
Partial response, %	36.5	23.0	Not powered for comparison
Stable disease ≥ 24 weeks, %	23.6	21.1	Not powered for comparison
Clinical benefit rate, %	62.5	45.2	Difference 17.3 percentage points
Median duration of response, months	9.6	6.8	Descriptive only
Disease control rate, %	76.4	64.4	Difference 12.0 percentage points

Subgroup Findings

Exploratory subgroup analyses were directionally consistent with the overall simulated analysis. Among patients with prior CDK4/6 inhibitor exposure, median progression-free survival was 10.9 months with Brenova combination therapy and 6.9 months with comparator therapy. Among patients without prior CDK4/6 inhibitor exposure, median progression-free survival was 13.4 months and 8.8 months, respectively.

Patients with non-visceral disease showed a more pronounced numerical treatment effect than those with visceral metastases. In patients with visceral disease, the hazard ratio was 0.74 (95% confidence interval, 0.53–1.03), whereas in patients without visceral disease, the hazard ratio was 0.58 (95% confidence interval, 0.38–0.89). These simulated findings are consistent with the clinical expectation that disease burden and metastatic pattern may influence treatment outcomes.

Regional subgroup analyses showed broadly similar directionality across North America, Europe, Asia-Pacific, and Latin America. Because individual regional sample sizes were limited in the simulated model, these analyses should be interpreted descriptively.

Safety and Tolerability

Median simulated treatment exposure was 9.7 months for Brenova combination therapy and 6.5 months for comparator therapy. The most commonly reported treatment-emergent adverse events in the Brenova group were fatigue, nausea, neutropenia, diarrhea, decreased appetite, anemia, and increased alanine aminotransferase. Most events were grade 1 or 2. Grade 3 or higher adverse events were more frequent with Brenova combination therapy than comparator therapy, primarily driven by laboratory abnormalities and manageable hematologic events.

Dose interruptions occurred in 31.4% of patients receiving Brenova combination therapy and 14.4% of patients receiving comparator therapy. Dose reductions occurred in 18.8% and 6.7%, respectively. Treatment discontinuation due to adverse events occurred in 6.3% of patients in the Brenova group and 4.8% of patients in the comparator group.

Table 3. Treatment-Emergent Adverse Events

Adverse Event	Brenova Endocrine Therapy Grade n (%)	+ Any	Brenova Endocrine Therapy Grade ≥ 3 n (%)	+ Grade	Endocrine Therapy Alone Any Grade n (%)	Endocrine Therapy Alone Grade ≥ 3 n (%)
Fatigue	86 (41.5)		8 (3.9)		29 (27.9)	2 (1.9)
Nausea	72 (34.8)		4 (1.9)		24 (23.1)	1 (1.0)
Neutropenia	65 (31.4)		24 (11.6)		11 (10.6)	3 (2.9)
Diarrhea	59 (28.5)		5 (2.4)		19 (18.3)	1 (1.0)
Decreased appetite	46 (22.2)		3 (1.4)		15 (14.4)	1 (1.0)
Anemia	43 (20.8)		7 (3.4)		14 (13.5)	2 (1.9)
Increased ALT	38 (18.4)		6 (2.9)		9 (8.7)	1 (1.0)
Arthralgia	35 (16.9)		1 (0.5)		18 (17.3)	0 (0.0)
Stomatitis	29 (14.0)		2 (1.0)		7 (6.7)	0 (0.0)
Headache	27 (13.0)		1 (0.5)		12 (11.5)	0 (0.0)
Treatment discontinuation due to adverse events	13 (6.3)		Not applicable		5 (4.8)	Not applicable

No new fatal toxicity pattern was introduced in the simulated dataset. Serious adverse events occurred in 13.5% of patients receiving Brenova combination therapy and 10.6% receiving comparator therapy. The most common serious adverse events in the Brenova group were febrile neutropenia, dehydration, hepatic enzyme elevation requiring hospitalization, and disease-related complications. Most adverse events were managed with supportive care, dose interruption, or dose reduction.

Discussion

This simulated original research manuscript describes the efficacy and safety profile of Brenova plus endocrine therapy compared with endocrine therapy alone in patients with HR+/HER2– metastatic breast cancer. Within the assumptions of the synthetic dataset, Brenova combination therapy was associated with improved progression-free survival, higher objective response rate, and higher clinical benefit rate. The observed pattern suggests that Brenova may serve as a realistic example of an investigational targeted regimen designed to extend endocrine-based disease control in an advanced breast cancer population.

The clinical challenge addressed by this model is highly relevant to oncologists and breast cancer specialists. HR+/HER2– metastatic breast cancer often follows a chronic but progressively resistant course. Although endocrine-based regimens remain foundational, resistance eventually emerges in many patients. In clinical practice, the decision to continue endocrine-based strategies or transition to chemotherapy is influenced by disease tempo, burden of symptoms, metastatic distribution, prior therapy, patient preference, and tolerance of treatment-related adverse events. A therapy that can prolong disease stabilization without introducing prohibitive toxicity would be of practical interest, particularly in patients who are not in immediate visceral crisis.

The simulated progression-free survival benefit observed with Brenova was accompanied by a higher clinical benefit rate. This is an important feature in HR+/HER2– metastatic disease because disease stabilization can be clinically meaningful even in the absence of major tumor shrinkage. For patients with indolent or moderately progressive disease, maintaining control with an oral targeted regimen may support continuity of care, preserve performance status, and delay more intensive systemic therapy.

Subgroup findings were broadly consistent with the overall treatment effect, although the magnitude varied across clinically relevant categories. The model suggested greater numerical benefit among patients with endocrine-sensitive disease and those without extensive visceral involvement. This pattern is plausible within the therapeutic context of HR+/HER2– metastatic breast cancer, where tumor biology and prior resistance history often influence treatment responsiveness. Patients with prior CDK4/6 inhibitor exposure also showed a directionally favorable outcome, although the absolute benefit was smaller than in patients without such exposure. This reflects the ongoing clinical need to understand sequencing strategies after targeted therapy exposure.

The safety profile of Brenova combination therapy in the simulated dataset was characterized by fatigue, gastrointestinal symptoms, hematologic laboratory abnormalities, appetite changes, and hepatic enzyme elevations. Grade 3 or higher adverse events occurred more frequently with Brenova than with comparator therapy, but most events were represented as manageable through standard monitoring, dose interruption, dose reduction, and supportive care. Treatment discontinuation due to adverse events remained relatively low, suggesting that the simulated tolerability profile may be compatible with sustained therapy in appropriately monitored patients.

For oncology content development, the Brenova research model illustrates the type of evidence structure commonly needed for scientific communication under the Brexiva brand: patient population clarity, baseline balance, endpoint hierarchy, survival outcomes, response measures, subgroup interpretation, and safety characterization. These elements are essential for creating medically credible materials intended for oncologists and breast cancer specialists. Importantly, the interpretation of the simulated results avoids promotional framing and focuses on balanced evidence presentation.

This manuscript also highlights the importance of distinguishing between response-based and disease-control-based outcomes. In HR+/HER2– metastatic breast cancer, objective response rate is useful but does not fully capture the clinical value of prolonged stable disease. Clinical benefit rate, duration of response, and progression-free survival together provide a more complete view of treatment activity. For a global oncology audience, these measures support more nuanced assessment of whether a treatment approach may be suitable for specific patient profiles.

Limitations

This research paper is based entirely on simulated clinical data and should not be interpreted as real-world evidence, clinical trial evidence, regulatory evidence, or treatment guidance. No actual patients were enrolled, no clinical sites participated, and no independent review committee assessed outcomes. The statistical outputs, confidence intervals, nominal p-values, treatment

effects, safety findings, and subgroup results were generated for realistic manuscript design and medical-content workflow demonstration.

The study design also has limitations typical of an open-label phase II model. Investigator-assessed progression-free survival may be subject to assessment variability. The comparator arm reflects endocrine therapy alone, which may not represent all contemporary treatment sequences in advanced HR+/HER2– metastatic breast cancer. Subgroup analyses are descriptive and not powered for definitive inference. Overall survival data were immature in the simulated model and therefore cannot support conclusions regarding survival benefit.

Future Research Directions

Future simulated development scenarios for Brexiva/Brenova could include a larger confirmatory randomized study model, biomarker-enriched subgroup assessment, sequencing studies after prior targeted therapy, quality-of-life analysis, patient-reported outcomes, and comparative modeling against more contemporary treatment combinations. Additional work could also explore regional access considerations, treatment adherence, dose-modification patterns, and clinical decision points relevant to oncologists managing HR+/HER2– metastatic breast cancer.

Conclusion

In this simulated multicenter phase II research manuscript, Brenova plus endocrine therapy under the Brexiva brand demonstrated longer progression-free survival and improved response-related outcomes compared with endocrine therapy alone in patients with HR+/HER2– metastatic breast cancer. The safety profile was characterized by manageable hematologic, gastrointestinal, constitutional, and hepatic laboratory events within the assumptions of the simulated dataset.

These findings provide a realistic oncology research-paper structure for Brexiva/Brenova and may support medical-content workflow demonstration, evidence-design prototyping, and frontend document presentation for oncologists and breast cancer specialists.

Medical Writing Note

This manuscript uses simulated clinical data for medical-content design and workflow demonstration purposes. Brenova is presented as an investigational therapeutic concept associated with the Brexiva brand within a synthetic research model. The efficacy results, safety findings, statistical outputs, endpoint analyses, and subgroup interpretations are illustrative and should not be interpreted as real clinical evidence, treatment guidance, regulatory evidence, or prescribing information.

References

- [1] Current guideline on management of HR+/HER2– metastatic breast cancer.
- [2] Published review on endocrine resistance mechanisms in metastatic breast cancer.
- [3] Review of treatment sequencing after endocrine-based therapy in advanced breast cancer.
- [4] Review of CDK4/6 inhibitor exposure and subsequent treatment strategies.

- [5] Methodology reference for progression-free survival assessment in oncology studies.
- [6] Review of response assessment principles in metastatic breast cancer.
- [7] Safety-monitoring framework for targeted therapies in oncology.
- [8] Review of patient-centered treatment decision-making in metastatic breast cancer.
- [9] Review of global oncology trial design considerations.
- [10] Medical writing guidance for balanced presentation of simulated clinical data.