

Biomarker Patterns and Endocrine-Resistance Features Associated with Brenova-Based Therapy Under the Brexiva Brand in HR+/HER2- Metastatic Breast Cancer

Marina Volkova, MD¹; Sandeep Iyer, PhD²; Celeste Romano, MD³; Tobias Klein, PharmD⁴; Anika Desai, MSc⁵; Rafael Moretti, MD⁶

¹ Global Brexiva Breast Cancer Medical Strategy Group

² Synthetic Evidence and Translational Analytics Unit, Oncology Development Network

³ Advanced Breast Cancer Translational Research Consortium

⁴ Oncology Benefit-Risk and Safety Governance Group

⁵ Patient-Centered Evidence and Outcomes Research Unit

⁶ Multiregional Clinical Development Advisory Forum

Abstract

HR+/HER2- metastatic breast cancer is biologically heterogeneous, and treatment outcomes are influenced by endocrine sensitivity, prior targeted therapy exposure, metastatic pattern, and resistance-associated biomarker profiles. This simulated translational oncology research model evaluated clinical outcomes and exploratory biomarker patterns among adults receiving Brenova-based therapy under the Brexiva brand for HR+/HER2- metastatic breast cancer. The simulated cohort included 324 patients with available baseline clinical characteristics and biomarker-status categories derived from tissue or circulating tumor DNA-style assessment. Patients received Brenova-based therapy in combination with clinician-selected endocrine therapy. The primary descriptive endpoint was progression-free survival by biomarker subgroup. Secondary endpoints included objective response rate, clinical benefit rate, treatment persistence, and safety outcomes. Median progression-free survival in the overall cohort was 10.9 months. Patients with endocrine-sensitive disease had longer median progression-free survival than those with endocrine-resistant disease, at 13.2 months versus 8.1 months. Patients with low resistance-marker burden had a median progression-free survival of 12.7 months compared with 8.6 months among those with high resistance-marker burden. Objective response rate was 36.8% overall and was higher among patients with lower baseline resistance-marker burden. The most frequent treatment-emergent adverse events were fatigue, nausea, neutropenia, diarrhea, decreased appetite, anemia, and hepatic laboratory abnormalities. These simulated findings provide a realistic research-design example showing how biomarker context, endocrine-resistance features, and clinical outcomes may be integrated into a balanced oncology manuscript for Brenova under the Brexiva brand in HR+/HER2- metastatic breast cancer.

Keywords: Brexiva; Brenova; HR+/HER2- metastatic breast cancer; endocrine resistance; biomarkers; translational oncology; circulating tumor DNA; treatment sequencing; simulated clinical data

Introduction

HR+/HER2- metastatic breast cancer is commonly managed with sequential endocrine-based and targeted systemic therapies. Although many patients achieve disease control with endocrine therapy combinations, resistance eventually develops in a substantial proportion of cases. The clinical course is influenced by tumor biology, prior treatment exposure, metastatic distribution,

endocrine sensitivity, molecular adaptation, and patient-specific tolerability. As a result, the same treatment approach may produce different outcomes across apparently similar patients.

A central challenge in this disease setting is the identification of patients most likely to benefit from continued endocrine-based strategies. In routine oncology practice, clinicians often evaluate factors such as duration of benefit on prior endocrine therapy, presence of visceral disease, pace of progression, prior exposure to targeted agents, performance status, and patient preference. Increasingly, biomarker-informed interpretation is also becoming important. Resistance-associated alterations, pathway activation patterns, and tumor evolution may affect treatment sensitivity and duration of response.

Brenova is presented in this manuscript as an investigational therapeutic concept associated with the Brexiva brand for HR+/HER2– metastatic breast cancer within the Oncology therapeutic area. The clinical rationale for this simulated translational research model is based on the need to understand how a Brenova-based regimen may perform across biomarker-defined and clinically defined subgroups. Rather than evaluating only aggregate outcomes, this manuscript explores how endocrine-resistance features and baseline biomarker burden may influence progression-free survival, response, clinical benefit, and treatment continuation.

This manuscript is designed as a realistic original research paper for medical-content workflow demonstration. It uses synthetic clinical-style data and simulated biomarker categories to demonstrate how translational oncology findings could be structured for oncologists and breast cancer specialists. The manuscript does not represent real clinical evidence and should not be interpreted as treatment guidance.

Methods

Study Design

This was a simulated, global, multicenter translational oncology research model evaluating Brenova-based therapy under the Brexiva brand in patients with HR+/HER2– metastatic breast cancer. The model included clinical outcome data, baseline disease characteristics, endocrine-sensitivity classification, and exploratory biomarker categories. Patients were followed from treatment initiation until disease progression, treatment discontinuation, loss to follow-up within the simulated model, or completion of the analysis period.

The study was structured to resemble a prospective translational cohort analysis embedded within a broader oncology treatment program. All patients received Brenova-based therapy in combination with clinician-selected endocrine therapy. No randomized comparator arm was included. The objective was to describe outcome patterns according to biomarker and endocrine-resistance features rather than to establish comparative treatment efficacy.

Patient Population

Eligible simulated patients were adults with HR+/HER2– metastatic breast cancer who initiated Brenova-based therapy after progression on at least one prior endocrine-based regimen. Patients were required to have evaluable metastatic disease, ECOG performance status of 0 to 2, adequate baseline clinical documentation, and available biomarker-status classification from tissue or circulating tumor DNA-style assessment.

Patients with rapidly progressive disease requiring immediate cytotoxic chemotherapy were excluded from the primary simulated cohort. Patients with stable treated central nervous system disease were included if neurologically stable and able to complete routine follow-up assessments.

Biomarker Assessment Framework

Baseline biomarker information was organized into clinically interpretable categories. The simulated framework included endocrine-resistance marker burden, cell-cycle pathway activation pattern, tumor-suppression alteration status, and detectable circulating tumor DNA-style signal at baseline. These categories were created for manuscript-design demonstration and do not represent validated diagnostic definitions.

Endocrine-resistance marker burden was classified as low, intermediate, or high. Low burden represented limited resistance-associated features, intermediate burden represented one or two resistance-associated features, and high burden represented multiple resistance-associated features or rapidly evolving disease biology. Circulating tumor DNA-style signal was classified as low, moderate, or high based on simulated baseline molecular burden.

Treatment Approach

Patients received Brenova-based therapy with clinician-selected endocrine therapy. The endocrine partner was selected according to prior treatment exposure, menopausal status, metastatic pattern, tolerance history, and clinician judgment. Treatment continued until disease progression, unacceptable toxicity, patient withdrawal, clinical deterioration, or completion of the simulated observation period.

Dose interruptions, dose reductions, safety monitoring, and supportive care were permitted according to simulated clinical-management rules. Treatment modifications were recorded descriptively and evaluated in relation to persistence and adverse-event patterns.

Endpoints

The primary descriptive endpoint was progression-free survival by biomarker subgroup. Progression-free survival was defined as the time from initiation of Brenova-based therapy to documented disease progression or death from any cause within the simulated analysis framework.

Secondary endpoints included objective response rate, clinical benefit rate, disease control rate, treatment persistence at 24 weeks, safety outcomes, and exploratory associations between biomarker categories and clinical outcomes. Objective response rate was assessed among patients with measurable disease. Clinical benefit rate included complete response, partial response, or stable disease lasting at least 24 weeks.

Statistical Analysis

All analyses were descriptive and based on simulated data. Kaplan–Meier-style summaries were used to estimate median progression-free survival. Hazard ratios were generated for exploratory subgroup comparison and are presented for manuscript realism only. Categorical outcomes were summarized using counts and percentages. Confidence intervals were included for selected

outcomes to support a journal-style presentation. No confirmatory hypothesis testing was performed.

Results

Patient Disposition

A total of 324 patients were included in the simulated translational analysis population. All patients received at least one dose of Brenova-based therapy and had baseline biomarker-status classification. At the time of analysis, 116 patients remained on treatment, 151 had discontinued due to disease progression, 23 had discontinued due to adverse events, 16 had discontinued due to patient preference or logistical reasons, and 18 had discontinued due to physician decision or clinical deterioration not attributed to a specific adverse event.

The median follow-up duration was 13.8 months. Treatment persistence at week 24 was 64.2% in the overall cohort. Persistence was higher among patients with low resistance-marker burden than among those with high resistance-marker burden.

Baseline Characteristics

The median age was 59 years. Most patients had ECOG performance status of 0 or 1. Visceral metastases were present in 57.4% of patients, and liver involvement was present in 34.6%. Prior CDK4/6 inhibitor exposure was reported in 72.2% of patients. Endocrine-sensitive disease was documented in 51.5%, while endocrine-resistant disease was documented in 48.5%.

Table 1. Baseline Clinical and Biomarker Characteristics

Characteristic	Brenova-Based Therapy Cohort (n=324)
Median age, years (range)	59 (32–82)
Age ≥65 years, n (%)	106 (32.7)
ECOG performance status 0, n (%)	148 (45.7)
ECOG performance status 1, n (%)	139 (42.9)
ECOG performance status 2, n (%)	37 (11.4)
Visceral metastases, n (%)	186 (57.4)
Liver involvement, n (%)	112 (34.6)
Bone metastases, n (%)	228 (70.4)
Prior CDK4/6 inhibitor exposure, n (%)	234 (72.2)
Prior chemotherapy for metastatic disease, n (%)	82 (25.3)
Endocrine-sensitive disease, n (%)	167 (51.5)
Endocrine-resistant disease, n (%)	157 (48.5)
Low resistance-marker burden, n (%)	103 (31.8)
Intermediate resistance-marker burden, n (%)	137 (42.3)
High resistance-marker burden, n (%)	84 (25.9)
Low circulating tumor DNA-style signal, n (%)	96 (29.6)

Characteristic	Brenova-Based Therapy Cohort (n=324)
Moderate circulating tumor DNA-style signal, n (%)	151 (46.6)
High circulating tumor DNA-style signal, n (%)	77 (23.8)

Progression-Free Survival by Clinical Subgroup

Median progression-free survival in the overall cohort was 10.9 months. Patients with endocrine-sensitive disease had a median progression-free survival of 13.2 months compared with 8.1 months among patients with endocrine-resistant disease. The exploratory hazard ratio for progression or death in endocrine-sensitive versus endocrine-resistant disease was 0.63 (95% confidence interval, 0.48–0.84).

Patients receiving Brenova-based therapy in earlier metastatic treatment lines had longer progression-free survival than those receiving therapy in later lines. Median progression-free survival was 12.8 months in the second-line metastatic setting, 10.5 months in the third-line setting, and 7.9 months in the fourth or later setting. Patients without visceral metastases had a median progression-free survival of 12.4 months compared with 9.2 months among patients with visceral disease.

Progression-Free Survival by Biomarker Subgroup

Progression-free survival varied by simulated biomarker category. Patients with low resistance-marker burden had a median progression-free survival of 12.7 months, compared with 10.4 months in the intermediate-burden group and 8.6 months in the high-burden group. Patients with low circulating tumor DNA-style signal had a median progression-free survival of 13.1 months, compared with 10.6 months in the moderate-signal group and 7.8 months in the high-signal group.

The combination of endocrine sensitivity and low resistance-marker burden identified the subgroup with the longest median progression-free survival, at 14.6 months. The subgroup with endocrine-resistant disease and high resistance-marker burden had the shortest median progression-free survival, at 6.9 months.

Table 2. Progression-Free Survival and Response by Biomarker Category

Subgroup	n	Median PFS, months	Objective Response Rate, %	Clinical Benefit Rate, %	Week 24 Treatment Persistence, %
Overall cohort	324	10.9	36.8	61.7	64.2
Endocrine-sensitive disease	167	13.2	42.1	68.3	70.1
Endocrine-resistant disease	157	8.1	31.2	54.8	58.0
Low	103	12.7	43.7	70.9	72.8

Subgroup	n	Median PFS, months	Objective Response Rate, %	Clinical Benefit Rate, %	Week 24 Treatment Persistence, %
resistance- marker burden					
Intermediate resistance- marker burden	137	10.4	36.5	61.3	64.2
High resistance- marker burden	84	8.6	28.6	50.0	53.6
Low circulating tumor DNA- style signal	96	13.1	44.8	71.9	73.9
Moderate circulating tumor DNA- style signal	151	10.6	35.8	60.9	63.6
High circulating tumor DNA- style signal	77	7.8	27.3	48.1	51.9

Objective Response and Clinical Benefit

Among patients with measurable disease, the overall objective response rate was 36.8%. Objective response was higher among patients with endocrine-sensitive disease, low resistance-marker burden, and low circulating tumor DNA-style signal. Complete responses were uncommon, and most objective responses were partial responses.

The clinical benefit rate was 61.7% in the overall cohort. Patients with low resistance-marker burden had a clinical benefit rate of 70.9%, while patients with high resistance-marker burden had a clinical benefit rate of 50.0%. These findings suggest that baseline biological context may influence not only tumor shrinkage but also the probability of sustained disease stabilization.

Treatment Persistence

Treatment persistence at week 24 was 64.2% in the overall cohort. Persistence was 72.8% among patients with low resistance-marker burden, 64.2% among those with intermediate burden, and 53.6% among those with high burden. Patients with lower circulating tumor DNA-style signal also demonstrated higher treatment persistence.

The most common reason for discontinuation was disease progression. Discontinuation due to adverse events occurred in 7.1% of patients. Patients with high resistance-marker burden were more likely to discontinue because of progression, while discontinuation due to adverse events was broadly similar across biomarker categories.

Safety and Tolerability

The safety profile of Brenova-based therapy in this simulated translational cohort was consistent with a targeted systemic regimen requiring routine monitoring. The most frequent adverse events were fatigue, nausea, neutropenia, diarrhea, decreased appetite, anemia, and increased alanine aminotransferase. Most events were grade 1 or 2. Grade 3 or higher treatment-emergent adverse events occurred in 32.4% of patients.

Dose interruptions occurred in 34.0% of patients, and dose reductions occurred in 20.7%. Dose modifications were most commonly associated with neutropenia, fatigue, gastrointestinal symptoms, and hepatic laboratory abnormalities. Treatment discontinuation due to adverse events occurred in 23 patients.

Table 3. Treatment-Emergent Adverse Events and Dose Modification Outcomes

Safety Outcome	Brenova-Based Therapy Cohort (n=324)
Any treatment-emergent adverse event, n (%)	272 (84.0)
Grade 3 or higher adverse event, n (%)	105 (32.4)
Serious adverse event, n (%)	41 (12.7)
Dose interruption, n (%)	110 (34.0)
Dose reduction, n (%)	67 (20.7)
Discontinuation due to adverse event, n (%)	23 (7.1)
Fatigue, any grade, n (%)	132 (40.7)
Nausea, any grade, n (%)	103 (31.8)
Neutropenia, any grade, n (%)	98 (30.2)
Diarrhea, any grade, n (%)	89 (27.5)
Decreased appetite, any grade, n (%)	70 (21.6)
Anemia, any grade, n (%)	63 (19.4)
Increased ALT, any grade, n (%)	54 (16.7)
Stomatitis, any grade, n (%)	39 (12.0)

Discussion

This simulated translational oncology manuscript evaluated Brenova-based therapy under the Brexiva brand in HR+/HER2– metastatic breast cancer with emphasis on biomarker patterns and endocrine-resistance features. The model showed that clinical outcomes varied according to endocrine sensitivity, resistance-marker burden, and circulating tumor DNA-style signal. Patients with endocrine-sensitive disease and lower baseline resistance-marker burden experienced longer progression-free survival, higher response rates, higher clinical benefit rates, and greater treatment persistence compared with patients who had endocrine-resistant disease or higher resistance-marker burden.

These findings reflect a clinically plausible pattern in HR+/HER2– metastatic breast cancer. The disease is not biologically uniform, and resistance mechanisms can accumulate over time. Patients with longer prior endocrine benefit and lower evidence of resistance-associated tumor evolution may be more likely to remain sensitive to endocrine-based combinations. Conversely, patients with rapidly progressive disease, high molecular burden, or multiple resistance-associated features may have shorter disease control even when treated with a targeted regimen.

For oncologists and breast cancer specialists, the practical value of biomarker-informed interpretation lies in treatment individualization. Biomarker categories should not replace clinical judgment, but they can provide additional context when evaluating expected benefit, monitoring intensity, and sequencing decisions. In this simulated model, resistance-marker burden and circulating tumor DNA-style signal were not used as strict eligibility criteria; instead, they helped describe outcome heterogeneity. This approach is useful for realistic medical-content design because it mirrors how exploratory translational data are often presented in oncology publications.

The progression-free survival results suggest that Brenova-based therapy may be positioned in a simulated evidence framework as a treatment concept with greater apparent activity in patients who retain endocrine sensitivity or have lower resistance-marker burden. However, the findings also show that patients with higher resistance-marker burden may still experience disease control in a subset of cases. This distinction is important. A balanced scientific interpretation should avoid suggesting that a biomarker group is entirely responsive or nonresponsive. Instead, biomarker findings should be described as associated with differences in probability, duration, and magnitude of benefit.

The response and clinical benefit results further support this interpretation. Objective response rate was higher among patients with favorable biomarker profiles, but clinical benefit rate remained meaningful across multiple subgroups. In HR+/HER2– metastatic breast cancer, sustained stable disease can be clinically relevant, especially when accompanied by manageable tolerability and preserved quality of life. Therefore, translational interpretation should consider both tumor shrinkage and disease stabilization.

Treatment persistence was also associated with biomarker-defined disease behavior. Patients with high resistance-marker burden were more likely to discontinue due to progression, while discontinuation due to adverse events was relatively consistent across biomarker categories. This suggests that biological disease control, rather than tolerability alone, may have been the major driver of persistence differences in the simulated dataset. For clinical communication, this

reinforces the importance of linking persistence outcomes to disease biology, patient selection, and treatment goals.

The safety profile in this simulated cohort was consistent with prior Brenova-style research models. Fatigue, nausea, neutropenia, diarrhea, decreased appetite, anemia, and hepatic laboratory abnormalities were the most frequently reported events. Most events were manageable with monitoring, supportive care, interruption, or dose reduction. Grade 3 or higher events occurred in approximately one-third of patients, indicating that active toxicity management would be necessary in clinical practice. The balanced interpretation is that tolerability appears manageable in the simulated dataset but not negligible.

Clinical Relevance

This manuscript demonstrates how Brexiva/Brenova-related research content can be expanded beyond general efficacy and safety into a translational oncology narrative. For oncologists and breast cancer specialists, biomarker-informed analysis may help frame which patients are more likely to experience durable disease control and which patients may require closer monitoring or earlier consideration of alternative treatment strategies.

The simulated findings may also support realistic content generation for congress posters, field medical summaries, internal evidence decks, and scientific narrative development. A biomarker-focused manuscript allows downstream materials to discuss endocrine resistance, patient selection, disease heterogeneity, and treatment sequencing in a medically credible way.

Limitations

This manuscript is based entirely on simulated clinical data. No actual patients were enrolled, no biomarker testing was performed, and no clinical outcomes were observed in real practice. The resistance-marker categories, circulating tumor DNA-style signal groups, statistical outputs, treatment effects, adverse-event patterns, and subgroup findings were created for medical-content design and workflow demonstration.

The biomarker framework is exploratory and should not be interpreted as a validated diagnostic or predictive system. The categories are intentionally simplified for manuscript clarity and do not capture the full complexity of HR+/HER2- metastatic breast cancer biology. Additionally, the absence of a randomized comparator group prevents conclusions regarding comparative efficacy. Observed subgroup differences may reflect baseline prognosis rather than treatment-specific effect.

The simulated dataset may also be affected by selection bias, missing data, and artificial assumptions regarding follow-up consistency. Real-world biomarker testing availability, sample quality, turnaround time, and interpretation may vary significantly across global markets. These limitations should be considered when using the manuscript for medical-content workflow demonstration.

Future Research Directions

Future research-design scenarios for Brexiva/Brenova could include biomarker-stratified randomized models, longitudinal circulating tumor DNA-style monitoring, evaluation of

acquired resistance at progression, integration of patient-reported outcomes with biomarker status, and sequencing analyses after prior targeted therapy. Additional simulated studies could explore whether early molecular change predicts treatment persistence, response depth, or time to treatment discontinuation.

Conclusion

In this simulated translational oncology research manuscript, Brenova-based therapy under the Brexiva brand demonstrated variable clinical outcomes across endocrine-resistance and biomarker-defined subgroups in HR+/HER2– metastatic breast cancer. Patients with endocrine-sensitive disease, lower resistance-marker burden, and lower circulating tumor DNA-style signal showed longer progression-free survival, higher response measures, and greater treatment persistence. Safety findings were consistent with a targeted systemic therapy model requiring active monitoring and dose modification in a subset of patients. These findings provide a realistic research-paper structure for integrating biomarker context, endocrine-resistance features, efficacy outcomes, and safety data into oncology medical-content workflows for Brexiva/Brenova.

Medical Writing Note

This manuscript uses simulated clinical data for medical-content design and workflow demonstration purposes. Brenova is presented as an investigational therapeutic concept associated with the Brexiva brand within a synthetic research model. The biomarker categories, clinical outcomes, statistical outputs, safety findings, and subgroup interpretations are illustrative and should not be interpreted as real clinical evidence, treatment guidance, regulatory evidence, or prescribing information.

References

- [1] Current guideline on management of HR+/HER2– metastatic breast cancer.
- [2] Review of endocrine resistance mechanisms in metastatic breast cancer.
- [3] Review of biomarker-informed treatment strategies in advanced breast cancer.
- [4] Review of circulating tumor DNA applications in metastatic breast cancer.
- [5] Review of treatment sequencing after prior endocrine-based targeted therapy.
- [6] Methodology reference for translational oncology cohort analysis.
- [7] Review of progression-free survival interpretation in oncology studies.
- [8] Review of clinical benefit and disease stabilization in metastatic breast cancer.
- [9] Review of adverse event management for targeted systemic therapies.
- [10] Medical writing guidance for balanced presentation of simulated clinical data.