

Brexiva Safety Guideline Document

Important Safety Information for Brenova in HR+/HER2– Metastatic Breast Cancer

1. Indication

Brexiva is a fictional internal oncology brand for Brenova, intended for use in simulated medical-content workflows for patients with HR+/HER2– metastatic breast cancer. This document is designed for HCP-facing safety communication in a global campaign context for oncologists and breast cancer specialists.

2. Contraindications

Treatment with Brenova should be contraindicated in patients with a known history of serious hypersensitivity to Brenova or any component of its formulation. Brenova should not be administered to patients who previously experienced a life-threatening administration-related reaction associated with the product in the simulated treatment setting.

3. Warnings and Precautions

Brenova may be associated with clinically important safety risks requiring careful patient selection, baseline assessment, and ongoing monitoring. Key areas of concern include:

- Serious hypersensitivity or administration-related reactions
- Myelosuppression, including neutropenia, anemia, or thrombocytopenia
- Hepatic function abnormalities, including elevated liver enzymes
- Increased infection risk, particularly in immunocompromised patients
- Potential embryo-fetal risk when administered during pregnancy
- Additive toxicity when used with other myelosuppressive or hepatotoxic therapies
- Need for treatment interruption, dose delay, or discontinuation based on clinical severity

Healthcare professionals should evaluate patient-specific risk factors before initiating Brenova and monitor for new or worsening adverse events throughout treatment.

4. Administration-Related Reactions

Administration-related reactions may occur during or after Brenova administration. Possible symptoms may include rash, flushing, pruritus, chills, fever, nausea, dizziness, dyspnea, hypotension, chest discomfort, or swelling.

Patients should be observed during administration and for an appropriate period afterward based on clinical judgment. For mild or moderate reactions, administration may be interrupted and supportive care provided. For severe or life-threatening reactions, Brenova should be stopped immediately, urgent medical management should be initiated, and re-treatment should be carefully reassessed.

5. Hematologic Toxicity

Brenova may be associated with hematologic toxicity, including neutropenia, anemia, thrombocytopenia, or other blood-count abnormalities. Complete blood count assessment should be considered before treatment initiation and periodically during therapy.

Patients with significant baseline cytopenias or prior intensive anticancer therapy may require closer monitoring. Treatment delay, interruption, or discontinuation may be considered for clinically meaningful hematologic toxicity or complications such as febrile neutropenia or bleeding risk.

6. Hepatic Monitoring

Hepatic function abnormalities may occur during treatment with Brenova. Liver function tests should be assessed at baseline and monitored periodically during treatment.

Use caution in patients with pre-existing hepatic impairment, elevated transaminases, or concurrent use of hepatotoxic therapies. If clinically significant liver enzyme elevation or hepatic dysfunction occurs, treatment interruption and medical evaluation should be considered.

7. Infections

Patients receiving oncology therapies may be at increased risk for infection, especially those with immune compromise, neutropenia, prior chemotherapy exposure, or advanced disease burden. Patients should be monitored for signs and symptoms of infection during treatment.

Brenova administration should be delayed in patients with active serious infection until the infection is clinically controlled. Healthcare professionals should evaluate fever, respiratory symptoms, urinary symptoms, unexplained fatigue, or localized signs of infection promptly.

8. Embryo-Fetal Risk

Based on the intended oncology mechanism and potential effects on rapidly dividing cells, Brenova may cause fetal harm if administered during pregnancy. Pregnancy status should be assessed before treatment initiation when clinically applicable.

Patients of reproductive potential should be counseled on the potential risk to a fetus and the need for effective contraception during treatment according to internal medical guidance and local review standards. Breastfeeding considerations should be discussed due to the potential for serious adverse reactions in an exposed infant.

9. Adverse Reactions

Possible adverse reactions associated with Brenova may include:

- Fatigue
- Nausea
- Decreased appetite

- Diarrhea
- Rash or pruritus
- Administration-site or infusion-related reactions
- Laboratory abnormalities
- Neutropenia
- Anemia
- Thrombocytopenia
- Elevated liver enzymes
- Infection-related complications

Adverse reactions should be evaluated according to severity, duration, reversibility, and relationship to treatment.

10. Patient Counseling Information

Healthcare professionals should advise patients to report symptoms of hypersensitivity, fever, chills, persistent cough, shortness of breath, unusual bruising or bleeding, severe fatigue, yellowing of the skin or eyes, dark urine, severe diarrhea, rash, or signs of infection.

Patients should be counseled not to ignore new or worsening symptoms during treatment and to contact their healthcare professional promptly if concerning events occur.

11. Safety Monitoring Checklist

Before and during Brenova treatment, HCPs should consider:

- Review of hypersensitivity history
- Baseline clinical assessment and medication review
- Complete blood count monitoring
- Liver function test monitoring
- Pregnancy status assessment when applicable
- Infection screening and symptom review
- Evaluation of concomitant myelosuppressive or hepatotoxic therapies
- Documentation of adverse events and management actions
- Treatment interruption or discontinuation assessment when clinically required

12. Closing Disclaimer

This Safety Guideline Document is for internal fictional brand simulation and medical-content workflow demonstration only. It is not real prescribing information, not medical advice, and should not be used to guide actual patient treatment decisions.