



TIC 232261044

Eclipsing binary candidate dossier - Star Vetter

THIN MARGIN

BLS EPHEMERIS		CATALOG CROSS-MATCH (novelty)	
Period	0.810821 d (ALIASED, see below)		
-> corrected period: 1.621641 d		VSX	no match
Epoch (t0)	1468.3491 BTJD	ASAS-SN	no match
Duration	0.1200 d	Gaia eclipsing-binary table	no match
Depth	2.534 % (25,343 ppm)	Gaia variable classification	no match
SNR	88.5	TESS EB catalog (Prsa+2022)	no match
		ExoFOP TOI list	no match
INDEPENDENT CHECKS (f3i gates)		Verdict: NOVEL	
Eclipses observed	23 (gate >= 3)		
Robust depth ratio	1.07 (gate >= 0.50)		
Robust depth sigma	53.0 (gate >= 7.0)		
Odd/even z-score	5.53 (ALIAS FLAGGED)		
Secondary eclipse	not detected		
PIXEL-LEVEL SOURCE CHECK (TESScut difference imaging)			
Sector 6	light-curve dip in this sector: 53.8 sigma	t0 refined by 0.0025 phase	
Difference-image peak SNR	13.06 (gate >= 5.0)	Compactness	0.94 (gate >= 0.50)
Dimming centroid offset	0.12 px (3 arcsec)	(gate <= 1.0 px, i.e. on the target)	
Nearest bright-enough neighbour in cutout: TIC 713665868			
0.20 px away, dTmag +0.56 - farther from the dimming than the target			



Caveat (PERIOD ALIAS): Odd/even eclipse-depth z-score is 5.53 (gate >= 7.0), aliasing eclipses differ significantly in depth, the signature of a true period double what BLS found. Pixel check confirmation and eclipse timing are unaffected - report the corrected period, 1.621641 d, in any write-up. Caveat (THIN MARGIN): The ON_TARGET call rests on a neighbour-competition margin of just 0.075 px (dimming centroid 0.125 px from target vs 0.200 px from the nearest bright-enough neighbour) - inside diag_pixel.py's own stated centroid systematic of 0.10-0.3 px. TESS's resolution cannot confidently separate this star from that neighbour.