

1 Variants at *RNF212* and *RNF212B* are associated with
2 recombination rate variation in Soay sheep (*Ovis aries*).

3 Susan E. Johnston^{1,*}, Martin Stoffel¹, Josephine M. Pemberton¹

4 ¹Institute of Evolutionary Biology, School of Biological Sciences,
5 University of Edinburgh, Edinburgh, EH9 3FL, United Kingdom

6 ^{*}Corresponding author: Susan.Johnston@ed.ac.uk

7 Abstract

8 [to be written]

9 Introduction

Meiotic recombination is ubiquitous in sexually-reproducing organisms and is an important driver of genomic diversity in eukaryotes (BARTON and CHARLESWORTH, 1998; OTTO and LENORMAND, 2002). The process of crossing-over ensures the proper disjunction of homologous chromosomes during meiosis (HASSOLD and HUNT, 2001), with most species having a minimum requirement of at least one obligate crossover per chromosome pair (OTTO and PAYSEUR, 2019). It also has the advantage of generating novel haplotypes and preventing the accumulation of harmful mutations in the genome, allowing populations to respond to selection faster and to a greater degree (MULLER, 1964; HILL and ROBERTSON, 1966; BATTAGIN *et al.*, 2016). However, this can come at the cost of increased mutations at crossover sites (HALLDORSSON *et al.*, 2019) and the break up of favourable haplotypes (BARTON and CHARLESWORTH, 1998). Variation in recombination rate is likely to arise across taxa with variation in genome size and chromosome number (STAPLEY *et al.*, 2017); in addition, the evolutionary costs and benefits of recombination may vary depending on the selective context, again leading to the expectation of recombination rate variation within and between populations (OTTO and BARTON, 2001; OTTO and LENORMAND, 2002). Indeed, cytogenetic and linkage mapping studies have shown that recombination rates can vary by orders of magnitude within and between chromosomes (MYERS *et al.*, 2005), individuals (KONG *et al.*, 2004), populations (SAMUK *et al.*, 2020), sexes (LENORMAND and DUTHEIL, 2005) and species (STAPLEY *et al.*, 2017).

In mammals, genomic studies in different species (i.e. humans, cattle, sheep, pigs and deer) have shown that recombination rate is often heritable, with a significant proportion of the additive genetic variation underpinned by loci of moderate to large effects (SANDOR *et al.*, 2012; KONG *et al.*, 2014; MA *et al.*, 2015; KADRI *et al.*, 2016; PETIT *et al.*, 2017; JOHNSTON *et al.*, 2018; JOHNSON *et al.*, 2020). Genome-wide association studies (GWAS) have shown repeated associations with variants at ring finger protein 212 *RNF212* and its paralogue *RNF212B*, whose associated proteins are essential for the crossover designation process during meiosis (REYNOLDS *et al.*, 2013), as well as other loci with established roles in meiotic processes (e.g.

37 *HEI10*, *REC8* and *TOP2*; SANDOR *et al.* 2012; KONG *et al.* 2014; MA *et al.* 2015; KADRI *et al.*
38 2016; PETIT *et al.* 2017; JOHNSTON *et al.* 2018; JOHNSON *et al.* 2020). This heritable, oligogenic
39 architecture implies that recombination rate has the potential to evolve rapidly to selection
40 if variation in recombination rate is associated with individual fitness (OTTO and BARTON,
41 2001; STAPLEY *et al.*, 2017). Therefore, detailed examination of the genetic architecture of
42 recombination rate is a critical step in understanding how rates are evolving over different
43 evolutionary timescales.

44 The Soay sheep (*Ovis aries*) are a Neolithic breed of domestic sheep that have been living
45 wild on the St Kilda archipelago since the Bronze age. These sheep are a model system for
46 studies of ecology and evolution, and have been intensively studied since 1985 (CLUTTON-
47 BROCK *et al.*, 2004). A previous study integrated pedigree information with genotypic data
48 from ~39,000 SNPs to characterise autosomal crossover counts (ACCs) in ~3,300 gametes
49 transmitted from parents to their offspring (JOHNSTON *et al.*, 2016). A GWAS showed that a
50 single region on chromosome 6, corresponding to the locus *RNF212*, explained around 47%
51 of heritable variation in female ACC only. A broader-scale “regional heritability” analysis
52 identified a tentative association in both sexes at a 1.09 Mb region on chromosome 7, con-
53 taining the candidate loci *REC8* and *RNF212B*), but this was not significant in the GWAS
54 or in sex-specific regional heritability models. Since then, addition of more SNP genotyped
55 individuals and improvements in pedigree construction methodology (HUISMAN, 2017) now
56 allows us to characterise recombination rate in a further ~3,900 gametes. Furthermore, ad-
57 vances in genotype imputation methods in complex pedigrees has allowed us to increase the
58 SNP dataset for conducting genome-wide association study from ~39,000 to ~417,000 SNPs.

59 Here, we revisit our previous analysis to carry out GWAS of male and female recombina-
60 tion rate. Our motivation for this study was to: (a) confirm a sex-limited association with
61 *RNF212* on chromosome 6; (b) fine-map the association on chromosome 7 and determine the
62 action and effect size of this locus in both males and females; and (c) to identify any further
63 associations that could not be detected using the previous dataset.

64 Materials and Methods

65 Study Population.

66 Soay sheep living within the Village Bay area on the island of Hirta, St Kilda, Scotland
67 (57°49'N, 8°34'W) have been individually studied since 1985 (CLUTTON-BROCK *et al.*, 2004).
68 All sheep are ear-tagged at their first capture, including 95% of lambs born within the study
69 area, and the majority of animals are recaptured on an annual basis. DNA samples are rou-
70 tinely obtained from ear punches and/or blood sampling. All animal work is carried out
71 according to UK Home Office procedures and was licensed under the UK Animals (Scientific
72 Procedures) Act 1986 (License no. PPL60/4211).

73 Genotype and Pedigree Data.

74 All sheep were genotyped at 51,135 SNP markers on the Illumina Ovine SNP50 BeadChip
75 (KIJAS *et al.*, 2009). Quality control was carried out using the check.marker function in
76 GenABEL v1.8-0 with the following thresholds: SNP minor allele frequency (MAF) > 0.001,
77 SNP locus genotyping success > 0.99, individual sheep genotyping success > 0.95; a total
78 of 7,700 sheep and 39,398 SNPs remained. Pedigree relationships were previously inferred
79 using data from 438 SNP loci in the R package Sequoia v1.02 (HUISMAN, 2017) and from field
80 observations between mothers and their offspring born within the study area. Autosomal
81 SNP genotypes were used to calculate the \hat{F}_{III} genomic inbreeding coefficients using GCTA
82 v1.90.2 (YANG *et al.*, 2011).

83 A further 189 individuals were genotyped at 430,702 polymorphic SNPs on the Ovine In-
84 finium HD SNP BeadChip for imputation of genotypes into individuals typed on the 50K
85 chip (see JOHNSTON *et al.* 2016 for method and individual selection criteria and STOFFEL *et al.*
86 2020 for full imputation method for this dataset). Briefly, SNP genotypes from the HD Chip
87 were imputed into the SNP50 Chip typed individuals using AlphaImpute v1.98 (HICKEY *et al.*,
88 2012; ANTOLÍN *et al.*, 2017) resulting in a dataset with 7,691 individuals genotyped at 417,373

89 SNPs, with a mean genotyping rate per individual of 99.5% (range 94.8% - 100%). As there
90 were no X-chromosome SNPs in common between the two SNP Chips, only the SNP50
91 Chip SNPs were used for association analyses on the X chromosome. SNP positions for both
92 chips were are known relative to sheep genome assembly Oar_v3.1 (GenBank assembly ID
93 GCA_000298735.1).

94 **Estimation of autosomal crossover counts and linkage maps.**

95 Autosomal crossover positions were estimated using an identical protocol to that outlined in
96 JOHNSTON *et al.* (2016). Briefly, the method uses Ovine SNP50 BeadChip SNP data from a
97 focal individual's parents, mate and offspring to characterise crossovers that occurred in the
98 gametes transmitted from the focal individual to its offspring. Crossovers were determined
99 using the software CRI-MAP v2.504a (GREEN *et al.*, 1990). Here, we estimated ACC for a
100 further 3,908 gametes, leading to a dataset of ACC for 7,235 gametes transmitted from 1,632
101 unique focal individuals. Because of differences in the SNP quality control between analyses,
102 we used 37,853 SNPs in common with the previous recombination rate study. Simulation
103 studies showed that this method will identify >99% of crossovers, meaning that ACC es-
104 timated using this method can be used as a proxy for individual recombination rate. The
105 method also provides linkage maps for each chromosome, allowing us to update the existing
106 map for Soay sheep to include information from a much larger number of meioses.

107 **Animal Models**

108 A restricted maximum likelihood (REML) animal model approach (HENDERSON, 1975) was
109 used to estimate components of phenotypic variance for ACC, including the additive genetic
110 effect (i.e. heritability). A genomic relatedness matrix (GRM) was constructed for all poly-
111 morphic autosomal SNPs from the 50K chip using GCTA v1.90.2 (YANG *et al.*, 2011) and
112 was adjusted with the argument `--grm-adj 0`, which assumes a similar frequency spectra of
113 genotyped and causal loci. Related individuals were not removed from the GRM as there is
114 high relatedness in the study population, and there are were no significant parental effects

115 or common environment effects on individual recombination rate (JOHNSTON *et al.*, 2016).
 116 Animal models were constructed for both sexes combined, females only and males only, and
 117 were run in ASReml-R v4 (BUTLER *et al.*, 2009) in R v3.6.2. Fixed effects included sex (in
 118 the combined model) and the genomic inbreeding coefficient \hat{F}_{III} ; initial models fit age in
 119 years fit as a linear covariate, but as it was not significant, it was removed from the final mod-
 120 els (Figure 1). The significance of fixed effects was tested using Wald tests. Random effects
 121 included the additive genetic effect (as estimated using the GRM) and identity of the focal
 122 individual (i.e. “permanent environment”) effect to account for repeated measures. Despite
 123 not being significant, the fixed effect \hat{F}_{III} and the random permanent environment effect were
 124 retained in all models to account for potential underestimation of ACC due to long runs of
 125 homozygosity and pseudoreplication, respectively. Significance of the additive genetic effect
 126 was determined by dropping the additive genetic effect from each model and comparing with
 127 the full model using a likelihood ratio test distributed as χ_1^2 . Bivariate models examining
 128 the genetic correlation r_A between male and female ACC were run using the CORGH error
 129 structure in ASReml-R (correlation with heterogeneous variances). Models were set with r_A
 130 to be unconstrained. To test whether r_A was significantly different from 0 and 1, the model
 131 was compared to models with r_A fixed at a value of 0 or 0.999 using likelihood ratio tests.

132 **Genome-wide association studies.**

133 Genome-wide association studies of ACC with the imputed SNP dataset were carried out us-
 134 ing function *rGLS* in the package RepeatABEL v1.1.31 (RÖNNEGÅRD *et al.*, 2016) in R v3.6.2.
 135 This approach fits both repeated measures and the GRM, the latter accounting for population
 136 structure. Association statistics were corrected for any further inflation by dividing by the
 137 genomic control parameter λ , calculated as the median observed χ_1^2 statistic divided by the
 138 median χ_1^2 expected under a null distribution. The significance threshold at $\alpha = 0.05$ was
 139 calculated by STOFFEL *et al.* (2020) as $P < 1.28 \times 10^{-06}$, using a ‘simpleM’ approach using
 140 linkage disequilibrium information to determine the effective number of independent tests
 141 (GAO *et al.*, 2008).

142 For the most highly-associated SNPs, genotype effect sizes were estimated by fitting the SNP
143 genotype as a three level factor in the animal models described above. Then, in separate
144 models, the proportion of phenotypic variance explained by significant regions was then
145 determined using a regional heritability approach. For these models, GRMs were constructed
146 as above for the 20 SNPs from the SNP50 chip spanning the highest association SNP (i.e.
147 10 SNPs from either side of the association). In one case where the association was at the
148 end of the chromosome, the last 20 SNPs on that chromosome were used to construct the
149 GRM. These regional GRMs were then fit as additional random effects in the animal models
150 described above, with their significance determined using likelihood ratio tests distributed as
151 χ^2_1 .

152 In order to identify potential candidate genes for ACC, sheep gene IDs SNPs in the broad
153 associated regions were extracted from Ensembl (gene build ID Oar_v3.1.100) using the func-
154 tion *getBM* in the R package biomaRt v2.42.1 (DURINCK *et al.*, 2009) in R v3.6.2. Gene
155 orthologues in humans (*Homo sapiens*), cattle (*Bos taurus*), mouse (*Mus musculus*) and rat
156 (*Rattus norvegicus*) were extracted using the function *getLDS*. The associated gene ontology
157 (GO annotations) information for all genes and orthologues were then extracted, again using
158 the *getBM* function. All gene names, phenotype descriptions, GO terms and definitions were
159 then queried for terms associated with meiosis and recombination, using the R command *grep*
160 with the text strings *meio* and *recombin*.

161 Data Availability

162 Raw data will be publicly archived. Code for the analysis is archived at
163 https://github.com/susjoh/Soay_Recombination_2020.

164 Results

165 Variation and heritability of autosomal crossover count.

166 Males had higher recombination rates than females, with 7.24 more autosomal crossovers
167 observed per gamete ($SE = 0.18$. Wald $\chi^2_1 = 1613.6$, $P < 0.001$; Figure 1). There was no
168 association with ACC and age or the inbreeding coefficient \hat{F}_{III} ($P > 0.05$; Figure 1). Females
169 had higher phenotypic variance than males (female $V_P = 32.41$, male $V_P = 24.162$; Table 1).
170 ACC was heritable in both sexes ($h^2 = 0.148$) and in males and females separately ($h^2 =$
171 0.118 and 0.181 , respectively; Table 1); likelihood ratio tests confirmed that including the
172 additive genetic effect significantly improved the animal models ($\chi^2_1 = 139.4$, 19.72 and 119.9
173 for both sexes, males, and females, respectively; $P < 0.001$). The permanent environment
174 contribution to phenotypic variance was not significant in any models ($P > 0.05$), with the
175 remaining variance attributed to residual effects (Table 1). Bivariate models of ACC between
176 males and females showed a positive genetic correlation ($r_A = 0.555$, $SE = 0.138$) that was
177 significantly different from both $r_A = 0$ and 1 ($\chi^2_1 = 11.95$ and 17.91 , respectively; $P <$
178 0.001). Linkage map lengths from the ACC analysis in this study were strongly correlated
179 with the previous maps (linear regression, adjusted $R^2 > 0.999$, Figure S1). The updated map
180 had 15,787 unique centiMorgan positions compared with 12,359 in the previous study; it is
181 provided in Table S2 and Figure S2.

182 Genome-wide Association Study.

183 There was a strong association between SNPs at the sub-telomeric region of chromosome 6
184 and female ACC (*oar3_OAR6_116402578*, $P = 2.45 \times 10^{-17}$; Figures 2 & S4; Tables 2 & S3).
185 This locus explained 45.9% of the heritable variation in female ACC, with a difference of
186 5 crossovers per gamete between the two homozygotes (Figure S3, Table 2). There was no
187 significant association with this locus and male ACC, confirming that its action is likely to be
188 sex-limited. The same SNP showed the highest genome-wide association in the previous study

189 (JOHNSTON *et al.*, 2016) and occurs 25.2 kb from the putative location of *RNF212*. It should
 190 be noted that *RNF212* is not annotated on the Oar_v3.1 sheep genome due to a gap in the
 191 sequence, but is predicted to be positioned between 116,427,802 and 116,446,904 bp (JOHN-
 192 STON *et al.*, 2016); *RNF212* is present in this position on the Oar_rambouillet_v1.0 genome
 193 assembly (GenBank assembly ID GCA_002742125.1). A second association was observed in
 194 both sexes on chromosome 7 (*oar3_OAR7_21368818* and *oar3_OAR7_21347355*, $P = 6.11 \times$
 195 10^{-13} , 1.83×10^{-10} and 7.20×10^{-7} in both sexes combined, females and males, respectively);
 196 the association statistic in males was also identical for locus *oar3_OAR7_21116299* (Figures 2
 197 & S5; Tables 2 & S3). The action of this locus was similar between males and females, with
 198 a difference of ~4.5 crossovers per gamete between the two homozygotes (Figure S3, Table
 199 2) and explaining between 19.7% and 24.8% of the heritable variation in ACC. These SNPs
 200 spanned the candidate locus *RNF212B* (21,241,831 to 21,273,165 bp). GO term analysis indi-
 201 cated that both *RNF212* and *RNF212B* were the closest genes to each hit that were associated
 202 with meiotic processes (Table S4; Figures S4 & S5).

203 Discussion

204 This study revisited a previous analysis investigating the genetic architecture of ACC in Soay
205 sheep, with more than double the number of gametes and more than ten times the number
206 of SNP loci. We confirmed that ACC is ~ 1.27 times higher in males than in females, and is
207 not associated with age or inbreeding in neither males nor females. We identified two candi-
208 date genes associated with ACC: *RNF212* on chromosome 6, which is associated with female
209 recombination rate only; and its paralogue *RNF212B* on chromosome 7, which is associated
210 with recombination rate in both sexes. Both loci have repeatedly been associated with recom-
211 bination rate variation in mammals, with all studies conducted to date implicating either one
212 or both of these loci; these include humans (KONG *et al.*, 2008, 2014), cattle (SANDOR *et al.*,
213 2012; MA *et al.*, 2015; KADRI *et al.*, 2016), pigs (JOHNSON *et al.*, 2020), deer (JOHNSTON *et al.*,
214 2018) and domestic sheep (PETIT *et al.*, 2017). Functional studies in mice indicate that the
215 protein RNF212 is essential for crossover formation during meiosis, and that it has a dosage
216 sensitive effect (REYNOLDS *et al.*, 2013). We observe an additive effect of alleles on ACC at
217 both loci in this study, suggesting that a similar dosage sensitive effect is the mechanism driv-
218 ing rate differences in Soay sheep.

219 The current study confirms the previous association at *RNF212* and does not gain any novel
220 insights into the nature of its association with ACC. However, the observation at *RNF212B*
221 builds on tentative evidence of an association in this genomic region. In the 2016 study, a
222 genome-wide regional heritability analysis identified an association in both sexes combined
223 in a 1.09Mb region, which contained *RNF212B* and another candidate locus, meiotic recom-
224 bination protein *REC8*. However, there was no corresponding association using GWAS and
225 there was no significant effect observed in the sex-specific regional heritability analyses, pro-
226 viding limited insight into the action of this region on ACC variation. In this study, we have
227 shown that adding a larger number of gametes and SNPs has increased the power to detect
228 an association at *RNF212B*, and indicating that ACC variation is more likely to be attributed
229 to *RNF212B* than *REC8* (although see further discussion below). The MAF of *RNF212B* is
230 relatively low (0.082), suggesting that lower sample sizes in the 2016 study were not sufficient

231 to detect the effects of rare loci within the population. Using the SNP50 Chip alone, the
232 increased number of gametes in current study would shown a significant on chromosome 7
233 in females (but not in males; Figure S5), however, the highest association would be around
234 345Kb away from the that observed in the current study. Our work shows that increasing
235 marker density using imputation is likely to be an increasingly important approach in con-
236 ducting GWAS in natural populations.

237 Soay sheep have extensive linkage disequilibrium (LD) throughout the genome due to recent
238 population bottlenecks, a small effective population size and a high prevalence of inbreeding
239 (CLUTTON-BROCK *et al.*, 2004; STOFFEL *et al.*, 2020). An advantage of this is that there is a
240 higher change of typing SNP loci that are in LD with causal loci, and that genotype imputa-
241 tion can be carried out with high accuracy. However, a disadvantage is that high LD within
242 the population makes it difficult to separate the effects of linked loci that also contribute to
243 phenotypic variation in the population. The associated regions on both chromosomes 6 and
244 7 extend over several megabases, with several other loci associated with meiotic processes
245 occurring within these regions (Figures S4 & S5, Table S4); we cannot rule out that these
246 loci also affect variation in ACC in Soay sheep. Functional validation of the role of *RNF212*
247 and *RNF212B* is not possible within this wild system as experimental manipulation and in-
248 vasive sampling are prohibited. However, further studies in domestic sheep (e.g. PETIT *et al.*
249 2017) and related species may further elucidate the role of loci within these regions and the
250 relative importance of regulatory and protein-coding variation in driving recombination rate
251 variation.

252 Overall, this study has highlighted the merits of genotype imputation and increasing sample
253 sizes to determine the genetic architecture of recombination rate variation in a natural system.
254 Identifying candidate loci and their sex-specific effect sizes provides a stepping stone for future
255 studies investigating the evolution of recombination rates within this and related systems,
256 through modelling the temporal dynamics of these loci, their origin and their association
257 with individual fitness.

258 Acknowledgements

259 We thank Jill Pilkington, Ian Stevenson and all Soay sheep project members and volunteers
260 for collection of data and samples. Camillo Bérénos, Philip Ellis and Hannah Lemon prepared
261 DNA samples for Ovine SNP50 BeadChip genotyping and Jisca Huisman assisted with the
262 pedigree construction. This work has made extensive use of the resources provided by the
263 Edinburgh Compute and Data Facility (<http://www.ecdf.ed.ac.uk/>). Permission to work on
264 St Kilda is granted by The National Trust for Scotland, and logistical support is provided
265 by QinetiQ and Eurest. The Soay sheep project has been supported by grants from the UK
266 Natural Environment Research Council, with the SNP genotyping funded by a European
267 Research Council Advanced Grant to JMP. SEJ is supported by a Royal Society University
268 Research Fellowship.

269 Author Contributions

270 S.E.J and J.M.P. conceived the study. J.M.P organised the collection of samples. M.S. con-
271 ducted the genotype imputation. S.E.J. analysed the data and wrote the paper. All authors
272 contributed to revisions.

273 Figures and Tables




Figure 1: Association between autosomal crossover count and: A. sex; B. inbreeding coefficient \hat{F}_{III} ; and C. age in years. Plots were made using the raw data. Points are red for females and blue for males. Lines in B. and C. are general additive model smoothing parameters as fit by the plotting library ggplot2 v3.3.2 (WICKHAM, 2016) in R v3.6.2. Points in plot C are slightly jittered between the sexes on the x-axis to more clearly visualise the sex differences with age. Sample sizes are provided in Table 1.




Figure 2: Genome-wide association plots of autosomal crossover counts for A. both sexes combined, B. females only and C. males only. The dashed line is the significant threshold equivalent to $\alpha < 0.05$. The left panels show the association for individual SNPs relative to their genomic position, with points colour-coded by chromosome. The right plots show the distribution of the observed $-\log_{10}P$ values against those expected under a null distribution. All association statistics have been corrected for genomic control using the λ parameter. All results are provided in Table S3. Sample sizes are provided in Table 1.

Table 1: Data information and animal model results for autosomal crossover count (ACC) in Soay sheep. Numbers in parentheses are the standard error unless otherwise stated. N_{OBS} is the observed number of gametes, with numbers in parentheses the number of unique focal individuals. N_{xovers} are the total number of crossovers in the dataset. The mean ACC and the variance (V_{OBS}) were calculated from the raw data. V_P is phenotypic variance. h^2 , pe^2 and e^2 are the narrow-sense heritability, the permanent environment effect, and the residual effect, respectively; all are calculated as the proportion of V_P .

Analysis	N_{OBS}	N_{xovers}	Mean	V_{OBS}	V_P	h^2	pe^2	e^2
Both Sexes	7626 (1483)	227646	29.851	41.32	29.514 (0.588)	0.148 (0.019)	0.025 (0.013)	0.827 (0.014)
Males	3043 (434)	103914	34.149	23.974	24.162 (0.72)	0.118 (0.032)	0 (0.024)	0.882 (0.02)
Females	4583 (1049)	123732	26.998	32.439	32.41 (0.797)	0.181 (0.024)	0 (0.017)	0.819 (0.017)

Table 2: The most highly associated SNPs from GWAS of ACC in Soay sheep. MAF is the minor allele frequency (corresponding to the A allele for each locus). Sex indicates if the model was run in both sexes combined, or in females or males only. P is the P-value for the association statistic after correction using genomic control λ . Effect sizes are provided for animal models where the genotype fit as a fixed factor, and are given relative to the model intercept of 0 at genotype AA. Prop. V_A is the proportion of the additive genetic variance explained by the genomic region containing the SNP locus. All values in parentheses are standard errors. Full GWAS results are provided in Table S3. Effect sizes and associated sample sizes for each genotype are shown in Figure S3.

SNP Information	MAF	Sex	P	Effect AG	Effect GG	Prop. V_A
oar3_OAR6_116402578	0.275	Both Sexes	2.45×10^{-17}	2.201	3.837	0.399
Chr. 6				(0.332)	(0.361)	(0.118)
Position 116,402,578		Females	1.68×10^{-18}	3.038	4.985	0.459
Candidate Gene: <i>RNF212</i>				(0.397)	(0.423)	(0.115)
		Males	0.0317	0.168	0.884	0.049
				(0.577)	(0.607)	(0.067, ns)
oar3_OAR7_21368818*	0.082	Both Sexes	6.11×10^{-13}	-2.383	-4.520	0.242
Chr. 7				(0.922)	(0.952)	(0.102)
Position 21,368,818		Females	1.83×10^{-10}	-2.705	-4.757	0.197
Candidate Gene: <i>RNF212B</i>				(1.152)	(1.194)	(0.09)
		Males	7.20×10^{-07}	-2.479	-4.394	0.248
				(1.454)	(1.447)	(0.145)

* This association was identical to hits at SNP loci oar3_OAR7_21347355 at position 21,347,355 (in both sexes, males and females) and oar3_OAR7_21116299 at position 21,116,299 (in males).

274 References

- 275 ANTOLÍN, R., C. NETTELBLAD, G. GORJANC, D. MONEY, and J. M. HICKEY, 2017 A hybrid
276 method for the imputation of genomic data in livestock populations. *Genetics Selection*
277 *Evolution* : 1–17.
- 278 BARTON, N. H., and B. CHARLESWORTH, 1998 Why sex and recombination? *Science* **74**:
279 187–195.
- 280 BATTAGIN, M., G. GORJANC, A.-M. FAUX, S. E. JOHNSTON, and J. M. HICKEY, 2016 Effect of
281 manipulating recombination rates on response to selection in livestock breeding programs.
282 *Genet. Select. Evol.* **48**: 44.
- 283 BUTLER, D. G., B. R. CULLIS, A. R. GILMOUR, and B. J. GOGEL, 2009 Mixed Models for S
284 language Environments: ASReml-R reference manual.
- 285 CLUTTON-BROCK, T., J. PEMBERTON, T. N. COULSON, I. R. STEVENSON, and A. D. C. MAC-
286 COLL, 2004 The sheep of St Kilda. In *Soay Sheep: Dynamics and Selection in an Island*
287 *Population*.2. Cambridge University Press, 17–51.
- 288 DURINCK, S., P. T. SPELLMAN, E. BIRNEY, and W. HUBER, 2009 Mapping identifiers for the inte-
289 gration of genomic datasets with the R/Bioconductor package biomaRt. *Nature protocols*
290 **4**: 1184–1191.
- 291 GAO, X., J. STARMER, and E. R. MARTIN, 2008 A multiple testing correction method for
292 genetic association studies using correlated single nucleotide polymorphisms. *Genetic epi-*
293 *demiology* **32**: 361–369.
- 294 GREEN, P., K. FALLS, and S. CROOKS, 1990 *Documentation for CRIMAP, version 2.4*. Washing-
295 ton University School of Medicine.
- 296 HALLDORSSON, B. V., G. PALSSON, O. A. STEFANSSON, H. JONSSON, M. T. HARDARSON, *et al.*,
297 2019 Characterizing mutagenic effects of recombination through a sequence-level genetic
298 map. *Science* **363**: eaau1043.

299 HASSOLD, T., and P. HUNT, 2001 To err (meiotically) is human: the genesis of human aneu-
 300 ploidy. *Nat. Rev. Genet.* **2**: 280–291.

301 HENDERSON, C. R., 1975 Best linear unbiased estimation and prediction under a selection
 302 model. *Biometrics* **31**: 423–447.

303 HICKEY, J. M., B. P. KINGHORN, B. TIER, J. H. J. V. D. WERF, and M. A. CLEVELAND, 2012
 304 A phasing and imputation method for pedigreed populations that results in a single-stage
 305 genomic evaluation. *Genet Sel Evol* **44**: 9.

306 HILL, W. G., and A. ROBERTSON, 1966 The effect of linkage on limits to artificial selection.
 307 *Genet. Res.* **8**: 269–294.

308 HUISMAN, J., 2017 Pedigree reconstruction from SNP data: parentage assignment, sibship
 309 clustering and beyond. *Mol. Ecol. Resour.* **17**: 1009–1024.

310 JOHNSON, M., A. WHALEN, R. ROS-FREIXEDES, G. GORJANC, C.-Y. CHEN, *et al.*, 2020 Genetics
 311 of recombination rate variation in the pig. *bioRxiv* .

312 JOHNSTON, S. E., C. BÉRÉNOIS, J. SLATE, and J. M. PEMBERTON, 2016 Conserved genetic ar-
 313 chitecture underlying individual recombination rate variation in a wild population of soay
 314 sheep (*Ovis aries*). *Genetics* **203**: 583–598.

315 JOHNSTON, S. E., J. HUISMAN, and J. M. PEMBERTON, 2018 A genomic region containing
 316 *REC8* and *RNF212B* is associated with individual recombination rate variation in a wild
 317 population of red deer (*Cervus elaphus*). *BioRxiv* .

318 KADRI, N. K., C. HARLAND, P. FAUX, N. CAMBISANO, L. KARIM, *et al.*, 2016 Coding and
 319 noncoding variants in HFM1, MLH3, MSH4, MSH5, RNF212, and RNF212B affect re-
 320 combination rate in cattle. *Genome research* **26**: 1323–1332.

321 KIJAS, J. W., D. TOWNLEY, B. P. DALRYMPLE, M. P. HEATON, J. F. MADDOX, *et al.*, 2009 A
 322 genome wide survey of SNP variation reveals the genetic structure of sheep breeds. *PLoS*
 323 *ONE* **4**: e4668.

324 KONG, A., J. BARNARD, D. F. GUDBJARTSSON, G. THORLEIFSSON, G. JONSDOTTIR, *et al.*, 2004
 325 Recombination rate and reproductive success in humans. *Nat. Genet.* **36**: 1203–1206.

326 KONG, A., G. THORLEIFSSON, M. L. FRIGGE, G. MASSON, D. F. GUDBJARTSSON, *et al.*, 2014
 327 Common and low-frequency variants associated with genome-wide recombination rate.
 328 Nat. Genet. **46**: 11–6.

329 KONG, A., G. THORLEIFSSON, H. STEFANSSON, G. MASSON, A. HELGASON, *et al.*, 2008 Se-
 330 quence variants in the RNF212 gene associate with genome-wide recombination rate. Sci-
 331 ence **319**: 1398–1401.

332 LENORMAND, T., and J. DUTHEIL, 2005 Recombination difference between sexes: a role for
 333 haploid selection. PLoS biology **3**: e63.

334 MA, L., J. R. O’CONNELL, P. M. VANRADEN, B. SHEN, A. PADHI, *et al.*, 2015 Cattle Sex-Specific
 335 Recombination and Genetic Control from a Large Pedigree Analysis. PLoS Genet. **11**:
 336 e1005387.

337 MULLER, H., 1964 The relation of recombination to mutational advance. Mutat. Res. **1**: 2–9.

338 MYERS, S., L. BOTTOLO, C. FREEMAN, G. McVEAN, and P. DONNELLY, 2005 A fine-scale map
 339 of recombination rates and hotspots across the human genome. Science **310**: 321–4.

340 OTTO, S. P., and N. H. BARTON, 2001 Selection for recombination in small populations.
 341 Evolution **55**: 1921–1931.

342 OTTO, S. P., and T. LENORMAND, 2002 Resolving the paradox of sex and recombination. Nat.
 343 Rev. Genet. **3**: 252–261.

344 OTTO, S. P., and B. A. PAYSEUR, 2019 Crossover Interference: Shedding Light on the Evolu-
 345 tion of Recombination. Annual review of genetics **53**: 19–44.

346 PETIT, M., J.-M. ASTRUC, J. SARRY, L. DROUILHET, S. FABRE, *et al.*, 2017 Variation in Recom-
 347 bination Rate and Its Genetic Determinism in Sheep Populations. Genetics **207**: 767–784.

348 REYNOLDS, A., H. QIAO, Y. YANG, J. K. CHEN, N. JACKSON, *et al.*, 2013 RNF212 is a dosage-
 349 sensitive regulator of crossing-over during mammalian meiosis. Nat. Genet. **45**: 269–78.

350 RÖNNEGÅRD, L., S. E. MCFARLANE, A. HUSBY, T. KAWAKAMI, H. ELLEGREN, *et al.*, 2016 Increas-
 351 ing the power of genome wide association studies in natural populations using repeated
 352 measures - evaluation and implementation. Methods in Ecology and Evolution **7**: 792–799.

353 SAMUK, K., B. MANZANO-WINKLER, K. R. RITZ, and M. A. F. NOOR, 2020 Natural Selection
354 Shapes Variation in Genome-wide Recombination Rate in *Drosophila pseudoob-*
355 *scura*. *Current Biology* **30**: 1517–1528.e6.

356 SANDOR, C., W. LI, W. COPPIETERS, T. DRUET, C. CHARLIER, *et al.*, 2012 Genetic variants in
357 *REC8*, *RNF212*, and *PRDM9* influence male recombination in cattle. *PLoS Genet.* **8**: 1–13.

358 STAPLEY, J., P. G. D. FEULNER, S. E. JOHNSTON, A. W. SANTURE, and C. M. SMADJA, 2017
359 Variation in recombination frequency and distribution across Eukaryotes: patterns and
360 processes. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **372**: 20160455.

361 STOFFEL, M., S. JOHNSTON, J. PILKINGTON, and J. PEMBERTON, 2020 Genetic architecture and
362 lifetime dynamics of inbreeding depression in a wild mammal. *bioRxiv* .

363 WICKHAM, H., 2016 *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

364 YANG, J., S. H. LEE, M. E. GODDARD, and P. M. VISSCHER, 2011 GCTA: a tool for genome-
365 wide complex trait analysis. *Am. J. Hum. Genet.* **88**: 76–82.

366 Supplementary Information




Figure S1: Correlation between linkage map lengths determined in the previous study (JOHNSTON *et al.*, 2016) and in the current study. Each point indicates a chromosome, and lines indicate linear regressions. Points are coloured by sex-averaged or sex-specific map types. The black dashed line indicates a perfect correlation with slope = 1 and intercept = 0.




Figure S2: Sex-specific linkage maps for each sheep chromosome plotted relative to the sheep genome assembly Oar_v3.1. The underlying map data is provided in Table S2. Chromosome 27 is the X chromosome, with the short map segment in male sheep corresponding to the pseudoautosomal region (PAR).




Figure S3: Effect sizes for the most highly associated SNPs shown for males and females, as calculated from animal models including the SNP genotype as a fixed factor. Genotype AA is the model intercept. Numbers indicate the number of observations at each genotype, with the number of unique focal individuals in parentheses.




Figure S4: Regional association statistics for the significant region on chromosome 6. Each point represents a single SNP locus, coloured by their origin on the SNP50 (red) or SNPHD Chips (black, i.e. imputed genotypes). The positions of genes associated with meiotic processes are shown as black bars at the top of each plot (Gene information in Table S4).




Figure S5: Regional association statistics for the significant region on chromosome 7. Each point represents a single SNP locus, coloured by their origin on the SNP50 (red) or SNP50 Chips (black, i.e. imputed genotypes). The positions of genes associated with meiotic processes are shown as black bars at the top of each plot (Gene information in Table S4).

Table S1: Fixed effects in animal models of autosomal crossover count. Models were run in both sexes, males and females. All Wald statistics were equivalent to a χ^2 test with 1 degree of freedom.

Model	Effect	Estimate	SE	Z Ratio	Wald Statistic	P
Both Sexes	(Intercept)	26.966	0.106	253.849	112596.078	0.0000
	Fhat3	1.148	3.692	0.311	0.097	0.7557
	Sex (Male)	7.236	0.180	40.161	1613.634	0.0000
Males	(Intercept)	34.178	0.144	236.710	61239.042	0.0000
	Fhat3	-3.402	5.872	-0.579	0.336	0.5623
Females	(Intercept)	26.970	0.113	238.303	59962.846	0.0000
	Fhat3	0.853	4.610	0.185	0.034	0.8533

Table S2: Linkage map information for the current study. Order is the order of markers on the chromosome. SNP.Name is the Ovine SNP50 BeadChip identifier. Chr is the chromosome number, where 27 is the X chromosome. cMPosition, r, cMdiff are the sex-averaged centiMorgan position, recombination fraction and centiMorgan difference with the following locus, respectively; similarly this is provided for Female and Male maps. Oar3_Chr and Oar3_Pos are the relative chromosome number and position relative to the sheep genome (Oar_v3.1), cMPosition.2016, cMPosition.Female.2016 and cMPosition.Male.2016 indicate the centiMorgan positions as determined by the previous study (JOHNSTON *et al.*, 2016).

[File: Table_S2_Linkage_Map.txt]

Table S3: Association statistics for genome-wide association studies of autosomal chromosome counts. Sex indicates analyses for both sexes, females only and males only. SNP.Name is the Ovine SNP50 BeadChip identifier. Chromosome and Position (bp) are given relative to the sheep genome Oar_v3.1. A1 and A2 reference and alternate allele at each SNP. effB is the slope of the effect of allele A2, with the standard error se_effB. Chi2.1df and P1df is the association chi-squared statistic and associated P-value, respectively, before correction with genomic control λ . Pc1df is the corrected P-value after genomic control. Exp is the corresponding P-value for that SNP locus assuming a null distribution of P-values (see Figure 2). Q.2 is the minor allele frequency. Cumu is the cumulative genomic distance.

[File: Table_S3_ACC_GWAS.txt]

Table S4: Gene positions in the GWAS significant regions obtained using biomaRt v2.42.1. ensembl_gene_id is the Ensembl identifier for the sheep gene. external_gene_name is the sheep gene name. chromosome_name, start_position and end_position are the chromosome number, start and stop positions for each gene, respectively. Meiotic indicates whether this gene and/or its orthologues in other species have GO terms associated with meiotic processes. Orthos indicates the unique gene names associated with orthologues of each gene.

[File: Table_S4_Gene_List_Table.txt]