

Surgery

Susmit

2022-06-21

Contents

Deep Vein Thrombosis (DVT)	1
Virchow's triad	1
Factors	2
Common sites	2
Prevention	2
CABG	2
Investigations for IHD	2
Indications for surgery	2
Graft selection	3
Types	3
Transfusion	3
Indications	3
Complications	3
Single transfusion	3
Massive transfusion	3
Clinical factoids	3
Burns	4
Assessment	4

Deep Vein Thrombosis (DVT)

Formation of semisolid coagulum in a deep vein.

Virchow's triad

- Abnormal surface (endothelial damage)
- Abnormal flow (stasis / turbulence)
- Abnormal blood (thrombophilia)

Factors

- Immobility
 - Age
 - Obesity
 - Prolonged surgery
 - Pregnancy
 - Puerperium
 - Varicosity (effect of immobility, the rest are causes)
- Hormone-replacement therapy (high oestrogen)
- Previous DVT / PE
- Thrombophilia

Common sites

- Popliteal vein
- Femoral "
- Iliac "

Prevention

- Early mobilization
- Hydration
- Compression stockings
- Prophylactic LMW heparin
- Calf pumps
- Minimal use of tourniquets

CABG

Investigations for IHD

- ECG (first line)
- Cardiac enzymes (in acute coronary syndrome)
- Exercise tolerance test
- Echo: Evaluate
 - ventricular function
 - regional wall motion abnormalities
 - valvular lesions
- **Coronary angiography: gold std**
 - Extent, severity and location of stenoses
 - 70% reduction of diameter (i.e. >90% reduction of cross-sec) => severe

Indications for surgery

- 50% stenosis of the left coronary artery ("*left main stem*")
- 50% stenosis of the proximal *LAD*
- 2/3 main coronary arteries diseased (*RCA*, *LAD*, *LCx*)

Graft selection

Types

- **Venous:** long saphenous vein
- **Arterial:**
 - LIMA most common (left internal mammary / left internal thoracic artery)
 - Others
 - * RIMA
 - * Radial
 - * Gastroepiploic
 - * Inf epigastric

Transfusion

Indications

- Acute blood loss
- Periop anaemia
- Symptomatic chronic anaemia

Complications

Single transfusion

- Haemolysis (haemolytic transfusion reaction)
- Fever (febrile transfusion reaction)
- Allergic reaction
- Infections
 - Hep B, C
 - HIV
 - Malaria
 - Bacterial inf
- Air embolism
- Thrombophlebitis
- TRALI

Massive transfusion

- Coagulopathy
- Hypothermia
- Hypo-Ca
- Hypo-K
- Hyper-K

Clinical factoids

- Target Hb level: 10g/dL
- 1 unit transfusion = 1g/dL improvement

Burns

Assessment

- Rule of 9:
 - First approx
 - Adult
 - * Head-neck $\rightarrow 9\%$
 - * Each upper limb $\rightarrow 9\%$
 - * Torso front 18%
 - * Torso back 18%
 - * Each lower limb 18%
 - * Perineum 1%
- Lund and Browder chart
 - More accurate
- For smaller burns, a piece of small paper about the size of the hand to measure the burnt area directly.
Size of hand $\approx 1\%$.