

CONSTRUCTION OF A CLOSED HYPERBOLIC SURFACE OF ARBITRARILY SMALL EIGENVALUE OF PRESCRIBED SERIAL NUMBER

SUSOVAN PAL

ABSTRACT. In this paper we construct, for given any small positive number ϵ and given natural number n , and given any closed hyperbolic surface M , a closed hyperbolic covering surface \tilde{M} , such that its n -th eigenvalue is less than ϵ . An application of this result will also be discussed. The main result follows from the techniques used in B.Randol's paper in 1974 [Ran]. Here I give a new and geometric proof of the main result.

1. INTRODUCTION AND PRELIMINARIES

A closed hyperbolic surface is a compact surface without boundary whose Gaussian (or sectional) curvature is -1 .

One can show that, by considering the Euler characteristic of the surface and using Gauss-Bonnet theorem for closed surface that closed oriented surface can support a complete Riemannian metric with constant Gaussian curvature -1 (called the hyperbolic metric on the surface) if and only if its genus is greater than or equal to 2. In this paper, we will be primarily concerned with hyperbolic metrics.

Laplacian on M is the linear operator acting on the space of all smooth functions on M , defined by $\Delta f = \text{div}(\nabla f)$ and let λ_n be the n -th eigenvalue of the Laplace operator. For a closed oriented surface, it is known that, the spectrum, i.e. the set of all the eigenvalues of the Laplacian is always discrete, countable and the eigenvalues are non-negative[Ch], so we can talk about the n -th eigenvalue of Laplacian operator. Some authors define the Laplacian operator as $\Delta f = -\text{div}(\nabla f)$, so for them the eigenvalues would be non-positive. Eigenvalues of the Laplace operator has been an area of continuous study and research, because of its obvious connection with Physics and other areas in Mathematics. There have been much research on the upper bound on the eigenvalues and also on whether we can produce closed surfaces with small eigenvalues. Good references for the geometry and spectra for closed hyperbolic surfaces are the books by P. Buser[Bu] and I. Chavel[Ch], among others. Another good reference for both compact and non-compact Riemann surfaces is the book by Nicolas Bergeron[Bergeron]. I will generalize a result mentioned in [Bergeron].

Date: April 01 ,2012.

1991 *Mathematics Subject Classification.* 35P15(49G05 53C20)(primary), and 58J50(secondary).

Key words and phrases. Riemann Surfaces, Hyperbolic Geometry, Differential geometry.

The main result, theorem 3.4, follows from the techniques involving Selberg's trace formula used by Prof. Burton Randol's 1974 paper "Small Eigenvalues of the Laplace Operator on Compact Riemann Surfaces" [Ran]. He also mentioned the result in Chapter 11 of Issac Chavel's book "Eigenvalue in Riemannian Geometry". Here I am giving an alternative proof of his result using more geometric techniques and elementary methods.

2. ORGANIZATION OF THE PAPER

In section 3, I will state the the minimax principle (theorem 3.1) used in the context of eigenvalues, and the result mentioned in [Bergeron], the lemma used there to prove that result, which is crucial in the proof of the lemma. Then in the same section, I will state the main theorem 3.4, which is a generalization of the result in [Bergeron], and the lemma 3.5 used to prove the main theorem, which generalizes the previous lemma 3.3, and hinges on the minimax principle. I will give a complete proof of the lemma and the main theorem.

In section 4, I will mention an application of the main theorem to a result of R.Schoen, S. Wolpert and S. T. Yau [SWY], whose another proof is given in [DPRS]. I will state the relevant definitions.

I also take the pleasure to thanks Prof. Feng Luo, my doctoral thesis advisor for suggesting this problem, for various helpful discussions and encouragement in this subject, and my colleague Tian Yang, Dr. Marius Beceanau and Ali Maalaoui for giving me suggestions on typesetting this article.

3. STATEMENT AND PROOF OF THE MAIN THEOREM

We will start with the minimax principle. Let $W^{1,2}(M)$ denote the Sobolev space of functions on M whose first order distributional derivatives exist on M and are (globally) square-integrable on M .

Theorem 3.1 (Minimax principle) :

Let f_1, f_2, \dots, f_{k+1} be continuous functions on M such that they lie in the Sobolev space $W^{1,2}(M)$ and assume that volume of (support of $f_i \cap$ support of f_j) = 0 $\forall 1 \leq j \leq k+1$. Then the k -th eigenvalue λ_k of M satisfies the upper bound:

$$\lambda_k \leq \max_{1 \leq i \leq k+1} \frac{(\|\nabla f_i\|_2)^2}{(\|f_i\|_2)^2}$$

For a proof of Minimax principle, please see Peter Buser's book [Bu] or in Issac Chavel's book [Ch]. Let us remark here that some authors also use the symbol $H^1(M)$ or $W^1(M)$ in stead of $W^{1,2}(M)$ to denote the corresponding Sobolev spaces.

Next, let us state the theorem from [Bergeron] that we intend to generalize.

Theorem 3.2 (Bergeron):

Given any connected, closed, hyperbolic surface M , and given any $\epsilon > 0$, there exists a finite cover \tilde{M} of M such that its 1-st eigenvalue $\lambda_1(\tilde{M}) < \epsilon$

To prove his theorem, [Bergeron] used the following (technical) lemma which we will generalize as well :

Lemma 3.3 :

Let M be a closed hyperbolic surface such that $M = A \cup B$ where A and B are two connected compact sets satisfying $A \cap B = \bigcup_{i=1}^n \gamma_i$, where γ_i 's are simple closed geodesics in M . Let $l(\gamma_i)$ denote the length of γ_i and let :

$$h = \frac{\sum_{i=1}^n l(\gamma_i)}{\text{minimum}\{\text{area}(A), \text{area}(B)\}}$$

Let $\eta > 0$ be a positive number such that η -neighborhood of every γ_i is embedded in M . Then there exists a constant $C(\eta)$, depending only on η such that the first positive eigenvalue $\lambda_1(M)$ satisfies : $\lambda_1(M) \leq C(\eta)(h + h^2)$.

The proof of Theorem 3.2 [Bergeron] follows from lemma 3.3.

Finally, we state the main theorem of this paper :

Theorem 3.4. (Main Theorem):

Given any connected, closed, hyperbolic surface M , given any natural number n ; and given any $\epsilon > 0$; there exists a finite cover \tilde{M} of M such that its n -th eigenvalue $\lambda_n(\tilde{M}) < \epsilon$.

Proof of the main theorem will follow from the following lemma :

Lemma 3.5 :

Let \tilde{M} be a closed hyperbolic surface such that $\tilde{M} = \bigcup_{i=1}^{n+1} A_i$ and $A_1 \cap A_2 = \gamma_1, A_2 \cap A_3 = \gamma_2, \dots, A_n \cap A_{n+1} = \gamma_n, A_{n+1} \cap A_1 = \gamma_{n+1}$, where A_i 's are closed subsets of \tilde{M} and γ_i 's are pairwise disjoint simple closed geodesic in \tilde{M} , and $A_i \cap A_j = \emptyset \forall j \geq i + 2$ except that $A_1 \cap A_{n+1} = \gamma_{n+1}$. Further assume that areas of all A_i and lengths of all the γ_i 's are equal, and that η neighborhood of each γ_i is embedded in \tilde{M} . Then we have : $\lambda_n(\tilde{M}) \leq C(\eta)(h + h^2)$, where $C(\eta)$ is a positive constant depending only on η , and $h = (n + 1) \cdot \frac{l(\gamma_1)}{\text{area}(A_1)}$.

Proof:

We will use the minimax principle [Ch] to prove the lemma. We will produce $(n + 1)$ functions g_1, g_2, \dots, g_{n+1} on \tilde{M} such that $\frac{(\|\nabla g_i\|_2)^2}{(\|g_i\|_2)^2} \leq C(\eta)(h + h^2)$.

Define for t small positive,

$$A_i(t) = \{z \in A_i : \text{dist}(z, \gamma_i) \leq t\}$$

So $A_i(t)$ is a half-collar around the simple closed geodesic γ_i .

Next, define the functions $f_i : \tilde{M} \rightarrow \mathbb{R}$ by :

$$f_i(z) = \begin{cases} \frac{1}{t} \cdot \text{dist}(z, \gamma_i) & \text{if } z \in A_i(t) \\ 1 & \text{if } z \in A_i \setminus A_i(t) \\ 0 & \text{if } z \in \tilde{M} \setminus A_i. \end{cases}$$

Then , $(\|\nabla f_i\|_2)^2 = \frac{1}{t^2} \cdot \text{area}(A_i(t))$

And, $(\|f_i\|_2)^2 \geq \text{area}(A_i \setminus (A_i(t)))$

It is clear that $f_i \in C^0(M) \cap W^{1,2}(M) \forall 1 \leq i \leq n$. Now,


$$\begin{aligned}
& \frac{(\|\nabla f_i\|_2)^2}{(\|f_i\|_2)^2} \\
& \leq \frac{1}{t^2} \cdot \frac{\text{area}(A_i(t))}{\text{area}(A_i) - \text{area}(A_i(t))} \\
& \leq \frac{1}{t^2} \cdot \frac{l(\gamma_i) \cdot \sinh(t)}{\text{area}(A_i) - l(\gamma_i) \cdot \sinh(t)} \\
& \leq \frac{1}{t^2} \cdot \frac{1}{n+1} \cdot \frac{h \cdot \text{area}(A_i) \sinh(t)}{\text{area}(A_i) - \frac{1}{n+1} \cdot h \cdot \text{area}(A_i) \sinh(t)} \\
& \leq \frac{1}{t^2} \cdot \frac{1}{n+1} \cdot \frac{h \cdot \sinh(t)}{1 - \frac{1}{n+1} \cdot h \cdot \sinh(t)} \\
& \leq \frac{h \cdot \sinh(t)}{t^2(1 - \sinh(t))} \\
& \leq \frac{h}{t(1 - \sinh(t))} \\
& \leq \frac{2h}{\eta} \cdot \frac{1}{(1 - \sinh(t))} \\
& \leq \frac{2}{\eta} \cdot h(1 + h) \\
& = C(\eta) \cdot h(1 + h), \text{ where } C(\eta) = \frac{2}{\eta}
\end{aligned}$$

This completes the proof of the lemma 3.5.

Proof of the main theorem 3.4 :

Proof of the main theorem :

As genus of M is ≥ 2 , there exists a simple closed geodesic γ embedded in M such that $M \setminus \gamma$ is connected. Take η positive such that η -neighborhood of γ is embedded in M . Fix η once and for all. Now, for each natural number N , construct a cover \tilde{M} of M of degree $(n+1)N$ in the following way : Take $(n+1)N$ copies of $M \setminus \gamma$ and join them in a cyclical way, i.e. each copy of $M \setminus \gamma$ is joined to two other and different copies of $M \setminus \gamma$. Then there exists $(n+1)$ lifts of γ cutting \tilde{M} into $(n+1)$ pieces A_1, A_2, \dots, A_{n+1} ; each one formed by N fundamental domains for the action for the covering \tilde{M} such that $A_1 \cap A_2 = \gamma_1, A_2 \cap A_3 = \gamma_2, \dots, A_n \cap A_{n+1} = \gamma_n, A_{n+1} \cap A_1 = \gamma_{n+1}$, such that each A_i is a union of N copies of $M \setminus \gamma$, which is a disjoint union except for a set of measure zero. Then for each i , $\text{area}(A_i) = N \cdot \text{area}(M)$. (see the corresponding figure of the $(2+1) \cdot 2 = 6$ -fold covering surface of M for $n = 2, N = 2$ below).



Then, by the previous lemma,

$$\lambda_n(\tilde{M}) \leq C(\eta) \cdot \left[\frac{l(\gamma)}{N \cdot \text{area}(M)} + \left(\frac{l(\gamma)}{N \cdot \text{area}(M)} \right)^2 \right] \rightarrow 0 \text{ as } n \rightarrow \infty$$

This proves the main theorem of the article. \square

4. APPLICATION OF THE MAIN THEOREM

Let us first define some quantities already defined in [SWY],[DPRS].

We define the quantity l_n already defined in [SWY].

Definition 4.1 :

Fix a closed hyperbolic surface M . For fixed n let C stand for any (finite) collection of simple closed geodesics in M such that the complement of the union of the geodesics in C is a disjoint union of $(n + 1)$ components. Let \mathcal{C}_n denote the family of all such C 's. Let $l(C)$ denote the sum of lengths of all the geodesics in C . Let l_n denote the infimum of all $l(C)$ where C varies in \mathcal{C}_n . It is easy to show directly using the definition that, $l_n \leq l_{n+1}$.

In [SWY], [DPRS], it is shown that λ_n is related to a geometric quantity l_n , where l_n is the quantity defined in the definition 4.1. Then the results of the papers above show us that :

Theorem 4.2 (SWY, DPRS) :

With l_n defined as above, we have $C_1(g).l_n \leq \lambda_n \leq C_2(g).l_n \forall 1 \leq n \leq (2g-3)$, where $C_1(g), C_2(g)$ are constants depending on only the genus g of the surface.

For a proof, see [SWY] or [DPRS]. Here we can easily prove as a

Corollary of the main theorem :

In the theorem 3.1 above, we cannot make $C_1(g)$ independent of g .

Proof of corollary: For large N , the n -th eigenvalue of \tilde{M} that we just constructed is arbitrarily close to zero but $l_n(\tilde{M}) \geq l_1(\tilde{M}) \geq l_1(M)$; since image of any family of geodesics that cut \tilde{M} into two pieces cut M into two pieces as well, and the image of any geodesic in the family cutting \tilde{M} has the same length of its image, and two geodesics in \tilde{M} could be identified in M . But $l_1(M)$ is a fixed positive number since M is fixed once and for all. So $C_1(g)$ cannot be made independent of g : note that the genus of the covering surfaces go to infinity as ϵ is made arbitrarily smaller. \square

A way to prove the dependence of $C_2(g)$ on the genus g could be to construct a sequence of hyperbolic surfaces from M with n -th eigenvalues going to ∞ and their l_n being less than or equal to that of M .

REFERENCES

- [1] [Bergeron]Nicholas Bergeron: *Le spectre des surfaces hyperboliques*
- [2] [SWY] R. Schoen, S. Wolpert and S. T. Yau: *Geometric bounds on the low eigenvalues of a compact surface*, Proc. Sympos. Pure Math., Vol. 36, Amer. Math. Soc, Providence, RI, 1980, pp. 279-286.
- [3] [DPRS]J. Dodziuk, T. Pignataro, B. Randol and D. Sullivan: *Estimating small eigenvalues of Riemann surfaces*, Contemporary Mathematics 64 (1987), 93-121.
- [4] [Bu]Peter Buser : *Geometry and Spectra of Compact Riemann Surfaces* , Birkhauser, pp. 213-215.
- [5] [Ch]Issac Chavel : *Eigenvalues in Riemannian Geometry* : Academic Press, Chapter 1, 11.
- [6] [Ran] Burton Randol : *Small Eigenvalues of the Laplace Operator on Compact Riemann Surfaces* , Bulletin of the American Mathematical Society 80(1974)996-1000 .

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NJ 08854
E-mail address: susovan97@gmail.com