

SDG Goal 3

Good health and well-being

SDG Target 3.4

By 2030, reduce by one third premature mortality from non-communicable diseases through prevention and treatment and promote mental health and well-being

SDG Indicator 3.4.2

Suicide mortality rate

Time series

Mortality due to suicide

1. General information on the time series

- Date of national metadata: April 30, 2026
- National data: <https://sdg-indicators.de/3-4-2>
- Definition: The time series shows the number of deaths due to suicide per 100,000 inhabitants.

Classification is based on ICD-10 codes. ICD-10 is the 10th Revision of the International Statistical Classification of Diseases and Related Health Problems. In its German Modification (ICD-10-GM), it serves as the official classification system for coding diagnoses in outpatient and inpatient care in Germany. The following ICD-10 codes are included:

X60-X84: Intentional self-harm
Y87.0: Sequelae of intentional self-harm

- Disaggregation: sex, age group.

2. Comparability with the UN metadata

- Date of UN metadata: December 2024
- UN metadata: <https://unstats.un.org/sdgs/metadata/files/Metadata-03-04-02.pdf>
- The time series is compliant with the UN metadata.

3. Data description

- The number of deaths is taken from the Federal Statistical Office's statistics on causes of death and is based on an analysis of death certificates issued by medical doctors.

For the calculation of population data, the retroactively calculated average population provided by the Federal Statistical Office is used. For 2010, this is based on the 2011 census as well as migration, birth, and death statistics; for the years 2012 to 2021, the calculation is based on the 2022 census.

4. Access to data source

- GBE – Table: Deaths, Mortality figures [Selections: Hierarchy: ICD-10 V01-Y98 (4-Steller); ICD10: X60-X84, Y87.0]:
https://www.gbe-bund.de/gbe/isgbe.fundstellen?p_uid=gast&p_aid=99629820&p_sprache=E&p_thema_id=3900&p_action=TRT

- Average population – GENESIS online 12411-0041:
<https://www-genesis.destatis.de/genesis//online?operation=table&code=12411-0041&bypass=true&levelindex=1&levelid=1639396599054#abreadcrumb>
- Population data based on Census 2022 (only available in German):
https://www.destatis.de/DE/Themen/Gesellschaft-Umwelt/Bevoelkerung/Bevoelkerungsstand/_inhalt.html#sprg233540

5. Metadata on source data

- Quality Report – Causes of death statistics (only available in German):
<https://www.destatis.de/DE/Methoden/Qualitaet/Qualitaetsberichte/Gesundheit/todesursachen.pdf>
- Quality Report – Microcensus (only available in German):
<https://www.destatis.de/DE/Methoden/Qualitaet/Qualitaetsberichte/Bevoelkerung/einfuehrung.html>

6. Timeliness and frequency

- Timeliness: t + 8.5 months
- Frequency: Annual

7. Calculation method

- Unit of measurement: Per 100,000 inhabitants
Calculation:

$$\text{Mortality due to suicide} = \frac{\sum_i \text{Deaths with assigned ICD}_i \text{ [number]}}{\text{Population [number]}} \cdot 100,000$$

With: $i \in \{X60; \dots; X84; Y87.0\}$