

AI701 Assignment 1

Sultan Rizky Hikmawan Madjid (20214196)

Due: September 21, 2021

1. For a random variable X denoting the number of heads among n coin tosses which follows the binomial distribution with parameter θ ,

$$\mathbb{P}(X = x|\theta) = \binom{n}{x} \theta^x (1 - \theta)^{n-x}$$

- (a) Given that θ itself is a random variable following the beta distribution with Probability Density Function (PDF) with parameters $a, b > 0$,

$$f(\theta; a, b) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \theta^{a-1} (1-\theta)^{b-1} \mathbb{1}_{0 \leq \theta \leq 1}$$

for gamma function $\Gamma(t) = \int_0^\infty z^{t-1} e^{-z} dz$, the marginal likelihood of x under this setting can be expressed as

$$\begin{aligned} \mathbb{P}(X = x; a, b) &= \int_0^1 \mathbb{P}(X = x|\theta) f(\theta; a, b) d\theta \\ &= \binom{n}{x} \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \int_0^1 \theta^{x+a-1} (1-\theta)^{n+b-x-1} d\theta \\ &= \binom{n}{x} \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \left(\int_0^1 \frac{\theta^{x+a-1} (1-\theta)^{n+b-x-1}}{x+a} d\theta + \frac{n+b-x-1}{x+a} \int_0^1 \theta^{x+a} (1-\theta)^{n+b-x-2} d\theta \right) \\ &= \binom{n}{x} \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \frac{(n+b-x-1) \cdot (n+b-x-2) \cdot \dots \cdot 1}{(a+b+n-2) \cdot (a+b+n-3) \cdot \dots \cdot (x+a)} \int_0^1 \theta^{x+a} (1-\theta)^{n+b-x-2} d\theta \\ &= \binom{n}{x} \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \frac{(n+b-x-1) \cdot (n+b-x-2) \cdot \dots \cdot 1}{(a+b+n-1) \cdot (a+b+n-2) \cdot \dots \cdot (x+a)} \\ &= \binom{n}{x} \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \frac{\Gamma(n+b-x)\Gamma(x+a)}{\Gamma(a+b+n)} \end{aligned}$$

- (b) Assuming that we toss the coin $n = 10$ times and see $x = 9$ heads, for two prior distributions \mathcal{M}_1 and \mathcal{M}_2 for θ defined as:

$$\mathcal{M}_1 : \mathbb{P}(\theta) = \delta_{1/2}(\theta), \quad \mathcal{M}_2 : f(\theta; a, b) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \theta^{a-1} (1-\theta)^{b-1} \mathbb{1}_{\{0 \leq \theta \leq 1\}}.$$

Fixing $a = b = 1$, we'll have that Bayes factor

$$\begin{aligned} \frac{\mathbb{P}(X = x|\mathcal{M}_1)}{\mathbb{P}(X = x|\mathcal{M}_2)} &= \frac{\int \mathbb{P}(X = x|\theta) \mathbb{P}(\theta|\mathcal{M}_1) d\theta}{\int \mathbb{P}(X = x|\theta) \mathbb{P}(\theta|\mathcal{M}_2) d\theta} \\ &= \frac{\binom{n}{x} (1/2)^x (1-1/2)^{n-x}}{\binom{n}{x} \int_0^1 \theta^x (1-\theta)^{n-x} d\theta} \\ &= \frac{(1/2)^n \Gamma(n+2)}{\Gamma(x+1) \Gamma(n-x+1)} \\ &= \frac{(1/2)^{10} (10+1)!}{9! (10-9)!} = \frac{55}{512} = 0.107421875 \end{aligned}$$

2. For a set E and its collection of subsets \mathcal{A} , defining $\sigma(\mathcal{A})$ as the smallest σ -algebra containing \mathcal{A} or the intersection of all σ -algebras containing \mathcal{A} ,

- (a) Given $E = \{1, 2, 3, 4\}$ and $\mathcal{A} = \{\{1\}, \{2\}\}$, from the definition of σ -algebra itself, given $\mathcal{A} = \{A, B\}$, we have that $\sigma(\mathcal{A}) = \{\emptyset, E, A, B, E \setminus A, E \setminus B, A \cup B, E \setminus (A \cup B)\}$. In other words,

$$\sigma(\mathcal{A}) = \{\emptyset, \{1\}, \{2\}, \{1, 2\}, \{3, 4\}, \{2, 3, 4\}, \{1, 3, 4\}, \{1, 2, 3, 4\}\}$$

- (b) For a set E and its collection of subsets \mathcal{A}, \mathcal{C} ,

1. *Proof.* Given $\mathcal{A} \subset \mathcal{C}$, from the definition of σ -algebra generated by \mathcal{C} , $\sigma(\mathcal{C})$, which is the smallest σ -algebra on E that contains \mathcal{C} , we have that $\mathcal{C} \subset \sigma(\mathcal{C})$. Intuitively, this implies that $\mathcal{A} \subset \mathcal{C} \subset \sigma(\mathcal{C})$, or in short, $\mathcal{A} \subset \sigma(\mathcal{C})$. In other words, $\sigma(\mathcal{C})$ is a σ -algebra which contains the sets in \mathcal{A} . Thus, from the definition of σ -algebra generated by \mathcal{A} , which is the smallest σ -algebra on E that contains \mathcal{A} , we have that $\mathcal{A} \subset \sigma(\mathcal{A}) \subset \sigma(\mathcal{C})$, or in short $\sigma(\mathcal{A}) \subset \sigma(\mathcal{C})$. \square
2. *Proof.* Given $\mathcal{A} \subset \sigma(\mathcal{C})$ and $\mathcal{C} \subset \sigma(\mathcal{A})$, it can be noticed that given $\mathcal{A} \subset \sigma(\mathcal{C})$, from the definition of σ -algebra generated by \mathcal{A} , which is the smallest σ -algebra on E that contains \mathcal{A} , we have that $\mathcal{A} \subset \sigma(\mathcal{A}) \subset \sigma(\mathcal{C})$, or in short, $\sigma(\mathcal{A}) \subset \sigma(\mathcal{C})$. Similarly, given $\mathcal{C} \subset \sigma(\mathcal{A})$, from the definition of σ -algebra generated by \mathcal{C} , which is the smallest σ -algebra on E that contains \mathcal{C} , we have that $\mathcal{C} \subset \sigma(\mathcal{C}) \subset \sigma(\mathcal{A})$, or in short, $\sigma(\mathcal{C}) \subset \sigma(\mathcal{A})$. Since we have $\sigma(\mathcal{C}) \subset \sigma(\mathcal{A})$ and $\sigma(\mathcal{A}) \subset \sigma(\mathcal{C})$, therefore, it implies that $\sigma(\mathcal{A}) = \sigma(\mathcal{C})$. \square
3. *Proof.* Given $\mathcal{A} \subset \mathcal{C} \subset \sigma(\mathcal{A})$, from 2(b)1, we have that given $\mathcal{A} \subset \mathcal{C}$, $\sigma(\mathcal{A}) \subset \sigma(\mathcal{C})$, which further implies that $\mathcal{A} \subset \sigma(\mathcal{A}) \subset \sigma(\mathcal{C})$. Now, from 2(b)2, since we have both $\mathcal{A} \subset \sigma(\mathcal{C})$ and $\mathcal{C} \subset \sigma(\mathcal{A})$, it means that $\sigma(\mathcal{A}) = \sigma(\mathcal{C})$, completing our proof. \square

3. For Moment Generating Function (MGF) of a \mathbb{R} -valued random variable X defined as

$$M_X(t) = \mathbb{E}[e^{tX}]$$

- (a) For a standard normal random variable X with PDF $f_X(x) = \frac{1}{\sqrt{2\pi}}e^{-\frac{x^2}{2}}$, we have that it's Moment Generating Function (MGF) is given by:

$$\begin{aligned} M_X(t) &= \mathbb{E}[e^{tX}] = \int_{-\infty}^{\infty} e^{tx} f_X(x) dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{tx - \frac{x^2}{2}} dx \\ &= e^{\frac{t^2}{2}} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(x-t)^2} dx \\ &= e^{\frac{t^2}{2}} \end{aligned}$$

- (b) Defining X_1, \dots, X_n for $n \in \mathbb{N}$ as i.i.d. \mathbb{R} -valued random variables with $M_{X_1}(t)$ defined as the MGF of X_1 , defining $Y_n := c(X_1 + X_2 + \dots + X_n)$, it's MGF can be mathematically expressed as:

$$\begin{aligned} M_{Y_n}(t) &= \mathbb{E}[e^{tY_n}] = \mathbb{E}[e^{tc(X_1 + X_2 + \dots + X_n)}] \\ &= \prod_{i=1}^n \mathbb{E}[e^{tcX_i}] = (\mathbb{E}[e^{tcX_1}])^n = (\mathbb{E}[e^{ctX_1}])^n \\ &= (M_{X_1}(ct))^n \end{aligned}$$

- (c) *Proof.* For i.i.d. \mathbb{R} -valued random variables X_i with mean μ and variance σ^2 , we first define for $n \in \mathbb{N}$ that

$$Y_i := \frac{X_i - \mu}{\sigma}, \quad Z_n := \frac{1}{\sqrt{n}} \sum_{i=1}^n Y_i.$$

Now, to simplify our calculations, we are going to use random variables Y_i which satisfy $\mathbb{E}[Y_i] = \mathbb{E}\left[\frac{X_i - \mu}{\sigma}\right] = 0$ and $\mathbb{E}[Y_i^2] = \mathbb{E}\left[\frac{(X_i - \mu)^2}{\sigma^2}\right] = \frac{\text{Var}[X_i]}{\sigma^2} = 1$. The MGF of Z_n can be calculated as:

$$\begin{aligned} M_{Z_n}(t) &= \mathbb{E}[e^{tZ_n}] \\ &= \mathbb{E}\left[\exp\left(\frac{t}{\sqrt{n}} \left(\sum_{i=1}^n Y_i\right)\right)\right] \\ &= \prod_{i=1}^n \mathbb{E}[e^{tY_i/\sqrt{n}}] = \left(M_{Y_i}\left(\frac{t}{\sqrt{n}}\right)\right)^n \end{aligned}$$

From Taylor Series expansion, by denoting $o(t^2)$ as a function that satisfies $\lim_{t \rightarrow 0} o(t^2)/t^2 = 0$, it is possible to express $M_{Z_n}(t)$ as

$$\begin{aligned} M_{Z_n}(t) &= \left(M_{Y_i}\left(\frac{t}{\sqrt{n}}\right)\right)^n = \left(M_{Y_i}(0) + \frac{\mathbb{E}[Y_i]t}{\sqrt{n}} + \frac{\mathbb{E}[Y_i^2]t^2}{2n} + o\left(\frac{t^2}{n}\right)\right)^n \\ &= \left(1 + \frac{t^2}{2n} + o\left(\frac{t^2}{n}\right)\right)^n \end{aligned}$$

Now, taking the limit as $n \rightarrow \infty$ and neglecting the higher-order terms represented by $o(\frac{t^2}{n})$ since it converges to 0 faster than the other existing terms, it is possible to deduce that

$$\begin{aligned} \lim_{n \rightarrow \infty} M_{Z_n}(t) &= \lim_{n \rightarrow \infty} \left(1 + \frac{t^2}{2n}\right)^n \\ &= e^{\frac{t^2}{2}} \end{aligned}$$

which is the MGF of a standard normal distribution as derived in [3a](#). Thus, Z_n converges in distribution to a standard normal distribution as $n \rightarrow \infty$. \square

4. For a random variable X with finite mean μ and variance σ^2 and its i.i.d. copies X_1, \dots, X_n , defining estimator for σ :

$$\hat{\sigma}_n^2 := \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2, \quad \bar{X}_n := \frac{1}{n} \sum_{i=1}^n X_i.$$

- (a) *Proof.* From the Weak Law of Large Numbers, it can be noticed that as $n \rightarrow \infty$, we have that $\bar{X}_n := \frac{1}{n} \sum_{i=1}^n X_i \xrightarrow{P} \mathbb{E}[X]$ and similarly, $\frac{1}{n} \sum_{i=1}^n X_i^2 \xrightarrow{P} \mathbb{E}[X^2]$. From the fact that

$$\hat{\sigma}_n^2 := \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}_n^2$$

Using both propositions deduced from the Weak Law of Large Numbers and the equation above, it can be implied that as $n \rightarrow \infty$, $\hat{\sigma}_n^2 \xrightarrow{P} \mathbb{E}[X^2] - (\mathbb{E}[X])^2 = \text{Var}[X] = \sigma^2$ or in short, $\hat{\sigma}_n^2 \xrightarrow{P} \sigma^2$. Thus, $\hat{\sigma}_n^2$ is consistent. \square

- (b) *Proof.* Using the expanded form of $\hat{\sigma}_n^2$ derived in 4a, it is possible to express that

$$\begin{aligned} \mathbb{E}[\hat{\sigma}_n^2] &= \frac{1}{n} \sum_{i=1}^n \mathbb{E}[X_i^2] - \mathbb{E}[\bar{X}_n^2] \\ &= \frac{1}{n} \sum_{i=1}^n \left(\text{Var}[X_i] + (\mathbb{E}[X_i])^2 \right) - \left(\text{Var}[\bar{X}_n] + (\mathbb{E}[\bar{X}_n])^2 \right) \\ &= \sigma^2 + \mu^2 - \frac{\sigma^2}{n} - \mu^2 \\ &= \frac{n-1}{n} \sigma^2 \neq \sigma^2 \end{aligned}$$

Due to the fact that $\mathbb{E}[\hat{\sigma}_n^2] \neq \sigma^2$, $\hat{\sigma}_n^2$ is biased. \square

- (c) In order to make $\hat{\sigma}_n^2$ unbiased, it is possible to redefine the estimator of σ as follows:

$$\hat{\sigma}_n^2 := \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2, \quad \bar{X}_n := \frac{1}{n} \sum_{i=1}^n X_i.$$

By this way, we are going to have an estimator $\hat{\sigma}_n^2$ which is *consistent* from the fact that given $n \rightarrow \infty$, $\hat{\sigma}_n^2 = \frac{n}{n-1} \left(\frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}_n^2 \right) \xrightarrow{P} (1) \cdot \sigma^2$, or in short, $\hat{\sigma}_n^2 \xrightarrow{P} \sigma^2$ and *unbiased*, using the fact that

$$\begin{aligned} \mathbb{E}[\hat{\sigma}_n^2] &= \frac{n}{n-1} \left(\sigma^2 + \mu^2 - \frac{\sigma^2}{n} - \mu^2 \right) \\ &= \sigma^2 \end{aligned}$$

5. *Proof.* For a sequence of \mathbb{R} -valued random variables $(X_n)_{n \geq 1}$ with distribution functions $(F_n)_{n \geq 1}$ and a \mathbb{R} -valued random variable X with distribution F , given $X_n \xrightarrow{d} c$, where c is a constant, from the definition of Convergence in Distribution that is being normally presented by the notation $X_n \xrightarrow{d} X$, where for all $x \in \mathbb{R}$, $\lim_{n \rightarrow \infty} F_n(x) = F(x)$, given $X = c$, we'll have that $F(x) = 0$ if $x < c$ and $F(x) = 1$ if $x \geq c$. Here, $F(x)$ is continuous everywhere except at c . This implies that for any $\epsilon > 0$, we have that

$$\begin{aligned}\lim_{n \rightarrow \infty} F_n(c - \epsilon) &= \lim_{n \rightarrow \infty} [\mathbb{P}(X_n \leq c - \epsilon)] = 0, \\ \lim_{n \rightarrow \infty} F_n\left(c + \frac{\epsilon}{2}\right) &= \lim_{n \rightarrow \infty} \left[\mathbb{P}\left(X_n \leq c + \frac{\epsilon}{2}\right)\right] = 1.\end{aligned}$$

Naturally, we have that $\lim_{n \rightarrow \infty} \mathbb{P}(|X_n - c| > \epsilon) \geq 0$ for every $\epsilon > 0$. However, it can be noticed that

$$\begin{aligned}\lim_{n \rightarrow \infty} \mathbb{P}(|X_n - c| > \epsilon) &= \lim_{n \rightarrow \infty} [\mathbb{P}(X_n < c - \epsilon) + \mathbb{P}(X_n > c + \epsilon)] \\ &= \lim_{n \rightarrow \infty} [\mathbb{P}(X_n < c - \epsilon)] + \lim_{n \rightarrow \infty} [\mathbb{P}(X_n > c + \epsilon)] \\ &= 0 + \lim_{n \rightarrow \infty} [\mathbb{P}(X_n > c + \epsilon)] \\ &\leq \lim_{n \rightarrow \infty} \left[\mathbb{P}\left(X_n > c + \frac{\epsilon}{2}\right)\right] \\ &= 1 - \lim_{n \rightarrow \infty} \left[F_n\left(c + \frac{\epsilon}{2}\right)\right] \\ &= 0\end{aligned}$$

Therefore, from both inequalities, we have that $\lim_{n \rightarrow \infty} \mathbb{P}(|X_n - c| > \epsilon) = 0$ for every $\epsilon > 0$, or equivalently, $X_n \xrightarrow{p} c$. \square