

PREPRINTS

10 papers

An overview of the papers released at github.com/suvrit/preprints. Except where marked, the mathematics in these papers was developed in sustained dialogue with an AI assistant (GPT Pro); none of the results is zero-shot. Claude Code was used extensively for writing, editing, and repository management. Results are grouped by subject; entry numbers do not indicate a ranking. Each summary is followed by the paper's title, linked to its latest timestamped PDF, together with its version and, where one exists, its arXiv identifier.

CONTENTS

Combinatorics	1
Matrix analysis	2
Complex geometry	2
Mathematical physics	2
Real and complex analysis	3



COMBINATORICS

1. McLeod's 1959 conjecture. The paper proves McLeod's conjecture that $(h_{\ell+r}/h_\ell)^{1/r}$ is subadditive for all $\ell \geq 0$ and $r \geq 1$, where h_k is the complete homogeneous symmetric polynomial of degree k . The first step shows, via a closed queueing network, that h_k/h_{k-1} is convex; a tensorization argument then couples neighboring ratios tightly enough to make their geometric mean convex.

A Proof of McLeod's 1959 Conjecture on Complete Homogeneous Symmetric Ratios · v1, 5 October 2026 · arXiv:2610.07302

2. Monotone Turán-ratios of Schur polynomials. The paper proves that, unless all n padded parts of λ are equal, the Turán-ratio $S_{\lambda,i}(x)^2 / (S_{\lambda,i-1}(x)S_{\lambda,i+1}(x))$ is strictly increasing for $x > 1$ and $1 \leq i < n$, where $S_{\lambda,i}(x) = s_\lambda(1^{[n-i]}, x^{[i]})$ is the Schur polynomial in n variables with i arguments equal to x and the rest equal to 1. For one-row shapes this answers a question of Ait-Haddou on complete homogeneous symmetric polynomials; in that case a second proof, through the zeros of Meixner polynomials, gives an explicit positive formula for the logarithmic derivative of the ratio. It conjectures the analogue for Macdonald polynomials.

Monotonicity of Turán-ratios of symmetric polynomials · v1, 7 October 2026

3. A Kotljanskii inequality for permanents. The paper proves that $\text{per}(A_{S \cup T}) \text{per}(A_{S \cap T}) \geq \text{per}(A_S) \text{per}(A_T)$ for all $S, T \subseteq [n]$ when A is an inverse M -matrix that becomes symmetric after a positive diagonal similarity; here A_S is the principal submatrix indexed by S . The proof writes permanents as moments of a complex Gaussian vector and applies Ginibre's correlation inequality. An explicit counterexample shows that symmetry cannot be dropped.

A Kotljanskii inequality for permanents · v1, 2 October 2026 · arXiv:2609.39979

4. Majorization for normalized Jack and Macdonald polynomials. The paper proves that if λ majorizes μ , then the Jack polynomial P_λ in n variables, normalized to equal one at $(1, \dots, 1)$, is at least the normalized P_μ on the positive orthant, for every n , and strictly so when the coordinates are distinct. For the conjectured Macdonald analogue, it proves the rank-two case, the case of a one-row partition against the hook below it for $t \leq q$ in every rank, and, by exact certificates, the full statement in three variables for partitions of size at most eight. It shows that the Harish-Chandra–Itzykson–Zuber mechanism behind the Schur and Jack proofs has no direct finite- q analogue. The full conjecture remains open.

On majorization inequalities for normalized Jack and Macdonald polynomials · v1, 8 October 2026

MATRIX ANALYSIS

5. Lin's singular-value conjecture. The paper proves Lin's conjecture that $\lambda_j((1-t)A + tB) \geq \sigma_j^{1/2}(A^{2(1-t)}B^{2t})$ for all positive semidefinite A, B , all j , and every $t \in [0, 1]$. The case $t = \frac{1}{2}$ is Drury's theorem, which answered a question of Bhatia and Kittaneh. The proof follows Drury's argument but replaces its block-matrix step, which works only at the midpoint, with an eigenvalue count based on an inertia identity and a one-variable inequality.

On Lin's Arithmetic–Geometric Means Conjecture for Singular Values · v1, 8 October 2026

6. Bellman's 1959 claim on Hua–Bellman matrices. The paper shows that Bellman's 1959 strengthening of Hua's results on positive definiteness of the matrices $[\det(I - A_i^* A_j)^{-\alpha}]_{ij}$, for strict contractions A_i , rests on an erroneous proof and, by an explicit example, that it is false. It gives conditions under which $\det(I - A^* B)^{-\alpha}$ is a positive definite function: for complex contractions, the known sharp condition, with the integer exponents below Hua's range reached by an elementary argument; for real symmetric contractions, the part of Bellman's claim that survives. It also introduces a Poincaré-like distance on noncommuting strict contractions.

Positive definite functions of noncommuting contractions, Hua–Bellman matrices, and a new distance metric · v1, 2 October 2026 · arXiv:2112.00056v2 · Not AI-assisted

COMPLEX GEOMETRY

7. Schur forms under decomposable curvature. Griffiths asked in 1969 whether positive curvature of a Hermitian holomorphic vector bundle forces positivity of its characteristic forms; Du's rank-nine example shows that a strictly Griffiths-positive metric can have a top Chern form that is negative on an open set. The paper proves that every Schur form is weakly nonnegative wherever the curvature is decomposable, that is, a sum of a completely positive map and a completely copositive map, with no splitting assumption. The algebraic input is the inequality $\Phi_n(H) \geq 0$ for Finski's double mixed discriminant of every decomposable map H , obtained from a quaternionic Gaussian determinant formula. Beyond the decomposable cone, $\Phi_k \geq 0$ on every square compression of H is equivalent to $\Phi_n(H + S) \geq 0$ for every entanglement-breaking map S ; the class of maps satisfying this condition contains indecomposable maps but is not convex for $n \geq 9$, since Du's map is an average of four of its members.

Weak positivity of Schur forms under decomposable curvature · v1, 8 October 2026

MATHEMATICAL PHYSICS

8. Petz's conjecture on the Kubo–Mori scalar curvature. The paper proves Petz's 1994 conjecture that the scalar curvature of the Kubo–Mori metric on the invertible density matrices increases as the state becomes more mixed. Dittmann (2000) reduced the conjecture to the concavity of an explicit symmetric function h of three positive variables, which he could support only by numerical tests; the paper proves that h is strictly concave. The proof writes $h = -1/(8M)$ with M positive and homogeneous of degree one, and shows that $1/M$ is a Stieltjes function along every line with positive base point and positive direction; Cauchy–Schwarz and homogeneity then give concavity, and the Stieltjes property reduces to an explicit inequality in two real variables, proved by elementary estimates for hyperbolic functions.

A Proof of Petz's Conjecture on the Scalar Curvature of the Kubo–Mori Metric · v1, 9 October 2026

9. The Gibilisco–Isola conjecture near the Kubo–Mori metric. The paper proves the Gibilisco–Isola conjecture that the scalar curvature of the Wigner–Yanase–Dyson metric with parameter p on the invertible density matrices increases under mixing for all p in some interval $(1, 1 + \varepsilon)$, together with the mirror statement for $p \in (1 - \varepsilon, 1)$, with $\varepsilon > 0$ independent of the dimension; the case $p \downarrow 1$ is Petz’s conjecture, result 8. For these p the curvature is a concave function of the eigenvalues. By Dittmann’s formula it suffices that one symmetric function H_p of three positive variables is concave, which the paper proves by showing that $-1/H_p$ is a Stieltjes transform along every line in the positive cone, uniformly in p near 1; the uniformity near the boundary of the eigenvalue domain comes from an exact decomposition of H_p . The value of ε is not explicit.

Wigner–Yanase–Dyson Metrics Have Monotone Curvature · v1, 9 October 2026

REAL AND COMPLEX ANALYSIS

10. Reverse regularity of the square-root theta kernel and Csordas’s conjecture. The paper proves that the Hankel kernel $K(x, y) = \Phi(\sqrt{x + y})$, where Φ is the Jacobi theta kernel in the Fourier representation of Ξ , is strictly reverse-regular through order four, and that order four is sharp. The coefficient minors give ordinary, double, and triple Turán inequalities, and Fourier inversion gives global multilinear integral inequalities for Ξ . The third-order sign gives an independent proof of Csordas’s conjecture that $s'^2 - ss''$ is strictly log-concave for $s(t) = \Phi(\sqrt{t})$. The proof combines exact polynomial estimates on the tail with interval computations.

Sharp order-four sign regularity of a Jacobi theta kernel · v1, 10 October 2026

