

Mixed Model

September 17, 2026

1 Fixed effect and random effect

Fixed effect: categorical variables which already represent the entire population.

Random effect: categorical variables which levels are randomly chosen from a larger population. This is used to correct correlated observations.

Consider there are g groups. Group 1 has n_1 observations $(X_1, X_2, \dots, X_{n_1})$ with a group mean μ_1 , group 2 has n_2 observations $(Y_1, Y_2, \dots, Y_{n_2})$ with a group mean μ_2 , and group g has n_g observations $(Z_1, Z_2, \dots, Z_{n_g})$ with a group mean μ_g . The total number of observations is $N = n_1 + n_2 + \dots + n_g$. The total mean is $\mu_{tot} = \frac{1}{g} \sum_{i=1}^g \mu_i$.

The variation within groups σ_{within}^2 is calculated as

$$\sigma_{\text{within}}^2 = \frac{1}{N} \left[\sum_{k=1}^{n_1} (X_k - \mu_1)^2 + \sum_{l=1}^{n_2} (Y_l - \mu_2)^2 + \dots + \sum_{m=1}^{n_g} (Z_m - \mu_g)^2 \right]$$

The variation among groups σ_{among}^2 is calculated as

$$\sigma_{\text{among}}^2 = \frac{1}{N} \left[n_1(\mu_1 - \mu_{tot})^2 + n_2(\mu_2 - \mu_{tot})^2 + \dots + n_g(\mu_g - \mu_{tot})^2 \right]$$

These equations assign equal amount of variation to each individual.

If group is treated as fixed effect, σ_{among}^2 is assumed infinite, meaning that group means are very different so knowing the mean of one group gives no information about mean of another group. If group is treated as random effect, we borrow information from other groups by estimating σ_{among}^2 . Fixed effects do not borrow information from other groups.