

DEFINITION  
 under Application method, system of linear equation  
 Chapter - V  
 Matrices

### 8) definition of matrix

An arrangement of certain numbers in an array of m rows and n columns, such as:

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{m1} & a_{m2} & a_{m3} & \dots & a_{mn} \end{bmatrix}_{m \times n}$$

is called as matrix of order  $m \times n$

where  $m =$  No. of rows

$n =$  No. of columns.

$a_{mn}$  = suffix  $mn$  represent position of an element

e.g.  $a_{11}$  = 1<sup>st</sup> row & 1<sup>st</sup> column.

The element with same row number and column number i.e.,  $a_{11}, a_{22}, a_{33}, \dots$  are said to be diagonal elements.

### Type of matrices

#### ① Row matrices:-

A matrix having only one row and n columns is called as row matrix.

e.g.  $A = [a_{11} \ a_{12} \ \dots \ a_{1n}]_{1 \times n}$

$$B = [1 \ 2 \ 3 \ 4]_{1 \times 4}$$

#### ② Column matrices:-

A matrix having only one column and m rows is called as column matrix.

$$\text{e.g. } A = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix}_{m \times 1}$$

$$(A^T)_{4 \times 1} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}_{4 \times 1}$$

### ③ zero matrix or null matrix

A matrix containing all zero elements is called a zero matrix

e.g.  $Z_{3 \times 3} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}_{3 \times 3}$

### ④ square matrix

A matrix containing number of rows = Number of columns is known as square matrix.

e.g.  $A_{3 \times 3} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$

### ⑤ transpose of matrix:

A matrix obtained by interchanging rows & columns of matrix is called as transpose

It is denoted by  $A^T$  or  $A'$

e.g.  $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$  then  $A^T = \begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix}$

### ⑥ symmetric matrix:

A square matrix is said to be symmetric if  $A = A^T$

e.g.  $A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & 3 \\ -1 & 3 & 2 \end{bmatrix}$

$A^T = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 1 & 3 \\ -1 & 3 & 2 \end{bmatrix}$

$\therefore A = A^T$

(b) symmetric matrix

A matrix is said to be skew symmetric matrix if

$$A^T = -A$$

e.g.  $A = \begin{bmatrix} 0 & 1 & 2 \\ -1 & 0 & 3 \\ -2 & -3 & 0 \end{bmatrix}$

$$A^T = \begin{bmatrix} 0 & -1 & -2 \\ 1 & 0 & -3 \\ 2 & 3 & 0 \end{bmatrix}$$

$$-A^T = \begin{bmatrix} 0 & 1 & 2 \\ -1 & 0 & 3 \\ -2 & -3 & 0 \end{bmatrix}$$

$$\therefore A = -A^T$$

(c) Diagonal matrix

A square matrix containing all non-diagonal elements as zero then it is called a diagonal matrix

e.g.  $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$

(d) Scalar matrix

If a square matrix has all diagonal elements equal i.e.  $a_{11} = a_{22} = a_{33} \dots$  then it is called scalar matrix

e.g.  $A = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{bmatrix}$

(e) Unit matrix or Identity matrix

A diagonal matrix, where all diagonal elements are unity is called identity matrix

$$I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

## (11) Upper Triangular matrix

It is square matrix in which all the elements below the principal diagonal are zero's

e.g.  $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 4 \end{bmatrix}$  or  $\begin{bmatrix} 0 & 2 & 0 \\ 0 & 0 & 3 \\ 0 & 0 & 0 \end{bmatrix}$

## (12) Lower Triangular matrix

A square matrix in which all the elements above principal diagonal are zeros.

## 8.3 Determinant of matrix

$|A|$  = determinant of A

If  $A = \begin{bmatrix} 1 & -2 & -3 \\ 4 & 5 & 2 \\ 1 & 1 & 2 \end{bmatrix}$

$$|A| = 1 \begin{vmatrix} 5 & 2 \\ 1 & 2 \end{vmatrix} - (-2) \begin{vmatrix} 4 & 2 \\ 1 & 2 \end{vmatrix} + (-3) \begin{vmatrix} 4 & 5 \\ 1 & 1 \end{vmatrix}$$

$$= 1 [10 - 2] + 2 [8 - 2] - 3 [4 - 5]$$

$$= 8 + 2 \times 6 + (-3)(-1)$$

$$= 8 + 12 + 3$$

$$|A| = 23$$

8.4 Minor and co-factor of  $|A|$ 

Let  $|A| = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$

i = row position

j = column position

Then minor of an element  $|A|$  is a determinant obtained by omitting the row and the column in which the element is present.

Minor of element is denoted by  $m_{ij}$

$$\text{Minor of } a_1 = M_{11} = \begin{vmatrix} b_2 & c_2 \\ b_3 & c_3 \end{vmatrix}$$

$$\begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

$$\text{Minor of } b_1 = M_{12} = \begin{vmatrix} a_2 & c_2 \\ a_3 & c_3 \end{vmatrix}$$

$$\begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

$$\text{Minor of } b_2 = M_{22} = \begin{vmatrix} a_1 & c_1 \\ a_3 & c_3 \end{vmatrix}$$

$$\begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

\* Now cofactor of an element is denoted by

$$A_{ij} = (-1)^{i+j} \cdot M_{ij}$$

Ex. cofactor of  $a_1$  from above  $|A|$

$$A_{11} = (-1)^{1+1} \cdot M_{11}$$

$$= (-1)^2 \cdot M_{11}$$

$$A_{11} = M_{11}$$

$$A_{11} = \textcircled{2} \begin{vmatrix} b_2 & c_2 \\ b_3 & c_3 \end{vmatrix} \text{ and so on...}$$

85 Adjoint of a matrix and  $A^{-1}$

If  $A$  is a matrix, then

Step 1: find minor matrix  $j.e M$

Step 2: find co-factor matrix  $b.e C$

Step 3: Take transpose of matrix  $C \rightarrow e C^T$   
then  $\text{adj } A = C^T$

Step 4  $\bar{A}^1 = \frac{1}{|A|} \cdot \text{adj} A$ , where  $|A| \neq 0$

Step 1: find

Ex

$$A = \begin{bmatrix} 1 & 1+2 & 1 \\ 1 & 2 & 3 \\ 1 & 4 & 9 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 4 & 1 \\ 3 & 9 & 1 \\ 1 & 1 & 1 \\ 2 & 4 & 1 \end{bmatrix}$$

Step 1 find minor matrix

$$M_{11} = 18 - 12 = 6(-1)^2$$

$$\begin{bmatrix} 6 & -5 & 1 \\ -6 & 8 & -2 \\ 2 & -3 & 1 \end{bmatrix}$$

$$M_{12} = 9 - 3 = 6(-1)^3$$

$$M_{31} = 3 \cdot 2 = 1(-1)^4$$

$$M_{13} = 4 - 2 = 2(-1)^4$$

$$M_{32} = 9 - 1 = 8(-1)^5$$

$$M_{21} = 9 - 4 = 5(-1)^3$$

$$M_{33} = 2 - 1 = 1(-1)^6$$

$$M_{22} = 9 - 1 = 8(-1)^4$$

Step 2:

find co-factor matrix

$$C = \begin{bmatrix} 6 & -6 & 2 \\ -5 & 8 & -3 \\ 1 & -2 & 1 \end{bmatrix}$$

$$\text{Step 3: } \text{adj} A = C^T = \begin{bmatrix} 6 & -5 & 1 \\ -6 & 8 & -2 \\ 2 & -3 & 1 \end{bmatrix}$$

Step 4

$$|A| = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 4 & 9 \end{vmatrix}$$

$$\therefore |A| = 1(18 - 12) - 1(9 - 3) + 1(4 - 2)$$

$$= 6 - 6 + 2$$

$$|A| = 2$$



$$\therefore \bar{A} = \frac{1}{|A|} \cdot \text{adj. } A$$

$$\bar{A} = \frac{1}{2} \begin{bmatrix} 6 & -5 & 1 \\ -6 & 8 & -2 \\ 2 & 3 & 1 \end{bmatrix}.$$

### operation of matrices

#### ① Equality of matrices

Two matrices A and B are equal, their order is same and their corresponding elements are same  
i.e  $a_{ij} = b_{ij}$

$$\text{i.e } \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

#### ② multiplication of matrix by scalar

$$\text{If } A = \begin{bmatrix} 2 & 3 \\ 4 & 5 \\ 6 & 7 \end{bmatrix} \text{ then } 2A = \begin{bmatrix} 4 & 6 \\ 8 & 10 \\ 12 & 14 \end{bmatrix}$$

#### ③ Addition of matrix

$$A+B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} + \begin{bmatrix} -1 & 2 & 4 \\ -3 & 2 & 9 \\ 11 & -5 & 3 \end{bmatrix}$$

$$A+B = \begin{pmatrix} (1)+(-1) & 2+2 & 3+4 \\ 4+(-3) & 5+2 & 6+9 \\ 7+11 & 8+(-5) & 4+8+3 \end{pmatrix}$$

$$A+B = \begin{bmatrix} 0 & 4 & 7 \\ 1 & 7 & 15 \\ 18 & 3 & 12 \end{bmatrix}$$

## \* multiplication of matrix

If  $A = \begin{bmatrix} -1 & 2 & 9 \\ 3 & 4 & 1 \end{bmatrix}_{2 \times 3}$        $B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}_{3 \times 3}$

then  $A \cdot B = \begin{bmatrix} -1 & 2 & 9 \\ 3 & 4 & 1 \end{bmatrix}_{2 \times 3} \cdot \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}_{3 \times 2}$

$$A \cdot B = \begin{bmatrix} -1+8+63 & -2+10+72 & -3+12+61 \\ 3+14+7 & 6+20+18 & 9+24+9 \end{bmatrix}_{2 \times 3}$$

$$A \cdot B = \begin{bmatrix} 70 & 80 & 90 \\ 26 & 34 & 42 \end{bmatrix}$$

Note:

① In basic algebra  $a \cdot b = 0$

then  $a=0$  or  $b=0$

But in matrices if  $A \cdot B = 0$

that does not mean ~~not~~  $A=0$  or  $B=0$

e.g.  $A = \begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix}$      $B = \begin{bmatrix} 2 & 6 \\ -1 & -3 \end{bmatrix}$

Here  $A \cdot B = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ , but  $A \neq 0$ ,  $B \neq 0$

②  $AB \neq BA$ .

## 87 Elementary transformation of matrix

The following three types transformations performed on any non-zero matrix, are called as elementary transformation.

① The interchange of  $i^{\text{th}}$  and  $j^{\text{th}}$  row / column is denoted by  $R_i \leftrightarrow R_j$  /  $C_i \leftrightarrow C_j$

② The multiplication of each element of  $i^{\text{th}}$  row/column by a non-zero scalar  $k$  is denoted by  $k \cdot R_i$  /  $k \cdot C_i$

③  $R_i + k R_j \quad | \quad C_i + k C_j$

## 88 Rank of matrix

The matrix is said to be of rank  $r$  if there is

- ① At least one minor of order  $r$  which is not equal to 0 &
- ② Every minor of order  $(r+1)$  is equal to 0

e.g.  $A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 0 & 5 & 7 \end{bmatrix}$

Here  $|A| = 0$  but  $\begin{vmatrix} 1 & 2 \\ 2 & 3 \end{vmatrix} = -1 \neq 0$

$\therefore$  Rank cannot be 3

$\therefore \boxed{f(A) = 2}$

## 8.9 Rank By using normal form.

Def<sup>n</sup>: By performing elementary transformation any non-zero matrix A can be reduced to one of the following four forms, called as normal form.

$$\textcircled{1} \begin{bmatrix} I_r \\ 0 \end{bmatrix} \quad \textcircled{2} \begin{bmatrix} I_r & 0 \end{bmatrix} \quad \textcircled{3} \begin{bmatrix} I_r \\ 0 \end{bmatrix} \quad \textcircled{4} \begin{bmatrix} I_r & 0 \end{bmatrix}$$

where r represent rank of matrix

Working Rule for Normal form.

Step 1: Make  $a_{11}=1$

$$\begin{bmatrix} 1 & * & * \\ * & * & * \\ * & * & * \end{bmatrix}$$

Step 2 Get zero below  $a_{11}=1$  by Row transformation and R<sub>1</sub> only

$$\begin{bmatrix} 1 & * & * \\ 0 & * & * \\ 0 & * & * \end{bmatrix}$$

Step 3:

Get zero to the right of  $a_{11}=1$  by column transformation and C<sub>1</sub> only.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & * & * \\ 0 & * & * \end{bmatrix}$$

Step 4: Make  $a_{22}=1$  (without using R<sub>1</sub> and C<sub>1</sub>)

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & * \\ 0 & * & * \end{bmatrix}$$

Step 5: Get zero's below  $a_{22}=1$  by row transformation and R3 only.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & * \\ 0 & 0 & * \end{bmatrix}$$

and so on.

Q. find the rank of following matrix by using normal form.

$$A = \begin{bmatrix} 1 & 1 & -1 & 1 \\ 1 & -1 & 2 & -1 \\ 3 & 1 & 0 & 1 \end{bmatrix}$$

$$\xrightarrow{\quad} A = \begin{bmatrix} 1 & 1 & -1 & 1 \\ 1 & -1 & 2 & -1 \\ 0 & 1 & 0 & 1 \end{bmatrix}$$

$$R_2 = R_2 - R_1 \quad \text{and} \quad R_3 = R_3 - 3R_1$$

$$R_2 = R_2 - R_1$$

$$= 1 - (1) = 0$$

$$= (-1) - (1) = -2$$

$$= 2 - (-1) = 3$$

$$= (-1) - 1 = -2$$

$$R_3 = R_3 - 3R_1$$

$$= 3 - 3(1) = 0$$

$$= 1 - 3(-1) = -2$$

$$= 0 - 3(-1) = 3$$

$$= 1 - 3(1) = -2$$

$$A = \begin{bmatrix} 1 & 1 & -1 & 1 \\ 0 & -2 & 3 & -2 \\ 0 & -2 & 3 & -2 \end{bmatrix}$$

$$C_2 = C_2 - C_1$$

$$C_3 = C_3 + C_1$$

$$C_4 = C_4 - C_1$$

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -2 & 3 & -2 \\ 0 & -2 & 3 & -2 \end{bmatrix}$$

$$C_2 = -\frac{1}{2}C_2$$

$$\therefore A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 3 & -2 \\ 0 & 1 & 8 & -2 \end{bmatrix}$$

$$R_3 = R_3 - R_2$$

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 3 & -2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$C_3 \rightarrow C_3 - 3C_2, C_4 = C_4 + 2C_2$$

$$A = \left[ \begin{array}{cc|cc} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \end{array} \right]$$

$$A = \begin{bmatrix} I_2 & 0 \\ 0 & 0 \end{bmatrix}$$

$\therefore f(A) = 2$  by normal form.

Ex find the rank of following matrix by using Normal form.

$$A = \begin{bmatrix} 1 & 1 & 2 & 3 & 5 \\ 3 & 2 & 7 & 5 & 12 \\ 3 & 3 & 6 & 9 & 15 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 1 & 2 & 3 & 5 \\ 3 & 2 & 7 & 8 & 12 \\ 3 & 3 & 6 & 9 & 15 \end{bmatrix}$$

$$R_2 \rightarrow R_2 - 3R_1 \quad \text{&} \quad R_3 \rightarrow R_3 - 3R_1$$

$$= 3 - 3(1) = 0$$

$$= 2 - 3(1) = -1$$

$$= 9 - 3(0) = 1$$

$$= 5 - 3(3) = -4$$

$$= 12 - 3(5) = -3$$

$$= 3 - 3(1) = 0$$

$$= 3 - 3(1) = 0$$

$$= 6 - 3(2) = 0$$

$$= 9 - 3(3) = 0$$

$$= 15 - 3(5) = 0$$

$$A = \begin{bmatrix} 1 & 1 & 2 & 3 & 5 \\ 0 & -1 & 1 & -4 & -3 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$C_2 = C_2 - C_1$$

$$C_3 = C_3 - 2C_1$$

$$C_4 = C_4 - 3C_1$$

$$C_5 = C_5 - 5C_1$$

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & -4 & -3 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$R_2 \rightarrow -1(R_2)$$

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 4 & 3 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$C_3 \rightarrow C_3 + C_2, \quad C_4 \rightarrow C_4 - 4C_2, \quad C_5 \rightarrow C_5 + 3C_2$$

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$A = \begin{bmatrix} I_2 & 0 \\ 0 & 0 \end{bmatrix}$$

$\therefore f(A) = 2$  by Normal form.

Homework:

i)  $A = \begin{bmatrix} 2 & -3 & 4 & 4 \\ 1 & 1 & 1 & 2 \\ 3 & -2 & 3 & 6 \end{bmatrix}$  Ans 3

ii)  $A = \begin{bmatrix} 2 & -1 & 1 & 3 \\ 2 & 4 & -1 & 0 \\ 4 & 3 & 2 & 1 \end{bmatrix}$  Ans 3

iii)  $A = \begin{bmatrix} 8 & -6 & 4 & -3 & 2 \\ 2 & -4 & 3 & 1 & 0 \\ 0 & 1 & -1 & 3 & 1 \\ 4 & -7 & 4 & -4 & 5 \end{bmatrix}$

8.10 Echelon form (eh-shuh-tawn)

In Echelon form we have to make elements below the diagonals zero. Diagonal may be or may not be equal to 1 unlike in normal form, where diagonal must be equal to 1.

Note: For Echelon form only row transformation is allowed

Matrix A must be reduced to form such as

$$\begin{bmatrix} 8 & 2 & 3 & 4 & 5 \\ 0 & 1 & 2 & 3 & 4 \\ 0 & 0 & -5 & 0 & 9 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Ex find the rank of following matrix by using Echelon form

$$A = \begin{bmatrix} 2 & 3 & -1 & -1 \\ 1 & -1 & -2 & -4 \\ 3 & 1 & 0 & -2 \\ 6 & 8 & 0 & -7 \end{bmatrix}$$

$R_1 \leftarrow R_1$

$$A = \begin{bmatrix} 1 & -1 & -2 & -4 \\ 2 & 3 & 1 & -1 \\ 3 & 1 & 3 & -2 \\ 6 & 3 & 0 & -1 \end{bmatrix}$$

$$R_2 \rightarrow R_2 - 2R_1, R_3 \rightarrow R_3 - 3R_1, R_4 \rightarrow R_4 - 6R_1$$

$$A = \begin{bmatrix} 1 & -1 & -2 & -4 \\ 0 & 5 & 3 & 7 \\ 0 & 4 & 9 & 10 \\ 0 & 3 & 12 & 13 \end{bmatrix}$$

$$R_2 \rightarrow R_2 - R_3$$

$$A = \begin{bmatrix} 1 & -1 & -2 & -4 \\ 0 & 1 & -6 & -3 \\ 0 & 4 & 9 & 10 \\ 0 & 3 & 12 & 13 \end{bmatrix}$$

$$R_3 \rightarrow R_3 - 4R_2, \quad R_4 \rightarrow R_4 - 3R_2$$

$$A = \begin{bmatrix} 1 & -1 & -2 & -4 \\ 0 & 1 & -6 & -3 \\ 0 & 0 & 33 & 22 \\ 0 & 0 & 66 & 44 \end{bmatrix}$$

$$R_4 \rightarrow R_4 - 2R_3$$

$$A = \begin{bmatrix} 1 & -1 & -2 & -4 \\ 0 & 1 & -6 & -3 \\ 0 & 0 & 33 & 22 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

② find the rank of

$$A = \begin{bmatrix} 3 & -2 & 0 & -1 \\ 0 & 2 & 2 & 7 \\ 1 & -2 & -3 & 2 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

$$R_1 \rightarrow R_3$$

$$A = \begin{bmatrix} 1 & -2 & -3 & 2 \\ 0 & 2 & 2 & 7 \\ 3 & -2 & 0 & -1 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

$$R_3 \rightarrow R_3 - 3R_1$$

$$A = \begin{bmatrix} 1 & -2 & -3 & 2 \\ 0 & 2 & 2 & 7 \\ 0 & 4 & 9 & -7 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

$$R_2 \leftrightarrow R_4$$

$$A = \begin{bmatrix} 1 & -2 & -3 & 2 \\ 0 & 1 & 2 & 1 \\ 0 & 4 & 9 & -7 \\ 0 & 2 & 2 & 7 \end{bmatrix}$$

Homework  $A = \begin{bmatrix} 6 & 1 & 3 & 8 \\ 4 & 2 & 6 & -1 \\ 10 & 3 & 9 & 7 \\ 16 & 4 & 12 & 15 \end{bmatrix}$  Ans  $\text{r}(A) = 2$

8.11) finding non-singular matrices  $P$  and  $Q$  such that  $PAQ$  is in Normal form

Q. find non-singular matrix  $P$  and  $Q$  such that

$$A = \begin{bmatrix} 7 & 2 & 3 & 4 \\ 2 & 1 & 4 & 3 \\ 3 & 0 & 5 & -10 \end{bmatrix}$$

is reduced to normal form. Also find rank of  $A$

$$\begin{array}{c} I_{3 \times 3} \quad A_{3 \times 4} \quad I_{4 \times 4} \\ \hline 3 \times 1 \quad 4 \times 4 \\ \hline 3 \times 4 \end{array}$$

for a given matrix A

Total No. of rows = 3

$\therefore$  consider unit matrix  $I_3$

Total No. of columns = 4

$\therefore$  consider unit matrix  $I_4$

We can write  $A_{3 \times 4} = I_3 A I_4$

$\therefore$  we can write  $A_{3 \times 4} = I_3 A I_4$

$$\left[ \begin{matrix} 1 & 2 & 3 & 4 \\ 2 & 1 & 4 & 3 \\ 3 & 0 & 5 & -10 \end{matrix} \right] = \left[ \begin{matrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{matrix} \right] A \left[ \begin{matrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{matrix} \right].$$

Our aim is to reduce matrix A to normal form using both row and column transformation

### Note 1

① Apply ~~row~~ transformation on L.H.S and  $I_3$  (Keep  $I_4$  unchanged)

② Apply column transformation on L.H.S of  $I_4$  (Keep  $I_3$  unchanged)

$$R_2 \rightarrow R_2 - 2R_1, R_3 \rightarrow R_3 - 3R_1$$

$$\left[ \begin{matrix} 1 & 2 & 3 & 4 \\ 0 & -3 & -2 & -5 \\ 0 & -6 & -4 & -22 \end{matrix} \right] = \left[ \begin{matrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ -3 & 0 & 1 \end{matrix} \right] A \left[ \begin{matrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{matrix} \right]$$

$$C_2 = C_2 - 2C_1 \quad ; \quad C_3 = C_3 - 3C_1$$

$$C_4 = C_4 - 4C_1$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -3 & -2 & -5 \\ 0 & -6 & -4 & -22 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix} A \begin{bmatrix} 1 & -2 & -3 & -4 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$C_2 = -\frac{1}{3} C_2$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & -2 & -5 \\ 0 & 2 & -4 & -22 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix} A \begin{bmatrix} 1 & \frac{2}{3} & -3 & -4 \\ 0 & \frac{-1}{3} & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\therefore R_3 = R_3 - 2R_2$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & -2 & -5 \\ 0 & 0 & 0 & -12 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & -2 & 1 \end{bmatrix} A \begin{bmatrix} 1 & \frac{2}{3} & -3 & -4 \\ 0 & -\frac{1}{3} & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$C_3 = C_3 + 2C_2 \text{ & } C_4 = C_4 + 5C_2$$

$$\therefore \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -12 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & -2 & 1 \end{bmatrix} A \begin{bmatrix} 1 & \frac{2}{3} & -\frac{5}{3} & -\frac{2}{3} \\ 0 & -\frac{1}{3} & -\frac{1}{3} & -\frac{5}{3} \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$C_4 = -\frac{1}{12} C_4$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & -2 & 1 \end{bmatrix} A \begin{bmatrix} 1 & \frac{2}{3} & -\frac{5}{3} & \frac{1}{12} \\ 0 & -\frac{1}{3} & -\frac{1}{3} & \frac{5}{12} \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$C_3 \leftrightarrow C_4$$

$$\therefore \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & -2 & 1 \end{bmatrix} A \begin{bmatrix} 1 & \frac{3}{2} & \frac{1}{2} & -\frac{5}{2} \\ 0 & -\frac{1}{2} & \frac{5}{36} & -\frac{2}{3} \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$\therefore f(A) = 3$$

$\rightarrow PAQ$  is normal form

and  $P = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & -2 & 1 \end{bmatrix}$ ,  $Q = \begin{bmatrix} 1 & \frac{3}{2} & \frac{1}{2} & -\frac{5}{2} \\ 0 & -\frac{1}{2} & \frac{5}{36} & -\frac{2}{3} \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$

Note: ①  $P$  and  $Q$  are not unique. They vary depending upon elementary transformation used while writing the process.

However, we can cross check if  $PAQ = I$ , then answer is correct.

② For square matrix  $A$ ,  $PAQ = I \Rightarrow A^{-1} = QP$ .  
where,  $P, Q, A$  are non-singular matrices  
(i.e  $|A| \neq 0$ ,  $|P| \neq 0$ ,  $|Q| \neq 0$ )

## UNIT V

### Chapter 9: System of Linear Algebraic Eq's

(1)

simultaneous Eq's

consider an eq

$$2x + 4$$

This is one eq in one unknown (1)

We solve it, we get

$$x = \frac{1}{2} + 2$$

Now, consider two simultaneous eq's

$$2x + y = 3$$

$$x + 2y = 9$$

These are two eq in two unknowns (2)

We solve these eq to find unknowns x and y by various methods.

(1)

Elimination method

(2)

matrix method.

(1)

Elimination method.

$$2x + y = 3 \quad \text{--- (1)}$$

$$x + 2y = 9 \quad \text{--- (2)} \quad \times 2$$

$$2x + y = 3$$

$$\cancel{-2x + 4y = 18}$$

$$-3y = -15$$

$$\boxed{y = 5}$$

$$x + 2y = 9$$

$$x + 2 \times 5 = 9$$

$$x = 9 - 10$$

$$\boxed{x = -1}$$

Solution of system is  $x = -1, y = 5$

## g.2 Representation of simultaneous equation in matrix form

$$A \underset{\substack{\downarrow \\ \text{coefficient} \\ \text{matrix}}}{X} = \underset{\substack{\downarrow \\ \text{unknown} \\ \text{variable} \\ \text{matrix}}}{B}$$

↓  
constant  
matrix

Ex

$$\begin{aligned} x + y - z &= 2 \\ x + y + 3z &= 9 \\ x + y + z &= 5 \end{aligned}$$

$$AX = B$$

$$\left[ \begin{array}{ccc|c} 2 & 1 & -1 & 2 \\ 1 & 1 & 3 & 9 \\ 1 & 1 & 1 & 5 \end{array} \right]$$

Note while actually solving the equation by matrix method, we only consider coeff. and constant matrix separated by a line called as augmented form.

$$\text{Augmented form} = (A | B)$$

$$(A | B) = \left[ \begin{array}{ccc|c} 2 & 1 & -1 & 2 \\ 1 & 1 & 3 & 9 \\ 1 & 1 & 1 & 5 \end{array} \right]$$

g.3 Type of eq.  
Equation

Non-homogeneous eq'

$$\begin{aligned} x + y + z &= 3 \\ x - y + 2z &= 4 \\ 2x + 3y - z &= 0 \end{aligned}$$


Homogeneous Eq'

$$\begin{aligned} x + y + z &= 0 \\ x - y + 2z &= 0 \\ 2x + 3y - z &= 0 \end{aligned}$$

Note.

- ① For Non-homo-eq<sup>n</sup> B matrix is a non-zero matrix i.e R.H.S must contain at least one non-zero number
- ② For homo-eq<sup>n</sup> B matrix is a zero matrix i.e R.H.S is all zero.

9.4 Flow chart to solve system of Algebraic Eq<sup>n</sup>



Note:

As we wing Echelon form to reduce the matrix, we can only use elementary row transformation.

Solve the following system of equations by matrix method

$$x + y + z = 0$$

$$2x + 3y - z = -5$$

$$x - y + z = 4$$

→ The system of equation in matrix form

$$AX = B$$

$$\begin{bmatrix} 1 & 1 & 1 \\ 2 & 3 & -1 \\ 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ -5 \\ 4 \end{bmatrix}$$

Augmented form,

$$\left[ \begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 2 & 3 & -1 & -5 \\ 1 & -1 & 1 & 4 \end{array} \right]$$

$$R_2 \rightarrow R_2 - 2R_1, R_3 \rightarrow R_3 - R_1$$

$$\left[ \begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 0 & 1 & -3 & -5 \\ 0 & -2 & 0 & 4 \end{array} \right]$$

$$R_3 \rightarrow R_3 + 2R_1$$

$$\left[ \begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 0 & 1 & -3 & -5 \\ 0 & 0 & -6 & -6 \end{array} \right]$$

$$\therefore f(A) = 3$$

$$f(A|B) = 3$$

$$\therefore f(A) = f(A|B) = n = 3$$

∴ system is consistent

but  $n=3$  (number of unknown)  
 $x, y, z$

$$\therefore n=3 \neq 1$$

system processes unique sol?

Q.2 Is the following system of eq consistent? If so find the solution.

$$2x - 3y + 5z = 1$$

$$3x + y - z = 2$$

$$x + 4y - 6z = 1$$

Given system of eq in matrix form

$$AX = B$$

$$\begin{bmatrix} 2 & -3 & 5 \\ 3 & 1 & -1 \\ 1 & 4 & -6 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$$

$$[A|B] = \begin{bmatrix} 2 & -3 & 5 & 1 \\ 3 & 1 & -1 & 2 \\ 1 & 4 & -6 & 1 \end{bmatrix}$$

$$R_3 \leftrightarrow R_1$$

$$[A|B] = \begin{bmatrix} 1 & 4 & -6 & 1 \\ 3 & 1 & -1 & 2 \\ 2 & -3 & 5 & 1 \end{bmatrix}$$

$$R_2 \leftarrow R_2 - 2R_1, R_3 \leftarrow R_3 - 3R_1$$

$$[A|B] = \begin{bmatrix} 1 & 4 & -6 & 1 \\ 0 & -11 & 17 & -1 \\ 0 & -11 & 17 & -1 \end{bmatrix}$$

$$R_3 = R_3 - R_1$$

$$\therefore [A|B] = \begin{bmatrix} 1 & 4 & -6 & 1 \\ 0 & -11 & 17 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\therefore f(A) = 2 \quad \therefore f(A|B) = 2$$

$$\therefore f(A) = f(A|B) = 2 \neq 0$$

System is consistent

but  $n=3$ , (No. of unknowns  $x, y, z$ )

$\therefore$  L.N

$\therefore$  system possesses infinite solution.

$$\text{Let } z=t$$

$\therefore$  By R<sub>2</sub>

$$-11y + 17z = -1$$

$$-11y + 17t = -1$$

$$-11y = -1 - 17t$$

$$y = \frac{-1 - 17t}{11}$$

$\therefore$  By R<sub>1</sub>

$$x + 4y - 6z = 1$$

$$x + 4\left(\frac{-1 - 17t}{11}\right) - 6t = 1$$

$$11x + 4(-1 - 17t) - 66t = 11$$

$$11x + 2t = -4$$

$$11x = 7 - 2t$$

$$x = \frac{7 - 2t}{11} \quad (\text{A})$$

$\therefore$  SoP is

$$x = \frac{7 - 2t}{11}, \quad y = \frac{-1 - 17t}{11}$$

$$z=t$$

Q.3 Examine for consistency of foll. eq<sup>n</sup> and solve them if consistent.

$$2x - 3y + 7z = 5$$

$$3x + y - 3z = 13$$

$$2x + 19y - 47z = 32$$

⇒ The Given eq<sup>n</sup> in matrix form

$$AX = B$$

$$\begin{bmatrix} 2 & -3 & 7 \\ 3 & 1 & -3 \\ 2 & 19 & -47 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5 \\ 13 \\ 32 \end{bmatrix}$$

In Augmented form

$$(A|B) = \left[ \begin{array}{ccc|c} 2 & -3 & 7 & 5 \\ 3 & 1 & -3 & 13 \\ 2 & 19 & -47 & 32 \end{array} \right]$$

$$R_1 = R_1 - R_2$$

$$(A|B) = \left[ \begin{array}{ccc|c} -1 & -4 & 10 & -8 \\ 3 & 1 & -3 & 13 \\ 2 & 19 & -47 & 32 \end{array} \right]$$

$$R_2 = R_2 + 3R_1; R_3 = R_3 + 2R_1$$

$$\therefore (A|B) = \left[ \begin{array}{ccc|c} -1 & -4 & 10 & -8 \\ 0 & -11 & 27 & -11 \\ 0 & 11 & -27 & 16 \end{array} \right]$$

$$R_3 \rightarrow R_3 + R_2$$

$$(A|B) = \left[ \begin{array}{ccc|c} -1 & -4 & 10 & -8 \\ 0 & -11 & 27 & -11 \\ 0 & 0 & 0 & 5 \end{array} \right]$$

$$\therefore \rho(A) = 2, \rho(A|B) = 3$$

$$\rho(A) \neq \rho(A|B)$$

∴ system is inconsistent  
Inconsistent system does not have any soln.

Q.4 Investigate the values of  $\lambda$  and  $\mu$  so that the following eq<sup>n</sup> have

- ① unique sol<sup>n</sup>
- ② infinite solution
- ③ no solution.

$$2x + 3y + 5z = 9$$

$$7x + 3y - 2z = 8$$

$$2x + 3y + \lambda z = \mu$$

⇒ matrix form  $AX = B$

$$\begin{bmatrix} 2 & 3 & 5 \\ 7 & 3 & -2 \\ 2 & 3 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 9 \\ 8 \\ \mu \end{bmatrix}$$

$$(A|B) = \left[ \begin{array}{ccc|c} 2 & 3 & 5 & 9 \\ 7 & 3 & -2 & 8 \\ 2 & 3 & 1 & \mu \end{array} \right]$$

$$R_2 = R_2 - 3R_1$$

$$(A|B) = \left[ \begin{array}{ccc|c} 2 & 3 & 5 & 9 \\ 1 & -6 & -17 & -19 \\ 2 & 3 & 1 & \mu \end{array} \right]$$

$$R_1 \leftrightarrow R_2$$

$$(A|B) = \left[ \begin{array}{ccc|c} 1 & -6 & -17 & -19 \\ 2 & 3 & 5 & 9 \\ 2 & 3 & 1 & \mu \end{array} \right]$$

$$R_2 \rightarrow R_2 - 2R_1, R_3 = R_3 - 2R_1$$

$$(A|B) = \left[ \begin{array}{ccc|c} 1 & -6 & -17 & -19 \\ 0 & 15 & 39 & 47 \\ 0 & 15 & 1+34 & \mu+38 \end{array} \right]$$

$$R_3 = R_3 - R_2$$

$$[A|B] \leftarrow \left[ \begin{array}{ccc|c} 1 & -6 & -14 & -19 \\ 0 & 15 & 39 & 47 \\ 0 & 0 & 1-5 & u-9 \end{array} \right]$$

Example on Homogeneous  
System of eqn by using  
Echelon form

i) unique sol<sup>n</sup>: for unique sol<sup>n</sup>

$$f(A) = f(A|B)$$

and  $r=n$

$\therefore$  Here  $n=3$

$\therefore r$  must be equal to 3  
for that

$1-5 \neq 0$  &  $u-9$  can have  
any value

$\therefore 1 \neq 5$  and  $u$  can have  
any value

ii) Infinite sol<sup>n</sup>:

for infinite sol<sup>n</sup>:

$$f(A) = f(A|B), \text{ if } r < n$$

$\therefore$  Here  $n=3$

$\therefore r$  must be equal to 2

for  $r=2$

$$1-5=0 \Rightarrow 1=5$$

$$-u-9=0 \Rightarrow u=9$$

iii) No sol<sup>n</sup>.

for No sol<sup>n</sup>  $f(A) \neq f(A|B)$

and for that,

$$1-5=0 \quad g-u-g \neq 0$$

$\therefore 1=5, u \neq g$

$$x+2y+3z=0$$

$$2x+3y+z=0$$

$$4x+5y+4z=0$$

$$x+2y-2z=0$$

matrix form

$$AX=B$$

$$\left[ \begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 2 & 3 & 1 & 0 \\ 4 & 5 & 4 & 0 \\ 1 & 1 & -2 & 0 \end{array} \right]$$

$$\left[ \begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 0 & 3 & 1 & 0 \\ 0 & 5 & 4 & 0 \\ 0 & 1 & -2 & 0 \end{array} \right]$$

$$R_2 \leftarrow R_2 - 2R_1$$

$$R_3 \leftarrow R_3 - 4R_1$$

$$R_4 \leftarrow R_4 - R_1$$

$$\left[ \begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 0 & -1 & -5 & 0 \\ 0 & -3 & -8 & 0 \\ 0 & -1 & -5 & 0 \end{array} \right]$$

$$R_3 \leftarrow R_3 - 3R_2, R_4 \leftarrow R_4$$

$$\left[ \begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 0 & -1 & -5 & 0 \\ 0 & 0 & 7 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$f(A)=3 \text{ and } f(A|B)=0$$

$\therefore$  system is consistent

$$r(A) = r(A|B) = 3 = r = n.$$

System possesses unique sol<sup>n</sup>.

$$\therefore \text{By R}_3 \quad 7z = 0$$

$$\therefore z = 0$$

$$\text{By R}_2 \quad -y - 5z = 0$$

$$\Rightarrow y = 0$$

$$\text{By R}_1 \quad x + 2y + 3z = 0$$

$$\Rightarrow x = 0$$

Hence  $x = y = z = 0$  is a trivial sol<sup>n</sup>.

Note:-

① Homogeneous system is always consistent i.e  $r(A) = r(A|B)$

② For homogeneous system, unique sol gives all unknown = 0 i.e  $x = y = z = 0$  is called trivial sol<sup>n</sup>.

③ For homogeneous system, infinite sol<sup>n</sup> gives infinite sol<sup>n</sup> of unknowns ( $x, y, z$ ) in terms of some arbitrary constant  $t$ , is called as non-trivial sol<sup>n</sup>.

Homogeneous system

$Ax = B$  always Consistent

$$m \neq n, m = n > 3$$

$$m = n = 3$$

$$r(A) = r = n$$
  
Trivial sol<sup>n</sup>  
unique sol<sup>n</sup>

$$r(A) = r < n$$
  
Infinite sol<sup>n</sup>

$$|A| \neq 0 \quad r(A) = n$$
  
Trivial sol<sup>n</sup>  
unique sol<sup>n</sup>

$$|A| = 0 \quad r(A) < n$$
  
Infinite sol<sup>n</sup>

9-8

## Linear Dependant and Independant vectors.

Let  $x_1, x_2, x_3, \dots, x_n$  be a system of  $n$  row (or column) matrices of the same order (also called vectors).

If there exist  $n$  scalar  $c_1, c_2, c_3, \dots, c_n$  not all zero such that  $c_1x_1 + c_2x_2 + \dots + c_nx_n = 0$ , then the system is linear dependant.

Whereas, if we get  $c_1, c_2, c_3, \dots, c_n$  all zero, then the system is called linearly independant.

Ex Examine for linear dependence of the following system of vectors. If dependant, find the relation bet' them.  
 $x_1 = (3, 1, -4)$ ,  $x_2 = (2, 2, -3)$ ,  $x_3 = (0, -4, 1)$

→ Given vector  $x_1 = (3, 1, -4)$   
 $x_2 = (2, 2, -3)$   
 $x_3 = (0, -4, 1)$

Now consider the matrix equation

$$c_1x_1 + c_2x_2 + c_3x_3 = 0$$

$$\therefore c_1(3, 1, -4) + c_2(2, 2, -3) + c_3(0, -4, 1) = 0$$

$$\therefore 3c_1 + 2c_2 + 0 \cdot c_3 = 0$$

$$3c_1 + 2c_2 - 4c_3 = 0$$

$$-4c_1 - 3c_2 + c_3 = 0$$

which is homogeneous system

Above equation in matrix form

$$AX = B$$

$$\begin{bmatrix} 3 & 2 & 0 \\ 1 & 2 & -4 \\ -4 & -3 & 1 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

In Augmented form

$$(A|B) = \left[ \begin{array}{ccc|c} 8 & 2 & 0 & 0 \\ 1 & 2 & -4 & 0 \\ -4 & -3 & 1 & 0 \end{array} \right]$$

$$\begin{matrix} R_1 \leftrightarrow R_2 \\ \left[ \begin{array}{ccc|c} 1 & 2 & -4 & 0 \\ 8 & 2 & 0 & 0 \\ -4 & -3 & 1 & 0 \end{array} \right] \end{matrix}$$

$$\begin{matrix} \text{By } R_2 \\ \left[ \begin{array}{ccc|c} 1 & 2 & -4 & 0 \\ 0 & -6 & 8 & 0 \\ -4 & -3 & 1 & 0 \end{array} \right] \\ \boxed{C_2 + 3C_3 = 0} \\ \boxed{C_2 + 3t = 0} \\ \boxed{C_3 = 3t} \end{matrix}$$

$$\begin{matrix} \text{By } R_1 \\ C_1 + 2C_2 - 4C_3 = 0 \\ C_1 + 2(3t) - 4t = 0 \\ \boxed{C_1 = -2t} \end{matrix}$$

$R_2 = R_2 - 3R_1$ , and  $R_4 = R_3 + 4R_1$ . As  $C_1, C_2, C_3 \neq 0$ , gives system of linearly dependent relation:

$$(A|B) = \left[ \begin{array}{ccc|c} 1 & 2 & -4 & 0 \\ 0 & -4 & 12 & 0 \\ 0 & 5 & -15 & 0 \end{array} \right]$$

$$R_2 = \frac{R_2}{4}, \quad R_3 = \frac{R_3}{5}$$

$$(A|B) = \left[ \begin{array}{ccc|c} 1 & 2 & -4 & 0 \\ 0 & -1 & 3 & 0 \\ 0 & 1 & -3 & 0 \end{array} \right]$$

$$R_3 = R_3 + R_2$$

$$(A|B) = \left[ \begin{array}{ccc|c} 1 & 2 & -4 & 0 \\ 0 & -1 & 3 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$\begin{aligned} C_1x_1 + C_2x_2 + C_3x_3 &= 0 \\ -2x_1 + 3x_2 + tx_3 &= 0 \\ t(3x_2 + x_3) &= t(2x_1) \end{aligned}$$

$$3x_2 + x_3 = 2x_1$$

Cross check:

$$3(2, 2, -3) + (0, -4, 1) = 2(3, -1)$$

$$(6, 6, -9) + (0, -4, 1) = (6, 2, -8)$$

$$(6, 2, -8) = (6, 2, -8)$$

$$\therefore r(A) = r(A|B) = r = 2$$

But,  $n=3 \rightarrow$  No. of unknowns  $C_1, C_2, C_3$

$$\therefore r < n$$

$\therefore$  System possesses infinite soln

$$\text{Let } C_3 = t$$

### 9.9 Linear Transformation:

A general linear transformation is represented by

$$y_1 = a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n.$$

$$y_2 = a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n.$$

!

$$y_n = a_{n1}x_1 + a_{n2}x_2 + \dots + a_{nn}x_n.$$

which in matrix form can be written as

$$Y = AX$$

$$\begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

Ex

Express each of the transformation

$$x_1 = 3y_1 + 2y_2, \quad x_2 = -y_1 + 4y_2$$

and  $y_1 = z_1 + 2z_2, \quad y_2 = 3z_1$ , In matrix form  
and find the composite transformation which express  
 $x_1, x_2$  in terms of  $z_1, z_2$

→ The transformation

$$x_1 = 3y_1 + 2y_2$$

$$x_2 = -y_1 + 4y_2$$

In matrix form can be written as

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ -1 & 4 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$$

$$\therefore x = AT \quad \text{--- (1)}$$

and the transformation

$$y_1 = z_1 + 2z_2$$

$$y_2 = 3z_1$$

In matrix form can be written as

$$\begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 3 & 0 \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \end{bmatrix}$$

$$y = Bz \quad \text{--- (2)}$$

From eq (1) & (2)

$$x = A(Bz)$$

$$x = ABz$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ -1 & 4 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 0 \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \end{bmatrix}$$

$$= \begin{bmatrix} 3(1)+2(3) & 3(2)+2(0) \\ (-1)(1)+4(3) & (-1)(2)+4(0) \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 9 & 6 \\ 11 & -2 \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \end{bmatrix}$$

$$x_1 = 9z_1 + 6z_2$$

$$x_2 = 11z_1 - 2z_2$$

is the required transformation.

### \* 9:10 Orthogonal Transformation.

The linear transformation  $y = Ax$  where,

$$y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}, \quad A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix}, \quad x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

is said to be orthogonal if it transforms

$$x_1^2 + x_2^2 + \cdots + x_n^2 \text{ into } y_1^2 + y_2^2 + y_3^2 + \cdots + y_n^2$$

Note 1

- ① A square matrix  $A$  is said to be Orthogonal if  $AA^T = I$
- ② If  $A$  is orthogonal then  $A^T = A^{-1}$

Ex ① show that  $A = \frac{1}{3} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix}$  is orthogonal.

$$\text{Let } A \cdot A^T = \frac{1}{3} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix} \cdot \frac{1}{3} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix}$$

$$\therefore A \cdot A^T = \frac{1}{9} \begin{bmatrix} 9 & 0 & 0 \\ 0 & 9 & 0 \\ 0 & 0 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\therefore \boxed{A \cdot A^T = I}$$

Homework

show that  $A$  is orthogonal  $A = \begin{bmatrix} \cos\theta & 0 & \sin\theta \\ 0 & 1 & 0 \\ -\sin\theta & 0 & \cos\theta \end{bmatrix}$

\* check whether following matrices is orthogonal or not

$$A = \begin{bmatrix} 2 & 2 & 1 \\ -2 & 1 & 2 \\ 1 & -2 & 2 \end{bmatrix}$$

\* If  $A$  is an orthogonal matrix, prove that  $|A| = \pm 1$

$$\Rightarrow A \cdot A^T = I$$

$$\therefore |A| \cdot |A^T| = |I|$$

$$\text{but } |A| = |A^T|$$

$$|A| \cdot |A| = |I|$$

$$|A|^2 = 1$$

$$\boxed{|A| = \pm 1}$$