

Assignment No.1

Suyog Tangade
MD/2020/710

Download all python codes from

<https://github.com/suyogtangade/ASSIGNMENT-01.git>

and latex-tikz codes from

<https://github.com/suyogtangade/ASSIGNMENT-01.git>

1 QUESTION No.16(B) (CBSE/2006/SET-2)

Find the co-ordinates of the point equidistant from three given points $\mathbf{A} \begin{pmatrix} 5 \\ 3 \end{pmatrix}$, $\mathbf{B} \begin{pmatrix} 5 \\ -5 \end{pmatrix}$ and $\mathbf{C} \begin{pmatrix} 1 \\ -5 \end{pmatrix}$

Solution:

Let the point equidistant from \mathbf{A} & \mathbf{B} & \mathbf{C} be

$$\mathbf{P} = \begin{pmatrix} x \\ y \end{pmatrix} \quad (1.0.1)$$

$$\|\mathbf{P} - \mathbf{A}\|^2 = \|\mathbf{P} - \mathbf{B}\|^2 \quad (1.0.2)$$

$$\mathbf{P} = \begin{pmatrix} x \\ y \end{pmatrix} \quad (1.0.3)$$

$$\|\mathbf{P}\|^2 + \|\mathbf{A}\|^2 - 2\mathbf{A}^T \mathbf{P} \quad (1.0.4)$$

$$= \|\mathbf{P}\|^2 + \|\mathbf{B}\|^2 - 2\mathbf{B}^T \mathbf{P} \quad (1.0.5)$$

$$(\mathbf{A} - \mathbf{B})^T \mathbf{P} = \frac{\|\mathbf{A}\|^2 - \|\mathbf{B}\|^2}{2} \quad (1.0.6)$$

$$(\mathbf{B} - \mathbf{C})^T \mathbf{P} = \frac{\|\mathbf{B}\|^2 - \|\mathbf{C}\|^2}{2} \quad (1.0.7)$$

$$\begin{pmatrix} (\mathbf{A} - \mathbf{B})^T \\ (\mathbf{B} - \mathbf{C})^T \end{pmatrix} \mathbf{P} = \frac{1}{2} \begin{pmatrix} \|\mathbf{A}\|^2 - \|\mathbf{B}\|^2 \\ \|\mathbf{B}\|^2 - \|\mathbf{C}\|^2 \end{pmatrix} \quad (1.0.8)$$

$$\mathbf{A} \begin{pmatrix} 5 \\ 3 \end{pmatrix}, \mathbf{B} \begin{pmatrix} 5 \\ -5 \end{pmatrix}, \mathbf{C} \begin{pmatrix} 1 \\ -5 \end{pmatrix} \quad (1.0.9)$$

$$\begin{pmatrix} \left(\begin{pmatrix} 5 \\ 3 \end{pmatrix} - \begin{pmatrix} 5 \\ -5 \end{pmatrix} \right)^T \\ \left(\begin{pmatrix} 5 \\ -5 \end{pmatrix} - \begin{pmatrix} 1 \\ -5 \end{pmatrix} \right)^T \end{pmatrix} \mathbf{P} = \frac{1}{2} \begin{pmatrix} \left\| \begin{pmatrix} 5 \\ 3 \end{pmatrix} \right\|^2 - \left\| \begin{pmatrix} 5 \\ -5 \end{pmatrix} \right\|^2 \\ \left\| \begin{pmatrix} 5 \\ -5 \end{pmatrix} \right\|^2 - \left\| \begin{pmatrix} 1 \\ -5 \end{pmatrix} \right\|^2 \end{pmatrix} \quad (1.0.10)$$

$$\begin{pmatrix} \begin{pmatrix} 5 & 3 \end{pmatrix} - \begin{pmatrix} 5 & -5 \end{pmatrix} \\ \begin{pmatrix} 5 & -5 \end{pmatrix} - \begin{pmatrix} 1 & -5 \end{pmatrix} \end{pmatrix} \mathbf{P} = \quad (1.0.11)$$

$$\frac{1}{2} \begin{pmatrix} (\sqrt{5^2 + 3^2})^2 - (\sqrt{5^2 + (-5)^2})^2 \\ (\sqrt{5^2 + (-5)^2})^2 - (\sqrt{1^2 + (-5)^2})^2 \end{pmatrix} \quad (1.0.12)$$

$$\begin{pmatrix} 0 & 8 \\ 4 & 0 \end{pmatrix} \mathbf{P} = \frac{1}{2} \begin{pmatrix} (\sqrt{25} + \sqrt{9})^2 - (\sqrt{25} + \sqrt{25})^2 \\ (\sqrt{25} + \sqrt{25})^2 - (\sqrt{1} + \sqrt{25})^2 \end{pmatrix} \quad (1.0.13)$$

$$\begin{pmatrix} 0 & 8 \\ 4 & 0 \end{pmatrix} \mathbf{P} = \frac{1}{2} \begin{pmatrix} (\sqrt{34})^2 - (\sqrt{50})^2 \\ (\sqrt{50})^2 - (\sqrt{26})^2 \end{pmatrix} \quad (1.0.14)$$

$$\begin{pmatrix} 0 & 8 \\ 4 & 0 \end{pmatrix} \mathbf{P} = \frac{1}{2} \begin{pmatrix} (\sqrt{-16})^2 \\ (\sqrt{24})^2 \end{pmatrix} \quad (1.0.15)$$

$$\begin{pmatrix} 0 & 8 \\ 4 & 0 \end{pmatrix} \mathbf{P} = \begin{pmatrix} -8 \\ 12 \end{pmatrix} \quad (1.0.16)$$

$$\begin{pmatrix} 0 & 8 \\ 4 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -8 \\ 12 \end{pmatrix} \quad (1.0.17)$$

$$8y = -8 \implies y = -1 \quad (1.0.18)$$

$$4x = 12 \implies x = 3 \quad (1.0.19)$$

co-ordinates for the point equidistant from three




Fig. 1.1: Graphical Solution

given points $\mathbf{A} \begin{pmatrix} 5 \\ 3 \end{pmatrix}$, $\mathbf{B} \begin{pmatrix} 5 \\ -5 \end{pmatrix}$ and $\mathbf{C} \begin{pmatrix} 1 \\ -5 \end{pmatrix}$ are

$$\mathbf{P} = \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 3 \\ -1 \end{pmatrix} \quad (1.0.20)$$