1

Assignment-07

Suyog Tangade MD/2020/710

Download latex-tikz codes from

https://github.com/suyogtangade/Assignment7.git

Download python codes from

https://github.com/suyogtangade/Assignment7.git

Question taken from

https://github.com/gadepall/ncert/blob/main/linalg /optimization/gvv_ncert_opt.pdf question 2.15

1 Question No 2.14

There are two types of fertilisers F_1 and $F_2.F_1$ consists of 10% nitrogen and 6% phosphoric acid and F_2 consists of 5% nitrogen and 10% phosphoric acid. After testing the soil conditions, a farmer finds that she needs atleast 14 kg of nitrogen and 14 kg of phosphoric acid for her crop. If F_1 costs Rs 6/kg and F_2 costs Rs 5/kg, determine how much of each type of fertiliser should be used so that nutrient requirements are met at a minimum cost. What is the minimum cost?

2 Solution

Fertiliser	Nitrogen	Phosphoric acid	Quantity	Cost
F1	10 Percentage	6 Percentage	X Kg	Rs 6
F2	5 Percentage	10 Percentage	Y Kg	Rs 5
Total	14 Kg	14Kg		

Lets consider fertiliser F1 requirement is x kg fertiliser F2 requirement is y kg

$$x \ge 0 \tag{2.0.1}$$

$$y \ge 0 \tag{2.0.2}$$

According to the question,

$$10x + 5y \le 14 \tag{2.0.3}$$

(2.0.4)

and,

$$6x + 10y \le 14 \tag{2.0.5}$$

$$\implies 3x + 5y \le 7 \tag{2.0.6}$$

.. Our problem is

$$\max_{\mathbf{x}} Z = \begin{pmatrix} 6 & 5 \end{pmatrix} \mathbf{x} \tag{2.0.7}$$

$$s.t. \quad \begin{pmatrix} 10 & 5 \\ 3 & 5 \end{pmatrix} \mathbf{x} \le \begin{pmatrix} 14 \\ 7 \end{pmatrix} \tag{2.0.8}$$

Lagrangian function is given by

$$L(\mathbf{x}, \lambda) = (6 \quad 5)\mathbf{x} + \{ [(10 \quad 5)\mathbf{x} + 14] + [(3 \quad 5)\mathbf{x} + 7] + [(-1 \quad 0)\mathbf{x}] + [(0 \quad -1)\mathbf{x}] \} \lambda$$

$$(2.0.9)$$

where,

$$\lambda = \begin{pmatrix} \lambda_1 \\ \lambda_2 \\ \lambda_3 \\ \lambda_4 \\ \lambda_5 \\ \lambda_6 \end{pmatrix} \tag{2.0.10}$$

Now,

$$\nabla L(\mathbf{x}, \lambda) = \begin{pmatrix} 6 + (10 & 5 & -1 & 0) \lambda \\ 6 + (3 & 5 & 0 & -1) \lambda \\ (10 & 5) \mathbf{x} + 14 \\ (3 & 5) \mathbf{x} + 7 \\ (-1 & 0) \mathbf{x} \\ (0 & -1) \mathbf{x} \end{pmatrix}$$
(2.0.11)

:. Lagrangian matrix is given by

$$\begin{pmatrix} 0 & 0 & 10 & 5 & -1 & 0 \\ 0 & 0 & 3 & 5 & 0 & -1 \\ 10 & 5 & 0 & 0 & 0 & 0 \\ 3 & 5 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \mathbf{x} \\ \lambda \end{pmatrix} = \begin{pmatrix} -6 \\ -5 \\ 14 \\ 7 \\ 0 \\ 0 \end{pmatrix} \quad (2.0.12)$$

Considering λ_1, λ_2 as only active multiplier,

$$\begin{pmatrix} 0 & 0 & 10 & 5 \\ 0 & 0 & 3 & 5 \\ 10 & 5 & 0 & 0 \\ 3 & 5 & 0 & 0 \end{pmatrix} \begin{pmatrix} \mathbf{x} \\ \lambda \end{pmatrix} = \begin{pmatrix} -6 \\ -5 \\ 14 \\ 7 \end{pmatrix}$$
 (2.0.13)

resulting in,

$$\implies \begin{pmatrix} \mathbf{x} \\ \lambda \end{pmatrix} = \begin{pmatrix} 0 & 0 & \frac{1}{7} & \frac{-1}{7} \\ 0 & 0 & \frac{-3}{35} & \frac{2}{7} \\ \frac{1}{7} & \frac{-1}{7} & 0 & 0 \\ \frac{-3}{35} & \frac{2}{7} & 0 & 0 \end{pmatrix} \begin{pmatrix} -6 \\ -5 \\ 14 \\ 7 \end{pmatrix}$$
(2.0.15)

$$\implies \begin{pmatrix} \mathbf{x} \\ \lambda \end{pmatrix} = \begin{pmatrix} 1 \\ \frac{4}{5} \\ \frac{-1}{7} \\ \frac{-32}{25} \end{pmatrix} \tag{2.0.16}$$

$$\therefore \lambda = \begin{pmatrix} \frac{-1}{7} \\ \frac{-23}{2\varepsilon} \end{pmatrix} > \mathbf{0}$$

.. Optimal solution is given by

$$\mathbf{x} = \begin{pmatrix} 1 \\ \frac{4}{5} \end{pmatrix} \tag{2.0.17}$$

$$Z = \begin{pmatrix} 6 & 5 \end{pmatrix} \mathbf{x} \tag{2.0.18}$$

$$= \begin{pmatrix} 6 & 5 \end{pmatrix} \begin{pmatrix} 1 \\ \frac{4}{5} \end{pmatrix} \tag{2.0.19}$$

$$= 10$$
 (2.0.20)

Hence the cost will be minimum If fertilizer F1 used x = 1kg and fertilizer F2 used $y = \frac{4}{5}$

Minimum cost Z = 10. This