

# Weak convergence proof v.2 (neater) (in progress)

Suzie Brown

January 20, 2021

**Lemma 1.**

$$t^l - \left( \sum_{s=1}^{\tau_N(t)} c_N(s)^2 \right) \binom{l}{2} (t+1)^{l-2} \leq \sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \prod_{j=1}^l c_N(s_j) \leq t^l + c_N(\tau_N(t))(t+1)^l. \quad (1)$$

*Proof.* As pointed out in Koskela et al. (2018, Equation (8)),

$$\sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \prod_{j=1}^l c_N(s_j) \geq \left( \sum_{s=0}^{\tau_N(t)} c_N(s) \right)^l - \binom{l}{2} \left( \sum_{s=0}^{\tau_N(t)} c_N(s)^2 \right) \left( \sum_{s=0}^{\tau_N(t)} c_N(s) \right)^{l-2}. \quad (2)$$

By definition of  $\tau_N$ ,

$$t \leq \sum_{s=0}^{\tau_N(t)} c_N(s) \leq t+1. \quad (3)$$

Substituting these bounds into the RHS of (2) yields the lower bound.

It is a true fact that

$$\sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \prod_{j=1}^l c_N(s_j) \leq \left( \sum_{s=0}^{\tau_N(t)} c_N(s) \right)^l, \quad (4)$$

as can be seen by considering the multinomial expansion of the RHS. This is further bounded by

$$\left( \sum_{s=0}^{\tau_N(t)} c_N(s) \right)^l \leq \left( \sum_{s=0}^{\tau_N(t)-1} c_N(s) + c_N(\tau_N(t)) \right)^l \leq [t + c_N(\tau_N(t))]^l, \quad (5)$$

again using the definition of  $\tau_N$ . A binomial expansion yields

$$[t + c_N(\tau_N(t))]^l = t^l + \sum_{i=0}^{l-1} \binom{l}{i} t^i c_N(\tau_N(t))^{l-i} = t^l + c_N(\tau_N(t)) \sum_{i=0}^{l-1} \binom{l}{i} t^i c_N(\tau_N(t))^{l-1-i}, \quad (6)$$

then since  $c_N(s) \leq 1$  for all  $s$ ,

$$\sum_{i=0}^{l-1} \binom{l}{i} t^i c_N(\tau_N(t))^{l-1-i} \leq \sum_{i=0}^{l-1} \binom{l}{i} t^i \leq (t+1)^l. \quad (7)$$

Putting this together yields the upper bound. □

**Lemma 2.** *Let  $B$  be a positive constant which may depend on  $n$ .*

$$\sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \prod_{j=1}^l [c_N(s_j) + B D_N(s_j)] \leq \sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \prod_{j=1}^l c_N(s_j) + \left( \sum_{s=1}^{\tau_N(t)} D_N(s) \right) (t+1)^{l-1} (1+B)^l. \quad (8)$$

*Proof.* We start with a binomial expansion:

$$\begin{aligned} \sum_{s_1 \neq \dots \neq s_l} \prod_{j=1}^l [c_N(s_j) + BD_N(s_j)] &= \sum_{s_1 \neq \dots \neq s_l} \sum_{\mathcal{I} \subseteq [l]} B^{l-|\mathcal{I}|} \left( \prod_{i \in \mathcal{I}} c_N(s_i) \right) \left( \prod_{j \notin \mathcal{I}} D_N(s_j) \right) \\ &= \sum_{\mathcal{I} \subseteq [l]} B^{l-|\mathcal{I}|} \sum_{s_1 \neq \dots \neq s_l} \left( \prod_{i \in \mathcal{I}} c_N(s_i) \right) \left( \prod_{j \notin \mathcal{I}} D_N(s_j) \right) \end{aligned} \quad (9)$$

where  $[l] := \{1, \dots, l\}$ . Since the sum is over all permutations of  $r_1, \dots, r_l$ , we may arbitrarily choose an ordering for  $\{1, \dots, l\}$  such that  $\mathcal{I} = \{1, \dots, |\mathcal{I}|\}$ :

$$\sum_{\mathcal{I} \subseteq [l]} B^{l-|\mathcal{I}|} \sum_{s_1 \neq \dots \neq s_l} \left( \prod_{i \in \mathcal{I}} c_N(s_i) \right) \left( \prod_{j \notin \mathcal{I}} D_N(s_j) \right) = \sum_{I=0}^l \binom{l}{I} B^{l-I} \sum_{s_1 \neq \dots \neq s_l} \left( \prod_{i=1}^I c_N(s_i) \right) \left( \prod_{j=I+1}^l D_N(s_j) \right). \quad (10)$$

Separating the term  $I = l$ ,

$$\begin{aligned} \sum_{I=0}^l \binom{l}{I} B^{l-I} \sum_{s_1 \neq \dots \neq s_l} \left( \prod_{i=1}^I c_N(s_i) \right) \left( \prod_{j=I+1}^l D_N(s_j) \right) &= \sum_{s_1 \neq \dots \neq s_l} \prod_{j=1}^l c_N(s_j) \\ &\quad + \sum_{I=0}^{l-1} \binom{l}{I} B^{l-I} \sum_{s_1 \neq \dots \neq s_l} \left( \prod_{i=1}^I c_N(s_i) \right) \left( \prod_{j=I+1}^l D_N(s_j) \right). \end{aligned} \quad (11)$$

In the second line, there is always at least one  $D_N$  term, and  $c_N(s) \leq D_N(s)$  for all  $s$ , so we can write

$$\begin{aligned} \sum_{I=0}^{l-1} \binom{l}{I} B^{l-I} \sum_{s_1 \neq \dots \neq s_l} \left( \prod_{i=1}^I c_N(s_i) \right) \left( \prod_{j=I+1}^l D_N(s_j) \right) &\leq \sum_{I=0}^{l-1} \binom{l}{I} B^{l-I} \sum_{s_1 \neq \dots \neq s_l} \left( \prod_{i=1}^{l-1} c_N(s_i) \right) D_N(s_l) \\ &\leq \sum_{I=0}^{l-1} \binom{l}{I} B^{l-I} \left( \sum_{s_1 \neq \dots \neq s_{l-1}} \prod_{i=1}^{l-1} c_N(s_i) \right) \sum_{s_l=1}^{\tau_N(t)} D_N(s_l) \\ &\leq \sum_{I=0}^{l-1} \binom{l}{I} B^{l-I} (t+1)^{l-1} \sum_{s=1}^{\tau_N(t)} D_N(s) \end{aligned} \quad (12)$$

using (4) and (3). Finally, by the Binomial Theorem,

$$\sum_{I=0}^{l-1} \binom{l}{I} B^{l-I} (t+1)^{l-1} \sum_{s=1}^{\tau_N(t)} D_N(s) \leq \left( \sum_{s=1}^{\tau_N(t)} D_N(s) \right) (t+1)^{l-1} (1+B)^l, \quad (13)$$

which, together with (11), concludes the proof.  $\square$

**Lemma 3.** *Let  $B$  be a positive constant which may depend on  $n$ .*

$$\sum_{s_1 \neq \dots \neq s_l} \prod_{j=1}^l [c_N(s_j) - BD_N(s_j)] \geq \sum_{s_1 \neq \dots \neq s_l} \prod_{j=1}^l c_N(s_j) - \left( \sum_{s=1}^{\tau_N(t)} D_N(s) \right) (t+1)^{l-1} (1+B)^l. \quad (14)$$

*Proof.* A binomial expansion and subsequent manipulation as in (9)–(11) gives

$$\begin{aligned}
\sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \prod_{j=1}^l [c_N(s_j) - BD_N(s_j)] &= \sum_{\mathcal{I} \subseteq [l]} (-B)^{l-|\mathcal{I}|} \sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \left( \prod_{i \in \mathcal{I}} c_N(s_i) \right) \left( \prod_{j \notin \mathcal{I}} D_N(s_j) \right) \\
&= \sum_{I=0}^l \binom{l}{I} (-B)^{l-I} \sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \left( \prod_{i=1}^I c_N(s_i) \right) \left( \prod_{j=I+1}^l D_N(s_j) \right) \\
&= \sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \prod_{j=1}^l c_N(s_j) + \sum_{I=0}^{l-1} \binom{l}{I} (-B)^{l-I} \sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \left( \prod_{i=1}^I c_N(s_i) \right) \left( \prod_{j=I+1}^l D_N(s_j) \right) \\
&\geq \sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \prod_{j=1}^l c_N(s_j) - \sum_{I=0}^{l-1} \binom{l}{I} B^{l-I} \sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \left( \prod_{i=1}^I c_N(s_i) \right) \left( \prod_{j=I+1}^l D_N(s_j) \right)
\end{aligned} \tag{15}$$

where the last inequality just multiplies some positive terms by  $-1$ . Then (12)–(13) can be applied directly (noting that an upper bound on negative terms gives a lower bound overall):

$$-\sum_{I=0}^{l-1} \binom{l}{I} B^{l-I} \sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \left( \prod_{i=1}^I c_N(s_i) \right) \left( \prod_{j=I+1}^l D_N(s_j) \right) \geq \left( \sum_{s=1}^{\tau_N(t)} D_N(s) \right) (t+1)^{l-1} (1+B)^l \tag{16}$$

which concludes the proof.  $\square$

**Lemma 4** (Basis step). *For any  $0 < t < \infty$ ,*

$$\lim_{N \rightarrow \infty} \mathbb{E} \left[ \prod_{r=1}^{\tau_N(t)} (1 - p_r) \right] = e^{-\alpha_n t} \tag{17}$$

where  $\alpha_n := n(n-1)/2$ .

*Proof.* We start by showing that  $\lim_{N \rightarrow \infty} \mathbb{E} \left[ \prod_{r=1}^{\tau_N(t)} (1 - p_r) \right] \leq e^{-\alpha_n t}$ .

From Koskela et al. (2018, Lemma 1 Case 1), taking  $\xi = \Delta$ , we have

$$1 - p_t = p_{\Delta\Delta}(t) \leq 1 - \alpha_n(1 + O(N^{-1})) [c_N(t) - B'_n D_N(t)] \tag{18}$$

where the  $O(N^{-1})$  term does not depend on  $t$ . Applying a multinomial expansion and then separating the positive and negative terms,

$$\begin{aligned}
\prod_{r=1}^{\tau_N(t)} (1 - p_r) &\leq 1 + \sum_{l=1}^{\tau_N(t)} (-\alpha_n)^l (1 + O(N^{-1})) \frac{1}{l!} \sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \prod_{j=1}^l [c_N(s_j) - B'_n D_N(s_j)] \\
&= 1 + \sum_{\substack{l=2 \\ \text{even}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} \sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \prod_{j=1}^l [c_N(s_j) - B'_n D_N(s_j)] \\
&\quad - \sum_{\substack{l=1 \\ \text{odd}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} \sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \prod_{j=1}^l [c_N(s_j) - B'_n D_N(s_j)].
\end{aligned} \tag{19}$$

This is further bounded by applying Lemma 3 and then both bounds of Lemma 1:

$$\begin{aligned}
\prod_{r=1}^{\tau_N(t)} (1 - p_r) &\leq 1 + \sum_{\substack{l=2 \\ \text{even}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} \sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \prod_{j=1}^l c_N(s_j) \\
&\quad - \sum_{\substack{l=1 \\ \text{odd}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} \left\{ \sum_{s_1 \neq \dots \neq s_l}^{\tau_N(t)} \prod_{j=1}^l c_N(s_j) - \left( \sum_{s=1}^{\tau_N(t)} D_N(s) \right) (t+1)^{l-1} (1 + B'_n)^l \right\} \\
&\leq 1 + \sum_{\substack{l=2 \\ \text{even}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} \{ t^l + c_N(\tau_N(t)) (t+1)^l \} \\
&\quad - \sum_{\substack{l=1 \\ \text{odd}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} \left\{ t^l - \left( \sum_{s=1}^{\tau_N(t)} c_N(s)^2 \right) \binom{l}{2} (t+1)^{l-2} - \left( \sum_{s=1}^{\tau_N(t)} D_N(s) \right) (t+1)^{l-1} (1 + B'_n)^l \right\}.
\end{aligned} \tag{20}$$

A bit of tidying up and we have

$$\begin{aligned}
\prod_{r=1}^{\tau_N(t)} (1 - p_r) &\leq \sum_{l=0}^{\tau_N(t)} (-\alpha_n)^l (1 + O(N^{-1})) \frac{1}{l!} t^l + c_N(\tau_N(t)) \sum_{\substack{l=2 \\ \text{even}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} (t+1)^l \\
&\quad + \left( \sum_{s=1}^{\tau_N(t)} c_N(s)^2 \right) \sum_{\substack{l=1 \\ \text{odd}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} \binom{l}{2} (t+1)^{l-2} \\
&\quad + \left( \sum_{s=1}^{\tau_N(t)} D_N(s) \right) \sum_{\substack{l=1 \\ \text{odd}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} (t+1)^{l-1} (1 + B'_n)^l \\
&\leq \sum_{l=0}^{\tau_N(t)} (-\alpha_n)^l (1 + O(N^{-1})) \frac{1}{l!} t^l + c_N(\tau_N(t)) \exp[\alpha_n (1 + O(N^{-1})) (t+1)] \\
&\quad + \left( \sum_{s=1}^{\tau_N(t)} c_N(s)^2 \right) \frac{1}{2} \alpha_n^2 \exp[\alpha_n (1 + O(N^{-1})) (t+1)] \\
&\quad + \left( \sum_{s=1}^{\tau_N(t)} D_N(s) \right) \exp[\alpha_n (1 + O(N^{-1})) (t+1) (1 + B'_n)].
\end{aligned} \tag{21}$$

Now, taking the expectation and limit, and applying Brown et al. (2021, Equations (3.3)–(3.5)) and Lemma ?? (Lemma 2 in the messy weakconv note; not written up here yet),

$$\begin{aligned}
\lim_{N \rightarrow \infty} \mathbb{E} \left[ \prod_{r=1}^{\tau_N(t)} (1 - p_r) \right] &\leq \sum_{l=0}^{\infty} (-\alpha_n)^l \frac{1}{l!} t^l \lim_{N \rightarrow \infty} \mathbb{P}[\tau_N(t) \geq l] + \lim_{N \rightarrow \infty} \mathbb{E}[c_N(\tau_N(t))] \exp[\alpha_n (t+1)] \\
&\quad + \lim_{N \rightarrow \infty} \mathbb{E} \left[ \sum_{s=1}^{\tau_N(t)} c_N(s)^2 \right] \frac{1}{2} \alpha_n^2 \exp[\alpha_n (t+1)] \\
&\quad + \lim_{N \rightarrow \infty} \mathbb{E} \left[ \sum_{s=1}^{\tau_N(t)} D_N(s) \right] \exp[\alpha_n (t+1) (1 + B'_n)] \\
&= \sum_{l=0}^{\infty} (-\alpha_n)^l \frac{1}{l!} t^l = e^{-\alpha_n t}.
\end{aligned} \tag{22}$$

It remains to show that  $\lim_{N \rightarrow \infty} \mathbb{E} \left[ \prod_{r=1}^{\tau_N(t)} (1 - p_r) \right] \geq e^{-\alpha_n t}$ .  
 From Brown et al. (2021, Equation (3.14)), taking  $\xi = \Delta$ , we have

$$1 - p_t = p_{\Delta\Delta}(t) \geq 1 - \alpha_n(1 + O(N^{-1})) [c_N(t) + B_n D_N(t)] \quad (23)$$

where  $B_n > 0$  and the  $O(N^{-1})$  term does not depend on  $t$ . In particular,

$$1 - p_t = p_{\Delta\Delta}(t) \geq 1 - \frac{N^{n-2}}{(N-2)_{n-2}} \alpha_n c_N(t) - \frac{N^{n-3}}{(N-3)_{n-3}} B_n D_N(t). \quad (24)$$

Since  $D_N(s) \leq c_N(s)$  for all  $s$  (Koskela et al., 2018, p.9), a sufficient condition for this bound to be non-negative is

$$E_r := \left\{ c_N(r) \leq \frac{(N-3)_{n-3}}{N^{n-3}} \left( \alpha_n \left( 1 + \frac{2}{N-2} \right) + B_n \right)^{-1} \right\}, \quad (25)$$

and we define  $E := \bigcap_{r=1}^{\tau_N(t)} E_r$ . We now apply a multinomial expansion to the product, and split into positive and negative terms:

$$\begin{aligned} \prod_{r=1}^{\tau_N(t)} (1 - p_r) &\geq \left\{ 1 + \sum_{l=1}^{\tau_N(t)} (-\alpha_n)^l (1 + O(N^{-1})) \frac{1}{l!} \sum_{s_1 \neq \dots \neq s_l} \prod_{j=1}^l [c_N(s_j) + B_n D_N(s_j)] \right\} \mathbb{1}_E \\ &= \left\{ 1 + \sum_{\substack{l=2 \\ \text{even}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} \sum_{s_1 \neq \dots \neq s_l} \prod_{j=1}^l [c_N(s_j) + B_n D_N(s_j)] \right. \\ &\quad \left. - \sum_{\substack{l=1 \\ \text{odd}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} \sum_{s_1 \neq \dots \neq s_l} \prod_{j=1}^l [c_N(s_j) + B_n D_N(s_j)] \right\} \mathbb{1}_E \end{aligned} \quad (26)$$

This is further bounded by applying Lemma 2 and both bounds in Lemma 1:

$$\begin{aligned} \prod_{r=1}^{\tau_N(t)} (1 - p_r) &\geq \left\{ 1 + \sum_{\substack{l=2 \\ \text{even}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} \sum_{s_1 \neq \dots \neq s_l} \prod_{j=1}^l c_N(s_j) \right. \\ &\quad \left. - \sum_{\substack{l=1 \\ \text{odd}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} \left[ \sum_{s_1 \neq \dots \neq s_l} \prod_{j=1}^l c_N(s_j) + \left( \sum_{s=1}^{\tau_N(t)} D_N(s) \right) (t+1)^{l-1} (1 + B_n)^l \right] \right\} \mathbb{1}_E \\ &\geq \left\{ 1 + \sum_{\substack{l=2 \\ \text{even}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} \left[ t^l - \left( \sum_{s=1}^{\tau_N(t)} c_N(s)^2 \right) \binom{l}{2} (t+1)^{l-2} \right] \right. \\ &\quad \left. - \sum_{\substack{l=1 \\ \text{odd}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} \left[ t^l + c_N(\tau_N(t)) (t+1)^l + \left( \sum_{s=1}^{\tau_N(t)} D_N(s) \right) (t+1)^{l-1} (1 + B_n)^l \right] \right\} \mathbb{1}_E. \end{aligned} \quad (27)$$

Tidying things up,

$$\begin{aligned}
\prod_{r=1}^{\tau_N(t)} (1 - p_r) &\geq \sum_{l=0}^{\tau_N(t)} (-\alpha_n)^l (1 + O(N^{-1})) \frac{1}{l!} t^l \mathbb{1}_E - \left( \sum_{s=1}^{\tau_N(t)} c_N(s)^2 \right) \sum_{\substack{l=2 \\ \text{even}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} \binom{l}{2} (t+1)^{l-2} \\
&\quad - c_N(\tau_N(t)) \sum_{\substack{l=1 \\ \text{odd}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} (t+1)^l \\
&\quad - \left( \sum_{s=1}^{\tau_N(t)} D_N(s) \right) \sum_{\substack{l=1 \\ \text{odd}}}^{\tau_N(t)} \alpha_n^l (1 + O(N^{-1})) \frac{1}{l!} (t+1)^{l-1} (1 + B_n)^l \\
&\geq \sum_{l=0}^{\tau_N(t)} (-\alpha_n)^l (1 + O(N^{-1})) \frac{1}{l!} t^l \mathbb{1}_E - \left( \sum_{s=1}^{\tau_N(t)} c_N(s)^2 \right) \frac{1}{2} \alpha_n^2 \exp[\alpha_n (1 + O(N^{-1})) (t+1)] \\
&\quad - c_N(\tau_N(t)) \exp[\alpha_n (1 + O(N^{-1})) (t+1)] \\
&\quad - \left( \sum_{s=1}^{\tau_N(t)} D_N(s) \right) \exp[\alpha_n (1 + O(N^{-1})) (t+1) (1 + B_n)]. \tag{28}
\end{aligned}$$

Now, taking the expectation and limit, and applying Brown et al. (2021, Equations (3.3)–(3.5)) and Lemma ?? (Lemma 2 in the messy weakconv note; not written up here yet),

$$\begin{aligned}
\lim_{N \rightarrow \infty} \mathbb{E} \left[ \prod_{r=1}^{\tau_N(t)} (1 - p_r) \right] &\geq \sum_{l=0}^{\infty} (-\alpha_n)^l (1 + O(N^{-1})) \frac{1}{l!} t^l \lim_{N \rightarrow \infty} \mathbb{P}[\{\tau_N(t) \geq l\} \cap E] \\
&\quad - \lim_{N \rightarrow \infty} \mathbb{E} \left[ \sum_{s=1}^{\tau_N(t)} c_N(s)^2 \right] \frac{1}{2} \alpha_n^2 \exp[\alpha_n (t+1)] \\
&\quad - \lim_{N \rightarrow \infty} \mathbb{E}[c_N(\tau_N(t))] \exp[\alpha_n (t+1)] \\
&\quad - \lim_{N \rightarrow \infty} \mathbb{E} \left[ \sum_{s=1}^{\tau_N(t)} D_N(s) \right] \exp[\alpha_n (t+1) (1 + B_n)] \\
&= \sum_{l=0}^{\infty} (-\alpha_n)^l \frac{1}{l!} t^l = e^{-\alpha_n t}. \tag{29}
\end{aligned}$$

Combining the upper and lower bounds in (22) and (29) respectively concludes the proof.  $\square$

## References

- Brown, S., Jenkins, P. A., Johansen, A. M. and Koskela, J. (2021), ‘Simple conditions for convergence of sequential Monte Carlo genealogies with applications’, *Electronic Journal of Probability* **26**(1), 1–22.
- Koskela, J., Jenkins, P. A., Johansen, A. M. and Spanò, D. (2018), Asymptotic genealogies of interacting particle systems with an application to sequential Monte Carlo, Mathematics e-print 1804.01811, ArXiv.